



Study of the tribe Heresiarchini (Hymenoptera: Ichneumonidae, Ichneumoninae) in northern Iran, with a new record for the Middle East

Farid Shirzadegan¹, Ali Asghar Talebi^{1*}, Matthias Riedel² and Hamidreza Hajiqanbar¹

¹ Department of Entomology, Faculty of Agriculture, Tarbiat Modares University, P. O. Box: 14115-336, Tehran, Iran.

² Zoologische Staatssammlung München, Münchenhausenstr. 21, D-81247 München, Germany.

Received:
06 February 2018

Accepted:
23 June 2018

Published:
25 September 2018

Subject Editor:
Samira Farahani

ABSTRACT. In this survey the tribe Heresiarchini (Hymenoptera: Ichneumonidae, Ichneumoninae) of Alborz and Guilan provinces (Iran) was taxonomically studied. The specimens were collected using Malaise traps during March to November 2010. Three species including: *Coelichneumon comitator* (Linnaeus, 1758), *Coelichneumon probator* Horstmann, 2000 and *Syspasis carinator* (Fabricius, 1798) were identified of which *C. probator* is recorded for the first time from the Middle East and Iran. An updated checklist of Heresiarchini in Iran, as well as diagnostic characters of the new recorded species is presented.

Key words: Ichneumonidae, Ichneumoninae, Heresiarchini, *Coelichneumon*, Iran

Citation: Shirzadegan, F., Talebi, A.A., Riedel, M. & Hajiqanbar, H. (2018) Study of the tribe Heresiarchini (Hymenoptera: Ichneumonidae, Ichneumoninae) in northern Iran, with a new record for the Middle East. *Journal of Insect Biodiversity and Systematics*, 4 (2), 113–122.

Introduction

The Ichneumonidae Latreille, 1802 with 25292 described species is one of the largest families in the insects order (Aguilar et al., 2013). Most ichneumonids are parasitoids of Lepidoptera, Coleoptera, Hymenoptera, Diptera and Arachnida. Thus ichneumonid wasps have important role in decrease of pest population and damages. The Ichneumoninae with 437 genera and about 4355 species is the largest subfamily (after Cryptinae) in the Ichneumonidae family, and has a worldwide distribution (Çoruh & Özbek, 2013; Yu et al., 2016). Almost all known species are parasitoids of Lepidoptera. So far 54 genera and 207

species were reported from Iran (Riedel & Aghadokht, 2017). Tribe Heresiarchini Ashmead, 1900 includes 88 genera and 595 species worldwide and 88 species have been recorded from the Western Palaearctic region (Yu et al., 2016). Members of the tribe Heresiarchini are recognized by sickle-shaped mandibles, upper tooth longer than lower one, third pleural area (area posteroexterna) extremely sharply shortened (the apices of areae dentiparae especially close approach to the hind coxae) (Tereshkin, 2009). So far 6 genera and 33 species of Heresiarchini have been recorded from Iran (Heinrich, 1929;

Corresponding author: Ali Asghar Talebi, E-mail: talebia@modares.ac.ir

Copyright © 2018, Shirzadegan et al. This is an open access article distributed under the terms of the Creative Commons Attribution License (CC BY NC 4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Kolarov & Ghahari, 2005; 2008; Masnadi & Sadeghi, 2006; Masnadi & Jussila, 2008; Ghahari & Jussila, 2010; Ghahari et al., 2010 a, 2010b; Riedel et al., 2010; Ghahari & Jussila, 2011a, 2011b, 2011c, 2011d; Ghahari & Schwarz, 2011; Ghahari, 2012, 2014; Ghahari & Jussila, 2014; Ghahari & Gadalla, 2015; Ghahari & Jussila, 2016; Riedel & Aghadokht, 2017). Despite recent advances in the taxonomy of Ichneumonidae in Iran (Mohammadi-Khoramabadi et al., 2011; 2013a, 2013b; Amiri et al., 2015a, 2015b, 2016a, 2016b, 2017; Mohebban et al., 2015, 2016; Shirzadegan et al., 2017; 2018), our knowledge on the biology, ecology, and their role in Biological control of insect pests in Iran is low. It is difficult to give a precise estimate of the number of species of Ichneumoninae occurring in Iran, but considering the great diversity in vegetation, monsoon forest, natural ecosystems and farmlands, the number of recorded species, in comparison to the known species of subfamily Ichneumoninae in the Palearctic region is low (1297 species) (Yu et al., 2016). The results may be useful for future bio control of insect pests and ecological studies.

Material and methods

Malaise traps were used in different places consisting Ghazichak, Eshman Kamachal, Orkom, Ziaz located in Guilan province and Arangeh, Sarzariat, Shahrestanak, Karaj in Alborz province during March to November 2010. For the preparation of samples, they were preserved in a mixture of 40% xylene and 60% alcohol 96% for two days, then transferred into amyl acetate for two days and finally placed on the filter paper for drying (AXA method, van Achterberg, 2009). The dried specimens were then card-mounted and labeled. Photographs were taken using an Olympus™ SZX9 stereomicroscope equipped with BMZ-04-DZTM digital imaging system (Behin Pajouhesh Co., Iran). A

series of 10-15 captured images were merged into a single in-focus image using the image-stacking software ZereneStacker version 1.04. Morphological terminology follows Tereshkin (2009). Nomenclature and distribution data are mainly taken from Yu et al. (2016). The specimens are deposited at the Collection of Department of Entomology, Tarbiat Modares University (TMUC).

Results

Three species belonging to two genera of the tribe Heresiarchini were collected and identified, of which *Coelichneumon probator* Horstmann, 2000 is newly recorded for the Iranian fauna and the Middle East. The results of this study and review of the previously recorded taxa documented the existence of 34 species of the tribe Heresiarchini in Iran.

Coelichneumon comitator (Linnaeus, 1758)

Material examined: Iran, Alborz province, Karaj, Sarzariat (35°55'12" N, 51°06'54" E, 1980 m a.s.l.), 01.VI.2010, 1♀, 06.VII.2010, 1♀, leg.: M. Khayrandish.

Distribution in Iran: Not defined (Heinrich, 1929), Golestan (Kolarov & Ghahari, 2008); Tehran (Masnadi & Jussila, 2008); Khorasan-e razavi (Ghahari & Jussila, 2014).

General distribution: Western and Eastern Palearctic region (Yu et al., 2016)

Coelichneumon probator Horstmann, 2000 (Figs. 1; 2 A-F)

Material examined: Iran, Guilan province, Rudsar, Ghazichak (36°45'54" N, 50° 19' 36" E, 1803 m a.s.l.), 10.V.2010, 1♂, Leg.: M. Khayrandish.

Description (Male): Body length 11 mm; vertex gradually sloping behind ocelli to occipital carina from lateral view (Fig. 2A); temples short, 0.7 X as wide as transverse diameter of the eye in lateral view, sharply roundly narrowed behind the eyes in dorsal view; occipital carina connected to hypostomal carina far from base of

mandibles; face 3.0 X as wide as its length, its surface densely punctured (Fig. 2B); clypeus almost flat, approximately 2.4 X as wide as its length, its surface coarsely punctated (Fig. 2B); clypeal foveae small; malar space 0.5 X as long as basal width of mandible; flagellum with 31 flagellomeres, flagellomeres 7-16 with tyloides; mesonotum moderately convex and scarcely punctuated; scutellum slightly convex, as long as its wide, its lateral margin carinated up to middle (Fig. 2D); mesopleuron densely punctured, without microsculpture (Fig. 2C); area supermedia hexagonal and merged with area basalis, costulae before middle; apices of area dentiparae slightly prominent (Fig. 2D); propodeal spiracle elongate, its length 3.5 X as long as its width; third pleural area (area postero-externa) extremely sharply shortened (Fig. 2C); hind femur 3.5 X as long as its maximum width; hind tibia 1.8 X longer than hind basitarsus; postpetiole slightly elevated at the middle, 2 X wider than lateral fields, its surface longitudinally striated (Fig. 2E); 2nd and 3rd tergites coarsely and densely punctate, with some longitudinal rugose sculpture at central parts; gastrocoeli developed with several

irregular longitudinal ridges; thyridia distinct, oblique, interspace between them longitudinal rugose sculptures (Fig. 2E).

Coloration: Body black; triangular ivory spots on vertex, facial orbits, frontal orbits, genal orbits, spots at lateral corners of clypeus, pronotal ridge, tegula and subalarum and lateral carinae of scutellum with ivory; Hind legs black with reddish pattern; coxae and trochanters of all legs black.

Distribution in Iran: New for Iran and the Middle East fauna.

General distribution: west palaeartic, Belgium, Czech Republic, France, France-main, Germany, Poland, Romania, Spain, Spain-main, Ukraine, United Kingdom (Yu et al., 2016).

***Syspasis carinator* (Fabricius, 1798)**

Material examined: Iran, Alborz province: Karaj, Shahrestanak (35°57'36" N, 51°22'18" E, 2301 m a.s.l.), 03.VIII.2010, 1♀, leg.: M. Khayrandish.

Distribution in Iran: Tehran province (Ghahari, 2014)

General distribution: Western Palaeartic region (Yu et al., 2016).



Figure 1. *Coelichneumon probator*, male lateral habitus.




Figure 2. *Coelichneumon probator*, male: **A.** Head, lateral view, **B.** Head, frontal view, **C.** Mesosoma, lateral view, **D.** Scutellum and propodeum, dorsal view, **E.** First and second tergites, dorsal view, **F.** Metasoma, dorsal view.

Discussion

As a crossroad between the Eastern Mediterranean area and the Oriental region, Iran is home to representatives of the Middle East and Central Asian parasitoid faunas (Rakhshani et al., 2008). According to previous studies, 33 species belong to tribe Heresiarchini has been reported from Iran (Table 1). Most species are attributed to the genus *Coelichneumon*. The members of this tribe attacking lepidopterous larvae (19 families) (Yu et al., 2016), sometimes coleoptera (Curculionidae) and Hymenoptera (Cimbicidae, Pamphilidae)

are reported as their hosts (Dalla Torre, 1902; Rudow, 1917). In this study, the biology of recorded species is unknown because the specimens were collected by Malaise trap; The number of Heresiarchini species were reported in neighboring countries including: former USSR 209 species, Turkey 35 species, Tajikistan four species, Turkmenistan one species, Armenia one species, Afghanistan nine species, Azerbaijan 20 species and Pakistan three species (Yu et al., 2016). We expected that the tribe Heresiarchini fauna of Iran be substantially increased in next studies.

Table 1. Updated checklist of the tribe Heresiarchini species known in Iran.

Species	Distribution in Iran (provinces)	References
<i>Amblyjoppa fuscipennis</i> (Wesmael, 1845)	East Azarbaijan	Ghahari & Jussila (2010)
<i>Callajoppa caspica</i> (Heinrich, 1929)	Fars	Kolarov & Ghahari (2005)
<i>Coelichneumon afghanicus</i> (Heinrich, 1957)	Mazandaran	Kolarov & Ghahari (2008)
<i>Coelichneumon biannulatus</i> (Gravenhorst, 1820)	Ardabil; Mazandaran; Zanjan	Kolarov & Ghahari (2008); Ghahari & Jussila (2010, 2016)
<i>Coelichneumon biguttulatus</i> (Kriechbaumer, 1875)	Guilan, Mazandaran	Kolarov & Ghahari (2008)
<i>Coelichneumon bohemani</i> (Holmgren, 1864)	Lorestan	Ghahari & Gadalla (2015)
<i>Coelichneumon comitator</i> (Linnaeus, 1758)	Not defined; Golestan; Tehran; Khorasan-e Razavi, Alborz	Heinrich (1929); Kolarov & Ghahari (2008); Masnadi & Jussila (2008); Ghahari & Jussila (2014); current study
<i>Coelichneumon desinatorius</i> (Thunberg, 1822)	Qazvin; Golestan; Guilan	Kolarov & Ghahari (2008)
<i>Coelichneumon deliratorius</i> (Linnaeus, 1758)	Guilan; Golestan; Qazvin	Kolarov & Ghahari (2005, 2008); Masnadi & Jussila (2008)
<i>Coelichneumon dorsosignatus</i> (Berthoumieu & Eversmann, 1894)	Mazandaran	Kolarov & Ghahari (2008)
<i>Coelichneumon erythromerus</i> (Rudow, 1888)	Ardabil	Ghahari et al. (2010b)
<i>Coelichneumon falsificus</i> (Wesmael, 1845)	Khuzestan	Kolarov & Ghahari (2008)
<i>Coelichneumon fasciatus</i> (Gmelin, 1790)	Mazandaran	Ghahari & Jussila (2010)

Table 1. continued

<i>Coelichneumon iranicus</i> (Riedel, 2017)	East Azarbaijan, West Azarbaijan	Riedel & Aghadokht (2017)
<i>Coelichneumon leucocerus</i> (Gravenhorst, 1820)	Hamedan; East Azarbaijan; West Azarbaijan	Kolarov & Ghahari (2008); Ghahari & Jussila (2011b)
<i>Coelichneumon melanocastaneus</i> (Riedel, 2017)	Golestan	Riedel & Aghadokht (2017)
<i>Coelichneumon nigerrimus</i> (Stephens, 1835)	West Azarbaijan; Tehran	Kolarov & Ghahari (2005, 2008); Ghahari (2014)
<i>Coelichneumon nigrifrons</i> (Riedel, 2017)	Golestan	Riedel & Aghadokht (2017)
<i>Coelichneumon nigratus</i> (Berthoumieu, 1894)	Guilan	Kolarov & Ghahari (2005)
<i>Coelichneumon nobilis</i> (Wesmael, 1857)	Golestan	Riedel & Aghadokht (2017)
<i>Coelichneumon nudicoxator</i> (Aubert, 1966)	Semnan	Ghahari (2012)
<i>Coelichneumon orbitator</i> (Thunberg, 1822)	Chaharmahal-o-Bakhtiari; Esfahan; Markazi; Fars; Mazandaran	Kolarov & Ghahari (2008); Masnadi & Sadeghi (2006); Masnadi & Jussila (2008)
<i>Coelichneumon probator</i> Horstmann, 2000*	Guilan	Current study
<i>Coelichneumon problematicus</i> (Riedel, Coruh & Ozbek, 2010)	Tehran	Riedel et al. (2010)
<i>Coelichneumon singularis</i> (Berthoumieu, 1892)	Sistan and Baluchestan; Fars; East Azarbaijan	Kolarov & Ghahari (2008); Masnadi & Jussila (2008); Masnadi & Sadeghi (2006); Ghahari & Jussila (2011b)
<i>Coelichneumon validus</i> (Berthoumieu, 1894)	Tehran	Masnadi & Jussila (2008)
<i>Lymantrichneumon disparis</i> (Poda, 1761)	Mazandaran	Kolarov & Ghahari (2005, 2008)
<i>Protichneumon fusorius</i> (Linnaeus, 1761)	Mazandaran	Ghahari et al. (2010a)
<i>Protichneumon persicus</i> Morley, 1915	Golestan	Kolarov & Ghahari (2005)
<i>Protichneumon pisorius</i> (Linnaeus, 1758)	Khorasan-e Razavi; Mazandaran	Ghahari & Jussila (2011a, c); Ghahari & Schwarz (2011)
<i>Protichneumon similatorius</i> (Fabricius, 1798)	Not defined	Kolarov & Ghahari (2005)
<i>Sypsis carinator</i> (Fabricius, 1798)	Tehran, Alborz	Ghahari (2014), current study
<i>Sypsis rufina</i> (Gravenhorst, 1820)	Guilan; East Azarbaijan	Kolarov & Ghahari (2005); Ghahari & Jussila (2011d)
<i>Sypsis scutellator</i> (Gravenhorst, 1829)	Mazandaran; West Azarbaijan; Tehran	Kolarov & Ghahari (2008); Masnadi & Jussila (2008)

* New records for the Iranian insect fauna.

Acknowledgments

We would like to thank the Department of Entomology, Tarbiat Modares University for providing financial support for this research. Many thanks to Dr. Mohammad Khayrandish for helping us with the collecting of the specimens. We are grateful to three anonymous reviewers for constructive comments and recommendations on the earlier version of this paper.

Conflict of Interests

The authors declare that there is no conflict of interest regarding the publication of this paper.

References

- Aguiar, A., Deans, A.R., Engel, M.S., Forshage, M., Huber, J.T., Jennings, J.T., Johnson, N., Lelej, A.S., Longino, J.T., Lohrmann, V., Miko, I., Ohl, M., Rasmussen, C., Taeger, A. & Yu, D.S.K. (2013) Order Hymenoptera. *Zootaxa*, 3703 (1), 51–62.
<http://doi.org/10.11646/zootaxa.3703.1.12>
- Amiri, A., Talebi, A.A., Riedel, M., Rakhshani, E. & Hajiqaanbar, H. (2015a) A survey of Metopiinae (Hymenoptera: Ichneumonidae) in southern Iran, with three new records. *Journal of Crop Protection*, 4 (4), 519–531.
- Amiri, A., Talebi, A.A., Jussila, R., Rakhshani, E. & Hajiqaanbar, H. (2015b) A study of the Iranian Cremastinae (Hymenoptera: Ichneumonidae). *Journal of Insect Biodiversity and Systematics*, 1 (2), 87–100.
- Amiri A., Talebi, A.A., Jussila, R., Rakhshani, E. & Hajiqaanbar, H. (2016a) Study of the subfamily Ophioninae (Hymenoptera: Ichneumonidae) in southern Iran. *Journal of Entomological Society of Iran*, 35 (4), 53–67.
- Amiri, A., Talebi, A.A., Castillo, C. R., Rakhshani, E. & Hajiqaanbar, H. (2016b) Study of the genus *Lissonota* (Hymenoptera: Ichneumonidae: Banchinae) in southern Iran. *Journal of Entomological Society of Iran*, 36 (2), 89–99.
- Amiri, A., Talebi, A.A., Rakhshani, E. & Hajiqaanbar, H. (2017) First report of the genus *Cymodusa* (Ichneumonidae: Campopleginae) from Iran. *Journal of Insect Biodiversity and Systematics*, 3 (2), 81–89.
- Çoruh, S. & Özbek, H. (2013) New and little known some Ichneumonidae species (Hymenoptera) from Turkey. *Munis Entomology and Zoology*, 8, 135–139.
- Dalla Torre, C.G.de. (1902) *Catalogus Hymenopterorum. Volumen III. Trigonalidae, Megalyridae, Stephanidae, Ichneumonidae, Agriotypidae, Evaniidae, Pelecinidae.* Guilelmi Engelmann. *Lipsiae*, 1901, 1–544. 1902, 545–1141.
- Ghahari, H. & Gadallah, N.S. (2015) A study on the ichneumonid wasps (Hymenoptera: Ichneumonidae) from the province of Lorestan, Iran. *Arquivos Entomoloxicos*, 13, 329–338.
- Ghahari, H. & Jussila, R. (2010) Some new records of Iranian Ichneumoninae (Hymenoptera: Ichneumonidae). *Linzer Biologische Beiträge*, 42 (2), 1373–1377.
- Ghahari, H. & Jussila, R. (2011a) A study on the subfamilies Cremastinae, Ichneumoninae, Pimplinae and Rhyssinae (Hymenoptera: Ichneumonidae) from the Mazandaran province, Iran. *Calodema*, 140, 1–6.
- Ghahari, H. & Jussila, R. (2011b) A contribution to the knowledge of Ichneumonidae (Hymenoptera) from Arasbaran and vicinity, Iran. *Calodema*, 166, 1–5.
- Ghahari, H. & Jussila, R. (2011c) A study on the ichneumonid wasps (Hymenoptera: Ichneumonidae) from some regions of Iran. *Linzer Biologische Beiträge*, 43 (1), 753–758.
- Ghahari, H. & Jussila, R. (2011d) Five species of Ichneumonidae (Hymenoptera) new to Iran. *Zoology in the Middle East*, 52, 123–124.
<https://doi.org/10.1080/09397140.2011.10638490>
- Ghahari, H. & Jussila, R. (2014) A study on the subfamily Ichneumoninae (Hymenoptera: Ichneumonidae) from Khorasan province, Iran. *Linzer biologische Beiträge*, 46 (2), 1367–1371.
- Ghahari, H. & Jussila, R. (2016) Contribution to the knowledge of the fauna of Ichneumonidae (Hymenoptera: Ichneumonoidea) from Iran. *Beiträge zur Entomologie*, 66 (1), 119–124.

- Ghahari, H. & Schwarz, M. (2011) A contribution to the knowledge of ichneumonid wasps from Mazandaran province, northern Iran (Hymenoptera, Ichneumonidae). *Spixiana*, 34 (2), 195–198.
- Ghahari, H. (2012) A study on the Ichneumonidae (Hymenoptera) from Jangal-e Abr, Semnan province, Iran. *Calodema*, 201, 1–4.
- Ghahari, H. (2014) A study on the subfamily Ichneumoninae (Hymenoptera: Ichneumonidae) from Varamin and vicinity, Iran. *Calodema*, 295, 1–2.
- Ghahari, H., Jussila, R., Kolarov, J. & Šedivý, J. (2010a) A Contribution to the Ichneumon wasps (Hymenoptera: Ichneumonidae) from the forests of Northern Iran. *Munis Entomology and Zoology*, 5 (1), 85–89.
- Ghahari, H., Cetin Erdogan, O., Šedivý, J. & Ostovan, H. (2010b) Survey of the Ichneumonidea and Chalcidoidea (Hymenoptera) parasitoids of Saturniidae (Lepidoptera) in Iran. *Efflatounia*, 10, 1–6.
- Heinrich, G.H. (1929) Beiträge zur Kenntnis der Insektenfauna Nordpersiens. Nach der Ausbeute der Elbrus-Expedition von G. Heinrich und Dr. M. Dammholz Anno 1927. Ichneumonidae (Hym.). *Deutsche Entomologische Zeitschrift*, 305–326.
- Kolarov, J. & Ghahari, H. (2005) A catalogue of Ichneumonidae (Hymenoptera) from Iran. *Linzer biologische Beiträge*, 37, 503–532.
- Kolarov, J. & Ghahari, H. (2008) A study of the Iranian Ichneumonidae (Hymenoptera) III. Ichneumoninae. *Acta Entomologica Serbica*, 13 (1/2), 61–76.
- Masnadi, A. & Jussila, R. (2008) Contribution to the knowledge of ichneumonid wasps of Iran. Subfamilies Ichneumoninae, Pimplinae and Diplazontinae (Hymenoptera, Ichneumonidae). *Entomofauna*, 29 (22), 293–320.
- Masnadi, A. & Sadeghi, S.E. (2006) New record on two species of ichneumonid wasps in Iran (*Coelichneumon singularis* Berthoumieu, *C. microstictus* Gravenhorst (Ichneumonidae: Ichneumoninae). *Applied Entomology and Phytopathology*, 73 (2), 46.
- Mohammadi-Khoramabadi, A., Talebi, A.A. & Farahani, S. (2011) *Hybrizon buccatus* (de Brebisson, 1825), the first record of the subfamily Hybrizontinae (Hymenoptera: Ichneumonidae) from northern Iran. *Biharean Biologist*, 5, 162–163.
- Mohammadi-khoramabadi, A., Talebi, A.A. & Zwakhals, K. (2013a) A study of the subfamily Pimplinae (Hymenoptera: Ichneumonidae) in the north of Iran, with eleven new species records. *Entomofauna*, 34(2), 29–56.
- Mohammadi-Khoramabadi, A., Talebi, A.A. & Zwakhals, K. (2013b) Study on Diplazontinae (Hymenoptera: Ichneumonidae) in the north central of Iran. *Journal of Crop Protection*, 2, 241–261.
- Mohebban, S., Takaloozadeh, H., Barahoei, H. & Madjdzadeh, M. (2015) New records of Cryptinae and Ichneumoninae (Hymenoptera: Ichneumonidae) species from Kerman province, Southeast Iran. *Journal of Crop Protection*, 4 (3), 337–349.
- Mohebban, S., Barahoei, H., Takaloozadeh, H.M., Madjdzadeh, S.M. & Riedel, M. (2016) A survey of the Ichneumonidae (Hymenoptera, Ichneumonoidea) of Kerman province, south-east Iran. *Journal of Insect Biodiversity and Systematics*, 2 (4), 419–437.
- Rakhshani, E., Talebi, A.A., Starý, P., Tomanović, Ž., Kavallieratos, N.G. & Manzari, S.A. (2008) Review of *Aphidius* Nees in Iran: host associations, distribution and taxonomic notes. *Zootaxa*, 1767, 37–54.
- Riedel, M. & Aghadokht, P. (2017) Contribution to the Ichneumoninae (Hymenoptera: Ichneumonidae) of Iran, with descriptions of three new species, *Zoology in the Middle East*, 63 (4), 348–355.
<http://doi.org/10.1080/09397140.2017.1361190>
- Riedel, M., Çoruh, S. & Özbek, H. (2010) Contribution to the Ichneumoninae (Hymenoptera, Ichneumonidae) fauna of Turkey, with description of three new species. *Türkiye Entomoloji Dergisi*, 34 (2), 133–156.

- Rudow, F. (1917) Die Ichneumonidengattung *Amblyteles* und ihre Wirte. *Entomologische Zeitschrift*, 31, 25–26, 31, 33–35.
- Shirzadegan, F., Talebi, A.A., Riedel, M. & Hajiqanbar, H. (2017) Two newly recorded species of the tribe Platylabini (Hymenoptera: Ichneumonidae, Ichneumoninae) from Iran. *Journal of Crop Protection*, 6 (3), 401–408.
- Shirzadegan, F., Talebi, A.A., Riedel, M. & Hajiqanbar, H.R. (2018) Study of the genus *Platylabus* Wesmael, 1845 (Hymenoptera: Ichneumonidae) in north of Iran, with a new record for the Middle East. *Journal of Insect Biodiversity and Systematics*, 4 (1), 1–11.
- Tereshkin, A.M. (2009) Illustrated key to the tribes of subfamilia Ichneumoninae and genera of the tribe Platylabini of world fauna (Hymenoptera, Ichneumonidae). *Linzer Biologische Beiträge*, 41 (2), 1317–1608.
- van Achterberg, C. (2009) Can Townes type Malaise traps be improved? Some recent developments. *Entomologische Berichten*, 69, 129–135.
- Yu, D.S., van Achterberg, K. & Horstmann, K. (2016) Home of Ichneumonoidea, 2011. Taxonomy, Biology, Morphology and Distribution. (CD-ROM) Taxapad (available at <http://www.taxapad.com/>).

مطالعه قبیله *Heresiarchini* (Hym., Ichneumonidae, Ichneumoninae) در
شمال ایران، به همراه یک گزارش جدید برای خاورمیانه

فرید شیرزادگان^۱، علی اصغر طالبی^{۱*}، ماتیاس ریدل^۲ و حمیدرضا حاجی قنبر^۱

۱ گروه حشره‌شناسی، دانشکده کشاورزی، دانشگاه تربیت مدرس، صندوق پستی ۳۳۶-۱۴۱۱۵، تهران، ایران

۲ موزه جانورشناسی مونیخ، خیابان مونیخ هاوزن ۲۱، ۸۱۲۴۷-d، آلمان

* پست الکترونیکی نویسنده مسئول مکاتبه: talebial@modares.ac.ir

تاریخ دریافت: ۱۷ بهمن ۱۳۹۶، تاریخ پذیرش: ۰۲ تیر ۱۳۹۷، تاریخ انتشار: ۰۳ مهر ۱۳۹۷

چکیده: در این تحقیق قبیله *Heresiarchini* (Hymenoptera: Ichneumonidae, Ichneumoninae) در استان‌های البرز و گیلان مورد مطالعه رده‌بندی قرار گرفت. نمونه‌ها با استفاده از تله مالیز طی ماه‌های اسفند تا آذر ۱۳۸۸ و ۱۳۸۹ جمع‌آوری شدند. سه گونه شامل *Coelichneumon comitator* (Linnaeus, 1758)، *Syspasis carinator* (Fabricius, 1798) و *Coelichneumon probator* Horstmann, 2000 (که از میان آنها گونه *Coelichneumon probator* برای اولین بار از ایران و خاورمیانه گزارش می‌شود. چک‌لیست به روز شده قبیله *Heresiarchini* به همراه خصوصیات افتراقی گزارش جدید ارائه شده است.

واژگان کلیدی: *Heresiarchini*, *Ichneumoninae*, *Ichneumonidae*, *Coelichneumon*, ایران