



## Taxonomical investigation on Braconidae (Hymenoptera: Ichneumonoidea) fauna in northern Cyprus, with twenty six new records for the country

Ahmet Beyarslan<sup>1\*</sup>, Celalettin Gözüaçik<sup>2</sup>, Mustafa Güllü<sup>3</sup> and Ayda Konuksal<sup>4</sup>

<sup>1</sup> Department of Biology, Faculty of Arts and Sciences, Bitlis Eren University, 13000 Bitlis, Turkey.

<sup>2</sup> Department of Plant Protection, Faculty of Agriculture, Iğdır University, Iğdır, Turkey.

<sup>3</sup> Biological Control Research Institute, Adana, Turkey.

<sup>4</sup> Agricultural Research Institute, Lefkoşa, Turkish Republic of Northern Cyprus.

**ABSTRACT.** In order to determine Braconidae fauna of Cyprus, adult specimens of Braconidae were collected from different habitats of Northern Cyprus between 2012 and 2016. All specimens were collected in both natural vegetation and agricultural areas using sweeping-net. A total of 42 species belonging to 14 genera of Agathidae, Braconinae, Cheloninae, Doryctinae, Euphorinae, Homolobinae, Opiinae and Rogadinae were reported from the Northern Cyprus, of which 26 species are recorded from Cyprus for the first time.

**Key words:** Pentadaktylos, Braconidae, fauna, Cyprus, parasitoid.

**Received:**  
25 September, 2017

**Accepted:**  
21 October, 2017

**Published:**  
24 October 2017

**Subject Editor:**  
Ehsan Rakhshani

**Citation:** Beyarslan, A., Gözüaçik, C., Güllü, M. & Konuksal, A. (2017) Taxonomical investigation on Braconidae (Hymenoptera: Ichneumonoidea) fauna in northern Cyprus, with twenty six new records for the country. *Journal of Insect Biodiversity and Systematics*, 3 (4), 319–334.

### Introduction


Braconidae is a large family of Hymenoptera, consisting of 1032 genera and approximately 17,963 known species. Thirty seven of 47 subfamilies of the Braconidae are recorded from the west Palaearctic region (Yu et al., 2012). The vast majority of the species are ectoparasitoids mainly among coleopterous and lepidopterous hosts, although a few attack Diptera, Hymenoptera-Symphyta and possibly Hemiptera (Wang et al., 2010).

Cyprus is divided into four geological zones: (a) the Pentadaktylos (Kyrenia) Zone; (b) the Troodos Zone or Troodos Ophiolite; (c) the Mamonia Zone or Complex, and (d) the Zone of autochthonous sedimentary

rocks (Fig. 1). The Pentadaktylos (Kyrenia) Zone, the northern-most geological zone of Cyprus, is considered to be the southern-most portion of the Tauro-Diranide Alpine Orogenic Zone. The base of the Zone is mostly composed of a series of allochthonous massive and recrystallized limestones, dolomites and marbles of Permian-Carboniferous to Lower Cretaceous age (between 350–135 Ma). These are stratigraphically followed by younger autochthonous sedimentary rocks of Upper Cretaceous to Middle Miocene age (67–15 Ma), on which the older allochthonous formations have been thrust southward (Anonymous, 2004).

Corresponding author: Ahmet Beyarslan, E-mail: [abeyars@gmail.com](mailto:abeyars@gmail.com)

Copyright © 2017, Beyarslan et al. This is an open access article distributed under the terms of the Creative Commons Attribution License (CC BY NC 4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.



**Figure 1.** Four geological zones of Cyprus (Anonymous, 2004)

The area of this research covers the Pentadaktylos (Kyrenia) Zone and partially the Troodos Zone or Troodos Ophiolite. Cyprus is located in the eastern Mediterranean basin, and is the third largest (9 251 km<sup>2</sup>) Mediterranean island after Sicily (25 710 km<sup>2</sup>) and Sardinia (24 090 km<sup>2</sup>). Cyprus is 70 km from Turkey (Syria 110 km, Egypt 418 km, and Greece 965 km). The Braconidae fauna of Cyprus consists of 124 species. It is similar with 65% (81 species) of Turkey, 64% (79 species) of Greece, 14.4% (18 species) of Egypt and 9.6% (12 species) of Syria (Yu et al., 2012).

The Braconidae fauna of Cyprus has been partially investigated. In Cyprus, there are known 124 species of Braconidae according to Terrestrial Arthropods of Cyprus (Anonymous, 2015). The first study of the Braconinae fauna of Cyprus was conducted by Papp (1990) and he worked the fauna of Greece and however, he also identified the Braconids of south Cyprus. We conducted a study on Braconidae fauna in northern Cyprus between 2012 and 2016. A total of 42 species belonging to 14 genera of Agathidinae, Braconinae, Cheloninae,

Doryctinae, Euphorinae, Homolobinae, Opiinae and Rogadinae were reported from Northern Cyprus. Twenty-six species are recorded from Cyprus for the first time.

#### Material and methods

Adult specimens of Braconidae were collected from different habitats of northern Cyprus by sweeping-nets between 2012 and 2016. Publications consulted for distributional data include: Fischer (1999), Nixon (1986), Papp (1982, 1990, 1998, 1999), Shenefelt (1978), Tobias (1986), and Yu et al. (2012). The following literature was used for taxonomic examination of the material: Matthews (1970), Shaw & Huddleston (1991), Fischer (1999), Achterberg (2003), Marsh & Shaw (2003), Fischer & Beyarslan (2011, 2012, 2013), Beyarslan et al. (2013) and Beyarslan (2014, 2015b).

Information on parasitoids, hosts and general distributions (in terms of zoogeographical region), and parasitoids of the determined species are given according to Yu et al. (2012). The subgenera and species are listed alphabetically. Known

host plants are given in square brackets (Yu et al., 2012). All specimens are deposited in the collection of the Department of Zoology, Bitlis Eren University, Bitlis, Turkey. An asterisk (\*) is used within the text for the species that are recorded as new from Cyprus.

## Results

### Subfamily: Agathidinae

#### Genus: *Agathis* Latreille, 1804

##### \**Agathis malvacearum* Latreille, 1805

**Material examined:** Northern Cyprus-İnönü, wild herb, 09.04.2015, 1♀; İskele, barley, 07.04.2015, 3♂♂.

**Hosts:** DIPTERA, **Tachinidae:** *Actia nudibasis* Stein, 1924; LEPIDOPTERA, **Coleophoridae:** *Coleophora galbulipennella* Zeller, 1838 [*Silene* sp.]; *C. graminicolella* Heinemann, 1876; **Gelechiidae:** *Metzneria aestivella* (Zeller, 1839); *M. lappella* (Linnaeus, 1758) [*Arctium minus*]; *Pexicopia malvella* (Hübner, 1805); **Pterophoridae:** *Hellinsia didactylites* (Ström, 1783).

**Distribution:** Holarctic, Cyprus (new record).

### Subfamily: Braconinae

#### Genus: *Bracon* Fabricius, 1804

##### Subgenus: *Bracon* 1804 s.str.

##### \**Bracon (Bracon) gusaricus* Telenga, 1933

**Material examined:** Northern Cyprus - İskele, barley, 07.04.2015, 4♀♀, 1♂.

**Hosts:** Unknown.

**Distribution:** Palaearctic, Cyprus (new record).

##### *Bracon (Bracon) intercessor* Nees, 1834

**Material examined:** Northern Cyprus - Beşparmak dağları, Taşocağı, wild herb, 08.04.2015, 1♂; Bostancı, wheat, 27.03.2012, 2♀♀; Dıpkarpaz, Büyük Kumsal, barley, 07.04.2015, 1♂; Girne, Çatalköy, wild herb, 08.04.2015, 1♂; Göçeri wild herb, 30.05.2012, 1♀; İnönü, barley, 09.04.2015,

1♂; İskele, wild herb, 28.03.2012, 2♂♂; Köprülü, wild herb, 01.06.2012, 2♂♂; Mehmetcik, wild herb, 31.05.2016, 1♂; Şirinevler, barley, 30.05.2012; 1♂; Taşpınar, wheat, 31.05.2012, 2♂♂; Türkmenköy, barley, 01.07.2012, 2♀♀; Yeşilırmak, barley, 01.6.2016, 2♀♀, 2♂♂.

**Hosts:** COLEOPTERA, **Apionidae:** *Oxystoma opeticum* Bach, 1854; **Attelabidae:** *Attelabus nitens* (Scopoli, 1763); **Cerambycidae:** *Agapanthia villosoviridescens* (De Geer, 1775); *A. violacea* (Fabricius, 1775); *Anthonomus (Anthonomus) pedicularius* (Linnaeus, 1758); *A. pomorum* Linnaeus 1758; *A. (A.) sorbi* Germar, 1821; *Hytoecia coerulescens* (Scopoli, 1763) [*Anchusasp.*]; *Curculio salicivorus* (Paykull, 1792); *C. crux* (Fabricius, 1776); *Lixus brevisrostris* Boheman, 1835; *L. incanescens* Boheman, 1835 [*Atriplex patula hastata*]; *L. (Compsolixus) juncii* Boheman, 1835; *Microlarinus lareynii* (Jacquelin du Val, 1852); *M. lypriformis* (Wollaston, 1861); *Sibinia (Sibinia) femoralis* Germar, 1824; **Rhynchitidae:** *Rhynchites bacchus* (Linnaeus, 1758); **DIPTERA, Agromyzidae:** *Liriomyza huidobrensis* (Blanchard, 1926); **HYMENOPTERA, Eurytomidae:** *Tetramesa hyalipennis* (Walker, 1832); *Tetramesa rossica* (Rimsky-Korsakov, 1914); **Tenthredinidae:** *Pontania bella* (Zaddach, 1876); *P. kriechebaumeri* Konow, 1901 [*Salix nigricans*]; *P. pedunculi* (Hartig, 1837); *P. vesicator* (Bremi-Wolf, 1849); *P. viminalis* (Linnaeus, 1758); *P. (Eupontania) acutifoliae* Zinovjev, 1985; **LEPIDOPTERA, Coleophoridae:** *Coleophora atraphaxidellum* Kuznetsov, 1957; **Elachistidae:** *Parametriotes theae* Kuznetsov, 1916; **Gelechiidae:** *Scrobipalpa obsoletella* (Fischer von Röslerstamm, 1841); **Sesiidae:** *Bembecia hungarica* (Tomala, 1910); *Chamaesphecia astatifformis* (Herrich-Schaffer, 1846); *Paranthrene tabaniformis* (Rottemburg, 1775); *Synanthedon culiciformis* (Linnaeus, 1758); **Tortricidae:** *Sparganothis pilleriana* (Denis & Schiffermüller, 1775); **Yponomeutidae:** *Argyresthia conjugella* Zeller, 1839.

**Distribution:** Palaearctic, Cyprus (Papp 1998).

***Bracon (Bracon) pectoralis* (Wesmael, 1838)**

**Material examined:** Northern Cyprus - Akdeniz, wheat, 01.06.2016, 2♀♀; Beşparmak dağları, Taşocağı, barley, 08.04.2015, 1♂; Mehmetcik, wild herb, 30.05.2016, 1♀; Serdarlı, wild herb, 30.05.2016, 4♀♀; Yeşil köy, barley, 30.05.2016, 3♀♀, 5♂♂.

**Hosts:** COLEOPTERA, Curculionidae: *Larinus flavescens* Germar, 1824; DIPTERA, Tephritidae: *Xyphosia miliari* (Schrank, 1781); HYMENOPTERA, Cynipidae: *Aphelonyx cerricola* (Girayd, 1859); LEPIDOPTERA, Alucitidae: *Alucita hexadactyla* (Linnaeus, 1758); *A. huebneri* Wallengren, 1859; Gelechiidae: *Sitotroga cerealella* (Olivier, 1789); Pyralidae: *Etiella zinckenella* (Treitschke, 1832); Tortricidae: *Aethes francillana* (Fabricius, 1794).

**Parasitoids:** HYMENOPTERA, Torymidae: *Exopristus trigonomerus* (Masi, 1916). Ichneumonidae: *Scambus elegans* (Woldstedt, 1877).

**Distribution:** Palaearctic, Cyprus (Papp 1990, 1998).

***Bracon (Bracon) luteator* Spinola, 1808**

**Material examined:** Northern Cyprus - Akdeniz, wild herb, 01.05.2016, 1♂; Dipkarpaz, barley, 31.05.2016, 2♀♀ 4♂♂; Serdarlı, wild herb, 30.30.05.2016, 1♀, 1♂; Taşpınar, wheat, 31.05.2012, 2♀♀; Türkmenköy Karpaz, barley, 01.07.2012, 1♀; wild herb, 31.05.2016, 1♂; Yeşilirmak, wild herb, 01.05.2016, 1♀, 1♂; Yeşilköy, barley, 30.5.2016, 1♀.

**Hosts:** DIPTERA, Tephritidae: *Urophora solstitialis* (Linnaeus, 1758); LEPIDOPTERA, Gelechiidae: *Metzneria aestivella* (Zeller, 1839); *Metzneria kerzhneri* Piskunov, 1979.

**Distribution:** Palaearctic, Cyprus (Papp, 1998).

**Subgenus: *CyanopteroBracon* Tobias, 1957  
*Bracon (CyanopteroBracon) armeniacus* Telenga, 1936**

**Material examined:** Northern Cyprus - Akdeniz, wild herb, 01.06.2016, 1♀.

**Hosts:** Unknown.

**Distribution:** Western Palaearctic, Cyprus (Papp, 1998).

**Subgenus: *GlabroBracon* Fahringer, 1927  
\**Bracon (GlabroBracon) ahngeri* Telenga, 1936**

**Material examined:** Northern Cyprus - Girne çatal köy, barley, 8.4.2015, 1♀.

**Hosts:** Unknown.

**Distribution:** Oriental, Palaearctic, Cyprus (new record).

***Bracon (GlabroBracon) atrator* Nees, 1834**

**Material examined:** Northern Cyprus - Ercan havaalanı, wheat, 8.4.2015, 1♀; Serhatköy, Güzelyurt kavşağı, barley, 09.04.2015, 1♀.

**Hosts:** COLEOPTERA, Brentidae: *Apion buddenbergi* Bedel (1885); *Gymnetron antirrhini* (Paykull, 1800); *G. campanulae* Schoenherr, 1825; *G. villosulum* Gyllenhal, 1838 [*Veronica* sp.]; DIPTERA, Agromyzidae: *Amauromyza flavifrons* Meigen, 1830; Tephritidae: *Tephritis conura* Loew, 1844; *T. neesii* Meigen, 1830, *T. separata* Rondani, 1871; LEPIDOPTERA, Coleophoridae: *Coleophora coronillae* Zeller, 1849.

**Distribution:** Palaearctic, Cyprus (Papp, 1998).

***Bracon (GlabroBracon) delibator* Haliday, 1833**

**Material examined:** Northern Cyprus - Lefkoşa, Dikkaya, wild herb, 09.04.2015, 1♀.

**Hosts:** COLEOPTERA, Curculionidae: *Gymnetron campanulae* Schoenherr, 1825; *Hylobius piceus* (De Geer, 1775); Phalacridae: *Olibrus aeneus* Fabricius, 1792 [*Tripleurospermum perforatum*]; DIPTERA, Tephritidae: *Urophora cuspidata* Meigen, 1826; LEPIDOPTERA, Tortricidae: *Cydia strobilella* (Linnaeus, 1758).

**Distribution:** Palaearctic, Cyprus (Papp, 1998).

**\**Bracon (GlabroBracon) lividus* Telenga, 1936**

**Material examined:** Northern Cyprus - Akdeniz, wild herb, 01.06.2016, 1♀; Türkmenköy, wild herb, 01.07.2012, 5♀♀.

**Hosts: HYMENOPTERA, Tenthredinidae:** *Pontania vesicator* (Bremer, 1849).

**Distribution:** Palaearctic, Cyprus (new record).

**\**Bracon (Glabrobracon) otiosus* Marshall, 1885 (syn. *macrurus*)**

**Material examined:** Northern Cyprus - Yeşilirmak, wild herb, 01.06.2016, 2♂♂; Yeşilköy, wild herb, 30.05.2016, 1♂.

**Hosts: COLEOPTERA, Curculionidae:** *Anthonomus pomorum* (Linnaeus, 1758); *Magdalis rufa* Porta, 1932; **HYMENOPTERA, Tenthredinidae:** *Nematus (Bacconematus) pumilio* (Konow, 1903); **LEPIDOPTERA, Agonoxenidae:** *Blastodacna atra* (Haworth, 1828).

**Distribution:** Palaearctic, Cyprus (new record).

**\**Bracon (Glabrobracon) parvulus* (Wesmael, 1838)**

**Material examined:** Northern Cyprus - Türkmenköy, wild herb, 01.07.2012, 1♀.

**Hosts: DIPTERA, Agromyzidae:** *Amauromyza flavifrons* Meigen, 1830; **Cecidomyiidae:** *Clinodiplosis strobili* Kieffer, 1909; **Tephritidae:** *Chaetostomella cylindrica* Robineau-Desvoidy, 1830 [*Chrysanthemum* sp.]; *Tephritis bardanae* Schrank, 1803; *T. conura* Loew, 1844 [*Cirsium oleraceum*]; *T. pulchra* Loew, 1844 [*Chrysanthemum leucanthemum*, *Scorzonera jaquiniana*]; *Trupanea stellata* (Fuessly, 1775).

**Distribution:** Oriental and Palaearctic, Cyprus (new record).

***Bracon (Glabrobracon) popovi* Telenga, 1936**

**Material examined:** Northern Cyprus - Mehmetcik, barley, 30.06.2016, 1♀.

**Hosts:** Unknown.

**Distribution:** Palaearctic, Cyprus (Papp, 1998).

***Bracon (Glabrobracon) variator* Nees, 1811**

**Material examined:** Northern Cyprus - Türkmenköy, wheat, 01.07.2012, 4♂♂;

Yeşilirmak, barley, 01.06.2016, 1♀.

**Hosts: COLEOPTERA, Anobiidae:** *Ernobius nigrinus* (Sturm, 1837); **Buprestidae:** *Coraebus florentinus* (Herbst, 1801); **Chrysomelidae:** *Bruchidius lividimanus* (Gyllenhal, 1833); *B. poupillieri* Allard, 1868; *Bruchus atomarius* (Linnaeus, 1761); *B. laticollis* Boheman, 1833; *B. lentis* Frölich, 1799; *B. viciae* Olivier, 1795; **Curculionidae:** *Anthonomus pomorum* (Linnaeus, 1758); *Baris chlorizans* Germar, 1837; *B. cuprirostris* Germar, 1817; *B. laticollis* (Marsham, 1802); *Ceutorhynchus assimilis* Paykull, 1792; *C. punctiger* (Gyllenhal, 1837); *Gymnetron asellus* (Gravenhorst, 1807); *G. campanulae* Schoenherr, 1825 [*Campanula* sp.]; *G. tetrum* (Fabricius, 1792); *Larinus flavescens* Germar, 1824; *L. jaceae* (Fabricius, 1775); *L. planus* Germar, 1824; *L. turbinatus* (Gyllenhal, 1836); *Magdalis rufa* Germar, 1824; *Pissodes validirostris* Gyllenhal, 1835; *Sibinia viscaria* Linnaeus, 1761; *S. viscaria* Linnaeus, 1761; *Sitona longulus* Gyllenhal, 1834; **DIPTERA, Anthomyiidae:** *Anthomyia albimana* Zetterstedt, 1845; *Botanophila seneciella* Meade, 1892; **Cecidomyiidae:** *Mikiola fagi* (Hartig, 1839); **Tephritidae:** *Sphenella marginata* Fallen, 1814; *Chaetostomella cylindrica* (Robineau-Desvoidy, 1830); *Noeta pupillata* Fallen, 1814; *Tephritis leontodontis* (DeGeer, 1776); *Urophora (Carduus) nutans* Fabricius, 1775); **HYMENOPTERA, Tenthredinidae:** *Hoplocampa minuta* (Christ, 1791); *H. brevis* (Klug, 1816); *H. flava* (Linnaeus, 1761); *Caliroa cerasi* (Linnaeus, 1758); **LEPIDOPTERA, Amphipyridae:** *Gortyna xanthenes* Germar, 1844 [*Cynara scolymus*]; **Carposinidae:** *Carposina niponensis* Walsingham, 1900; **Coleophoridae:** *Coleophora coronillae* Zeller, 1849; *C. medelichensis* Krone, 1908; **Gelechiidae:** *Mesophleps corsicella* Herrich-Schäffer, 1856; *Pexicopia malvella* (Huebner, 1805); *Platyedra subcinerea* (Haworth, 1828); **Geometridae:** *Perizoma lugdunaria* Herrich-Schäffer, 1855; **Gracillariidae:** *Phyllonorycter mespillella*

(Huebner, 1805); **Hadeninae:** *Hadena bicruris* (Hufnagel, 1766); **Lycaenidae:** *Lampides boeticus* (Linnaeus, 1767); **Noctuidae:** *Myelois circumvoluta* (Fourcroy, 1785) (*Onopordum acanthium*); **Nymphalidae:** *Vanessa cardui* Linnaeus, 1758; **Pyralidae:** *Etiella zinckenella* (Treitschke, 1832); *Myelois circumvoluta* (Fourcroy, 1785); *Dioryctria abietella* ([Denis & Schiffermüller], 1775); **Sesiidae:** *Synanthedon andrenaeformis* Laspeyres, 1801; **Tortricidae:** *Acleris variegana* ([Denis & Schiffermüller], 1775) [*Malus* sp.]; *Barbara herrichiana* Obraztsov 1960; *Cydia dorsana* Fabricius, 1775; *C. funebrana* (Treitschke, 1835); *Eucosma cana* (Haworth, 1811); *Gypsonoma aceriana* (Duponchel, 1843); *Pandemis cerasana* (Huebner, 1786); *Rhopobota naevana* (Malus) (Huebner, 1817); *Rhyacionia resinella* (Linnaeus, 1758). **Distribution:** Palaearctic, Cyprus (Papp, 1990, 1998).

**Subgenus: Orthobracon Fahringer, 1927**

**\*Bracon (Orthobracon) longigenis Tobias, 1957**

**Material examined:** Northern Cyprus - İskele, wild herb, 07.04.2015, 1♀.

**Hosts:** Unknown.

**Distribution:** Western Palaearctic, Cyprus (new record).

**Subgenus: Pigeria van Achterberg, 1985**

**Bracon (Pigeria) piger (Wesmael, 1838)**

**Material examined:** Northern Cyprus - Mehmetcik, barey, 30.06.2016, 1♀, 2♂♂; Girne, Çatalköy, wild herb, 08.04.2015, wheat, 3♂♂; Mağosa, Nergizli, wild herb, 09.04.2015, 1♀; Devlet Üretme çitliği, wheat, 09.04.2015, 1♂.

**Hosts:** COLEOPTERA, Curculionidae: *Pissodes validirostris* (Sahlberg, 1834); LEPIDOPTERA, Noctuidae: *Heliothis peltigera* ([Denis & Schiffermüller], 1775); Pyralidae: *Etiella zinckenella* (Treitschke, 1832); Tortricidae: *Cnephasia longana*

(Haworth, 1811); *Cydia nigricana* (Fabricius, 1794).

**Distribution:** Holarctic, Cyprus (Papp, 1998).

**Subgenus: Rostrobracon Tobias, 1957**

**Bracon (Rostrobracon) urinator (Fabricius, 1798)**

**Material examined:** Northern Cyprus - Akdeniz, barley, 01.06.2016, 4♀♀; Dipkarpaz, barley, 31.05.2016, 5♀♀; Göçeri wild herb, 30.05.2012, 1♀; Köprüköy, wild herb, 30.05.2016, 1♂; Serdarlı, barey, 30.05.2016, 1♀; Taşpınar, wheat, 31.05.2012, 2♀♀; Türkmenköy, barley, 01.07.2012, (4♀♀; Yeşilirmak, wild herb, 01.06.2016, 1♀.

**Hosts:** COLEOPTERA, Curculionidae: *Larinus flavescens* Germar, 1824; *L. saussureae* Marshall, 1924; *L. sibiricus* Gyllenhal, 1835 [*Xeranthemum annuum*]; *L. sturnus* (Schaller, 1783) [*Cirsium heterophyllum*]; *L. vulpes* (Olivier, 1807) [*Echinops ritro*]; *Lixus obesus* Petri, 1904 [*Prangos uloptera*]; *Rhinocyllus conicus* (Frölich, 1792) [*Carduus nutans*]; *R. latirostris* Schoenherr, 1826 [*Carduus nutans*]; DIPTERA, Lonchaeidae: *Protearomyia nigra* (Meigen, 1826); Tephritidae: *Tephritis pulchra* Loew, 1844 [*Scorzonera hispanica*].

**Distribution:** Afrotropical, Oriental and Palaearctic, Cyprus (Papp, 1990, 1998).

**Genus: Habrobracon Ashmead, 1895**

**\*Habrobracon concolorans (Marshall, 1900) (syn. nigricans)**

**Material examined:** Northern Cyprus - Akıncılar, wild herb, 09.04.2015, 2♀♀; Beşparmak dağı, Taşocağı, wheat, 08.04.2015, 1♀, 3♂♂; Devlet Üretme çitliği, wild herb, 09.04.2015, 1♂; Ercan Havaalanı, wild herb, 09.04.2015, 2♀♀; 1♂; Girne, Çatalköy, wheat, 08.04.2014, 1♀; Girne, Tepebaşı, wild herb, 09.04.2015, 1♂; İnönü, wild herb, 09.04.2015, 1♀, 1♂; İskele, wild herb, 08.04.2015, 3♂♂; Lefkoşa, Dilekkaya, wild herb, 09.04.2015, 7♂♂; Mağosa, Nergizli, wheat, 09.04.2015, 5♂♂; Türkmenköy, wheat, 09.04.2015, 2♂♂.



**Hosts: Leiodoptera: Gelechiidae:** *Pexicopia malvella* (Huebner, 1805); **Pyralidae:** *Etiella zinckenella* (Treitschke, 1832); **Crambidae:** *Loxostege sticticalis* (Linnaeus, 1761).

**Tortricidae:** *Cnephasia sedana* Constant 1884.

**Distribution:** Oriental, Palaearctic, Cyprus (new record).

***Habrobracon stabilis* (Wesmael, 1838)**

**Material examined:** Northern Cyprus - Ercan Hava Alanı, wheat, 09.04.2015, 2♀♀, 5♂♂; Gazimağusa, İnönü, wild herb, 09.04.2015, 4♀♀, 11♂♂; Gazimağusa, Nergizli, barley, 09.04.2015, 9♀♀, 2♂♂; Gazimağusa, Türkmen köy, wild herb, 09.04.2015, 9♀♀, 2♂♂; Girne, Başpınar (Beşparmak Taşocağı, 2), barley, 08.04.2015, 5♀♀; Girne, Tepebaşı, wheat, 09.04.2015, 1♀, 1♂; İskele, wheat, 07.04.2015, 2♀♀, 8♂♂; Lefkoşa, Dilekkaya, wheat, 09.04.2015, 2♀♀, 4♂♂; Lefkoşe, Akıncılar, weeds, 09.04.2015, 1♀; Lefkoşe, Çömlekçi (Devlet üretim çiftliği), barley, 09.04.2015, 4♀♀, 3♂♂.

**Hosts: COLEOPTERA, Anobiidae:** *Ernobius kiesewetteri* Schilsky, 1898; *E. nigrinus* (Sturm, 1837); **Curculionidae:** *Anthonomus pomorum* (Linnaeus, 1758); *Hylesinus crenatus* (Fabricius, 1787); *H. fraxini* (Panzer, 1779); *H. oleiperda* Fabricius, 1792; **Dermestidae:** *Attagenus pello* (Linnaeus, 1758). **Scolytidae:** *Ips typographus* (Linnaeus, 1758); **DIPTERA, Tephritidae:** *Chaetostomella cylindrica* Robineau-Desvoidy, 1830; *Noeeta pupillata* (Fallen, 1814); **LEPIDOPTERA, Agonoxenidae:** *Scrobipalpa brahmiella* Heyden, 1862; **Coleophoridae:** *Coleophora laricella* (Huebner, 1817); **Gelechiidae:** *Anarsia spartiella* (Schrank, 1802) [*Ulex minor*]; *Caryocolum marmorea* Haworth, 1829; *Exoteleia dodecella* (Linnaeus, 1758); *Gelechia senticetella* (Staudinger, 1859); *Mirificarma mulinella* (Zeller, 1839); **Lymantriidae:** *Euproctis chrysorrhoea* (Linnaeus, 1758) [*Quercus pubescens*]; **Oecophoridae:** *Agonopterix subpropinquella* (Stainton, 1849); **Pyralidae:**

*Ephestia terebellum* Zeller, ?; **Tortricidae:** *Archips rosana* (Linnaeus, 1758); *Avaria hyperana* Milliére, 1857; *Choristoneura murinana* (Huebner, 1799); *Clepsis rogana* Guenée, 1845 [*Veratrum album*]; *Cnephasia asseclana* ([Denis & Schiffermüller], 1775); *C. chrysantheana* (Duponchel, 1843); *C. longana* (Haworth, 1811); *Cydia strobilella* (Linnaeus, 1758) [*Abies excelsa*]; *Epinotia nigricana* (Herrich-Schaffer, 1851); *Rhyacionia buoliana* (Denis & Schiffermüller, 1775); *R. pinivora* (Lienig & Zeller, 1846); *Tortrix viridana* Linnaeus, 1758; *Zeiraphera griseana* (Hübner, 1799); **Yponomeutidae:** *Yponomeuta malinella* Zeller, 1838.

**Distribution:** Holarctic, Oriental Cyprus (Papp, 1998).

**Genus: *Glyptomorpha* Holmgren, 1868**

**\**Glyptomorpha (Glyptomorpha) pectoralis* (Brullé, 1832)**

**Material examined:** Northern Cyprus - Akıncılar, wild herb, 09.04.2015, 1♀, 1♂; Beşparmak, Taşocağı, wheat, 08.04.2015, 5♀♀, 1♂; Ercan Hava alanı, wild herb, 09.04.2015, 4♀♀, 6♂♂; Girne, Çatalköy, wheat, 08.04.2014, 1♀, 1♂; Girne, Tepebaşı, wheat, 09.04.2015, 1♀, 2♂♂; İskele, wild herb, 07.04.2015, 2♀♀, 12♂♂; Lefkoşa, Devlet Üretim Çiftliği, wheat, 09.04.2015, 3♀♀, 2♂♂; Lefkoşa, Devlet Üretim Çiftliği, wheat, 09.04.2015, 1♀; Lefkoşa, Dikkaya, wild herb, 08.04.2015, 2♀♀, 6♂♂; Magosa, İnönü, barley, 09.04.2015, 4♀♀, 7♂♂; Magosa, Nergizli, wild herb, 09.04.2015, 1♀, 3♂♂; Serhatköy, Güzelyurt kavşağı, barley, 09.04.2015, 1♂; Türkmenköy, wild herb, 09.04.2015, 9♀♀, 2♂♂.

**Hosts: COLEOPTERA, Buprestidae:** *Chrysobothris affinis* (Fabricius, 1794); *Sphenoptera gossypii* Kerremans, 1892; *S. laticollis* (Olivier, 1790) [*Onobrychis viciifolia*]; *S. montana* Dejean, 1833; **Cerambycidae:** *Plagionotus arcuatus* (Linnaeus, 1758).

**Distribution:** Afrotropical, Oriental and Palaearctic, Cyprus (New record).

**Genus:** *Iphiaulax* Förster, 1862

**\**Iphiaulax fastidiator* (Fabricius, 1781)**

**Material examined:** Northern Cyprus - Mehmetcik, wild herb, 30.06.2016, 1♀, 2♂♂.

**Hosts:** COLEOPTERA, Curculionidae: *Saperda populnea* (Linnaeus, 1758).

**Distribution:** Afrotropical and Western Palearctic, Cyprus (New record).

**\**Iphiaulax jacobsoni* Shestakov, 1927**

**Material examined:** Northern Cyprus - Taşpınar wheat, 31.5.2012, 1♀, 2♂♂.

**Hosts:** Unknown.

**Distribution:** Eastern Palearctic, Oriental, Western Palearctic: Cyprus (New record).

**Genus:** *Pseudovipio* Szépligeti, 1896

***Pseudovipio castrator* (Fabricius, 1798)**

**Material examined:** Northern Cyprus - Türkmenköy, Karpaz, wild herb, 31.05.2016, 1♀.

**Hosts:** COLEOPTERA, Buprestidae: *Chrysobothris affinis* (Fabricius, 1794);

**Cerambycidae:** *Plagionotus arcuatus* (Linné, 1758); **Curculionidae:** *Lixus juncii* Boheman, 1835; **LEPIDOPTERA, Amphipyrae:** *Gortyna xanthenes* Germar, 1844 [*Cynara scolymus*].

**Distribution:** Afrotropical, Palearctic, Cyprus (Papp, 1998).

***Pseudovipio inceptor* (Nees, 1834)**

**Material examined:** Northern Cyprus - Türkmenköy, Karpaz, wild herb, 31.05.2016, 1♀.

**Hosts:** LEPIDOPTERA, Crambidae: *Ostrinia nubilalis* (Hübner, 1796).

**Distribution:** Palearctic, Cyprus (Papp, 1998).

**Subfamily:** Cheloninae

**Genus:** *Chelonus* Panzer, 1806

**Subgenus:** *Chelonus* Panzer, 1806

**\**Chelonus (Chelonus) mirandus* Tobias, 1964**

**Material examined:** Northern Cyprus - Geçitköy, wild herb, 01.06.2016, 2♂♂.

**Hosts:** Unknown.

**Distribution:** Palearctic, Cyprus (New record).

**\**Chelonus (Chelonus) oculator* (Fabricius, 1775)**

**Material examined:** Northern Cyprus - Akıncılar, wild herb, 09.04.2015, 1♀, 1♂; Devlet Üretim Çitliği, barley, 09.04.2015, 3♂♂; Dilekkaya, wild herb, 09.04.2015, 1♀, Girne, Çatalköy, wheat, 09.04.2015, 1♀; İnönü, wild herb, 09.04.2015, 1♀; İskele, wild herb, 07.04.2015, 1♀; Lefkoşa, Dikkaya, barley, 09.04.2015, 1♀; Serhatköy, wild herb, 09.04.2015, 1♀.

**Hosts:** LEPIDOPTERA, Coleophoridae: *Coleophora anatipennella* (Hübner, 1796);

**Crambidae:** *Loxostege sticticalis* (Linnaeus, 1761); *Ostrinia nubilalis* (Hübner, 1796);

**Noctuidae:** *Agrotis segetum* (Denis & Schiffermüller, 1775); *Helicoverpa armigera*

(Hübner, 1805); *Heliothis peltigera* Denis & Schiffermüller, 1775; *H. viroplaca* (Hufnagel,

1766); *Leucania loreyi* (Duponchel, 1827); *Longalatedes elymi* (Treitschke, 1825) (syn.: *Chortodes elymi*);

*Spodoptera exigua* (Hübner, 1808); *S. littoralis* Boisduval, 1833;

**Pyralidae:** *Ephestia kuehniella* Zeller, 1879; *Etiella zinckenella* (Treitschke, 1832);

*Homoeosoma nebulella* Denis & Schiffermüller, 1775; **Tortricidae:** *Cydia corticana* Hübner

(?), *Zeiraphera isertana* (Fabricius, 1794).

**Distribution:** Western Palearctic, Cyprus (New record).

**Subgenus:** *Microchelonus* Szépligeti, 1908

**\**Chelonus (Microchelonus) phalloniae* (Telenga, 1941)**

**Material examined:** Northern Cyprus - Serdarlı, wild herb, 30.05.2016, 1♂.

**Host:** Unknown.



**Distribution:** Eastern Palaearctic, Cyprus (New record).

**Subgenus:** *Stylochelonus* Hellén, 1958

\**Chelonus (Stylochelonus) pusillus* Szépligeti, 1908

**Material examined:** Northern Cyprus - Mehmetcik, wild herb, 31.05.2016, 1♀.

**Host:** LEPIDOPTERA, Elachistidae: *Elachista* sp.

**Distribution:** Palaearctic, Cyprus (New record).

**Subfamily:** Doryctinae

**Genus:** *Rhaconotus* Ruthe, 1854

\**Rhaconotus (Rhaconotus) testaceus* (Szépligeti, 1908)

**Material examined:** Northern Cyprus - Ercan, Balıkesir, wild herb, 30.05.2016, 1♀.

**Hosts:** LEPIDOPTERA, Crambidae: *Chilo suppressalis* (Walker, 1863); *Scirpophaga incertulas* (Walker, 1863).

**Distribution:** Oriental, Palaearctic, Cyprus (New record).

**Subfamily:** Euphorinae

**Genus:** *Meteorus* Haliday, 1835

*Meteorus rubens* (Nees, 1811)

**Material examined:** Northern Cyprus - Ercan, Balıkesir, wild herb, 30.05.2016, 1♀, 2♂♂; İskele, barley, 07.04.2015, 1♀; Köprüküy, wild herb, 30.05.2016, 1♂; Magosa, İnönü, wild herb, 08.04.2015, 1♀.

**Hosts:** DIPTERA, Chloropidae: *Meromyza americana* (Fitch, 1856) [*Triticum aestivum*];

LEPIDOPTERA, Coleophoridae: *Coleophora malivorella* (Riley, 1878);

Gelechiidae: *Pexicopia malvella* (Common, 1958);

*Phthorimaea operculella*; (Zeller, 1873);

Geometridae: *Idaea muricata*; (Hufnagel, 1767);

Lasiocampidae: *Eriogaster lanestris* (Linnaeus, 1758);

Lymantriidae: *Orgyia antiqua* (Linnaeus, 1758);

Noctuidae: *Agrotis exclamationis* (Linnaeus, 1758); *A. gladiaria* (Morrison, 1875); *A. ipsilon* (Hufnagel, 1766);

*A. malefida* (Guenée, 1852); *A. obesa* (Boisduval, 1829); *A. orthogonia* (Morrison, 1876); *A. segetum* (Denis & Schiffermüller, 1775); *A. subterranea* (Fabricius, 1794); *A. tokionis* (Butler, 1881); *A. vestigialis* (Hufnagel, 1766); *Anticarsia gemmatalis* (Hübner, 1818); *Apamea devastator* (Brace, 1819); *A. lateritia* (Hufnagel, 1766); *A. zeta* (Treitschke, 1825); *Autographa californica*; (Speyer, 1875); *A. gamma* (Linnaeus, 1758); *Autoplusia egea* (Guenée, 1852); *Celaena leucostigma* (Hübner, 1808); *Cerapteryx graminis* (Linnaeus, 1758); *Discestra trifolii* (Hufnagel, 1766); *Euxoa auxiliaris* (Grote, 1873); *E. messoria* (Harris, 1841); *E. ochrogaster* (Guenée, 1852); *E. perexcellens* (Grote, 1875); *E. temera* (Hübner, 1808); *E. tristicula* (Morrison, 1875); *E. tritici* (Linnaeus, 1761); *Feltia jaculifera* (Guenée, 1852); *F. subgothica* (Haworth, 1809); *Helicoverpa armigera* (Hübner, 1809); *Lacanobia oleracea* (Linnaeus, 1758), *Lithophane antennata* (Walker, 1858); *Mamestra brassicae* (Linnaeus, 1758); *Mesapamea stipata* (Morrison, 1875); *Noctua pronuba* (Linnaeus, 1758); *Ochropleura fennica* (Tauscher, 1806) (Syn. *Actebia fennica*); *Peridroma saucia* (Hübner, 1808); *Spodoptera exigua* (Hübner, 1808); *S. exigua* (Hübner, 1808); *S. frugiperda* (J.E. Smith, 1797); *S. littoralis* (Boisduval, 1833); *S. praefica* (Grote, 1875); *Syngrapha epigaea* (Grote, 1874); *Xestia c-nigrum* (Linnaeus, 1758); **Nymphalidae:** *Vanessa cardui* (Linnaeus, 1758); **Pieridae:** *Achyra bifidialis* (Fabricius, 1794); *Colias eurytheme* (Boisduval, 1852); *Hellula undalis* (Fabricius, 1794); *Nomophila noctuella* (Denis & Schiffermüller, 1775); *Omphalocera cariosa* (Lederer, 1863); *Pococera militella* (Zeller, 1848); **Thaumetopoeidae:** *Thaumetopoea wilkinsoni* (Tams, 1926); **Tortricidae:** *Acleris maccana* (Treitschke, 1835); *Eupoecilia ambiguella* (Hübner, 1796); *Lobesia botrana* (Denis & Schiffermüller, 1775); *Pandemis heparana* (Denis & Schiffermüller, 1775); *Tortrix viridana* (Linnaeus, 1758); **Yponomeutidae:** *Yponomeuta malinella* (Zeller, 1838).

**Distribution:** Holarctic, Neotropical, Oriental, Cyprus (Papp, 1990, 1998).

**\*Meteorus versicolor (Wesmael, 1835)**

**Material examined:** Northern Cyprus - Yeşilirmak, barley, 01.06.2016, 1♀.

**Hosts: LEPIDOPTERA, Arctiidae:** *Hyphantria cunea* (Drury, 1773); *Ocnogyna baetica* (Rambur, 1836); **Argyresthiidae:** *Argyresthia bonnetella* (Linnaeus, 1758); **Geometridae:** *Abraxas pantaria* (Linnaeus, 1767); *Archiearis parthenias sajana* (Prout, 1912); *Ematurga atomaria* (Linnaeus, 1758); *Ennomos erosaria* ([Denis & Schiffermüller], 1775); *Eulithis testata* (Linnaeus, 1761); *Eupithecia exigua* (Hübner, 1813); *Geometra papilionaria* (Linnaeus, 1758); *Operophtera brumata* (Linnaeus, 1758); *Protoarmia porcelaria* (Guenée, 1857); *Tephрина arenacearia* (Denis & Schiffermüller, 1775); *Thera juniperata* (Linnaeus, 1758); **Lasiocampidae:** *Dendrolimus pini* Linnaeus 1758; *D. spectabilis* (Butler 1877); *Eriogaster lanestris* (Linnaeus, 1758); *E. philippi* Bartel 1911 (*Quercus ithaburensis*); *Lasiocampa quercussicula* (Staudinger, 1861); *Macrothylacia rubi* Linnaeus, 1758; *Malacosoma castrense* (Linnaeus, 1758); *M. neustria* (Linnaeus, 1758); *M. parallelum* Staudinger 1887; *Bombyx lobulina* Denis & Schiffermüller 1775; **Lycaenidae:** *Callophrys rubi* (Linnaeus, 1758); **Lymantriidae:** *Arctornis l-nigrum* (Müller, 1764); *Calliteara pudibunda* (Linnaeus, 1758); *Euproctis chrysoorrhoea* (Linnaeus, 1758); *E. flava* Bremen, 1861; *Leucoma salicis* (Linnaeus, 1758) [*Populus grandidentata*, *P. tremuloides*]; *Lymantria dispar* (Linnaeus, 1758); *L. monacha* (Linnaeus, 1758); *Orgyia antiqua* (Linnaeus, 1758); *O. leucostigma* (Smith, 1797); *O. (Clethrogyna) recens* (Hubner, 1819); *O. thyellina* Butler [1979]; *Parocneria detrita* Esper, 1785; *Teia antiquioides* Hubner 1822; **Noctuidae:** *Agrotis exclamationis* (Linnaeus, 1758); *A. segetum* ([Denis & Schiffermüller], 1775); *Anarta myrtilli* (Linnaeus, 1761); *Autographa gamma* (Linnaeus, 1758); *Brachionycha sphinx* (Hufnagel, 1766; *Cosmia*

*trapezina* (Linnaeus, 1758); *Diachrysia chrysitis* (Linnaeus, 1758); *Noctua porphyrea* Denis & Schiffermüller, 1775; *N. pronuba* (Linnaeus, 1758); *Nycteola asiatica* (Krulikovsky, 1904); *Orthosia carnipennis* Butler, 1878; *O. miniosa* (Denis & Schiffermüller, 1775); *Panolis flammea* ([Denis & Schiffermüller], 1775); *Xestia agathina* (Duponchel, 1827); *Nola cucullatella* (Linnaeus, 1758); **Notodontidae:** *Clostera anastomosis* (Linnaeus, 1758); **Nymphalidae:** *Maniola jurtina* Linnaeus, 1758; **Pieridae:** *Pieris rapae* (Linnaeus, 1758); **Pyralidae:** *Ercta orientalis* Yamanaka, 1972 [*Ipomoea paniculata*]; *Eurrhynpara hortulata* (Linnaeus, 1758); **Thaumetopoeidae:** *Thaumetopoea pityocampa* Denis & Schiffermüller, 1775 [*Pinus halepensis*]; *T. processionea* (Linnaeus, 1758); *T. wilkinsoni* Tams, 1924 [*Pinus halepensis*]; **Tortricidae:** *Archips oporana* (Linnaeus, 1758); *Pammene inquilina* Fletcher, 1938; *Pandemis cerasana* (Hübner, 1786).

**Distribution:** Holarctic, Oriental, Cyprus (New record).

**Subfamily: Homolobinae**

**Genus: Homolobus Förster, 1862**

**Subgenus: Apatia Enderlein, 1920**

**Homolobus (Apatia) truncator (Say, 1829)**

**Material examined:** Northern Cyprus - Ercan Hava alanı, Deneme, wild herb, 09.04.2015, 1♀; Ercan, Balıkesir, barley, 30.05.2016, 1♀; 09.04.2015, 2♀♀; Lefkoşa, Hamitköy, wheat, wild herb, 28.03.2012, 2♀♀; Lefkoşa, Devlet üretme çitliği, wheat, 09.04.2015, 1♀.

**Hosts: LEPIDOPTERA, Gelechiidae:** *Phthorimaea operculella* (Zeller, 1873); **Geometridae:** *Agriopsis bajaria* (Denis & Schiffermüller, 1775); *Anisodes lyciscaria* (Guenée, 1858); *Ascotis selenaria* (Denis & Schiffermüller, 1775); *Besma quercivoraria* (Guenée, 1857); *Casilda consecraria* (Rambur, 1858); *Chiasmia clathrata* (Linnaeus, 1758); *Glossotrophia annae* (Mentzer, 1990); *G. asellaria* (Herrich-Schäffer, 1847); *G.*

(Graslin, 1863); *Hypagyrtis piniata* (Packard, 1870); *Lambdina feroidaria* (Hübner, 1827); *Lambdina fiscellaria* (Guenée, 1857); *Lycia hirtaria* (Clerck, 1759); *L. zonaria* (Dennis & Schiffermüller, 1775); *Pachycnemia hippocastanaria* (Hübner, 1799); *Rhodometra sacraria* (Linnaeus, 1767); **Lasiocampidae:** *Malacosoma neustria* (Linnaeus, 1758); **Noctuidae:** *Agrotis ipsilon* (Hufnagel, 1766); *A. orthogonia* (Morrison, 1876); *A. segetum* (Denis & Schiffermüller, 1775); *A. subterranea* (Fabricius, 1794); *A. venerabilis* (Walker, 1857); *Anumeta cestina* (Staudinger, 1884); *Autographa gamma* (Linnaeus, 1758); *Calophasia platyptera* (Esper, 1788); *Catocala nymphagoga* (Esper, 1787); *Emmelia trabealis* (Scopoli, 1763); *E. tristicula* (Morrison, 1875); *Helicoverpa zea* (Boddie, 1850); *H. zea* (Boddie, 1850) [*Medicago sativa*]; *Lacanobia contigua* (Denis & Schiffermüller, 1775); *Mocis latipes* (Guenée, 1852); *M. latipes* (*Cynodon dactylon*) (Guenée, 1852); *Panolis flammea* (Denis & Schiffermüller, 1775); *Protorthodes smithii* (Dyar, 1904); *Pseudaletia unipuncta* (Haworth, 1809) [*Medicago sativa*]; *Rhyacia simulans* (Hufnagel, 1766); *Spodoptera exigua* (Hübner, 1808) [*Asparagus setaceus*, *Medicago sativa*]; *S. frugiperda* (J.E. Smith, 1797) [*Medicago sativa*, *Sorghum bicolor*]; *S. litura* (Fabricius, 1775); *S. ornithogalli* (Guenée, 1852); *S. ornithogalli* (Guenée, 1852) [*Medicago sativa*]; *Xestia c-nigrum* (Linnaeus, 1758); *X. smithii* (Snellen, 1896); **Pyralidae:** *Achyra rantalis* (Guenée, 1854); *Loxostege sticticalis* (Linnaeus, 1761).

**Distribution:** Afrotropical, Holarctic, Neotropical, Oriental, Cyprus (Papp, 1998).

**Subgenus: *Phylacter* Reinhard, 1863**

\**Homolobus (Phylacter) annulicornis* (Nees, 1834)

**Material examined:** Northern Cyprus - Magosa, İnönü, wild herb, 09.04.2015, 1♂.

**Hosts:** LEPIDOPTERA, **Noctuidae:** *Autographa gamma* (Linnaeus, 1758); *Cosmia diffinis* Linnaeus, 1767 [*Ulmus carpinifolia*];

*C. trapezina* (Linnaeus, 1758); *Dichonia convergens* (Denis & Schiffermüller, 1775); *Dryobotodes eremita* (Fabricius, 1775); *Fissipunctia ypsilon* Denis & Schiffermüller, 1775; *Lithophane lamda* (Fabricius, 1787); *Mamestra brassicae* (Linnaeus, 1758); *Orthosia populeti* Fabricius, 1775; *Panolis flammea* Denis & Schiffermüller, 1775; *Xestia triangulum* (Hufnagel, 1766).

**Distribution:** Oriental, Western Palearctic, Cyprus (**New record**).

**Subfamily: Opiinae**

**Genus: *Biosteres* Förster, 1862**

\**Biosteres (Chilotrachia) arenarius* (Stelfox, 1959)

**Material examined:** Northern Cyprus - Girne Çatalköy, barley, 8.4.2015, 1♀.

**Hosts:** Unknown.

**Distribution:** Palearctic, Cyprus (**New record**).

**Genus: *Opius* Wesmael, 1835**

**Subgenus: *Allotypus* Förster, 1862**

\**Opius (Allotypus) arenaceus* Jakimavicius, 1986

**Material examined:** Northern Cyprus - Girne, Çatalköy, wild herb, 08.04.2015, 1♀.

**Hosts:** Unknown.

**Distribution:** Western Palearctic, Cyprus (**New record**).

**Subgenus: *Gastrosema* Fischer, 1972**

\**Opius (Gastrosema) caucasi* Tobias, 1986

**Material examined:** Northern Cyprus - Beşparmak, Taşocağı, wheat, 08.04.2015, 1♀; İskele, 07.04.2015, wheat, 1♀.

**Host:** Diptera, **Agomyzidae:** *Phytomyza horticola* Goureau, 1851.

**Distribution:** Palearctic, Cyprus (**New record**).

**Subgenus: *Nosopoea* Förster, 1862**

\**Opius (Nosopoea) sigmodus* Papp, 1981

**Material examined:** Northern Cyprus - Girne tepebaşı, wild herb, 09.04.2015, 1♀.

**Hosts:** Unknown.

**Distribution:** Western Palaearctic.

**\**Opius (Nosopoea) speciosus* Fischer, 1959**

**Material examined:** Northern Cyprus - Lefkoşa, Devlet üretim çiftliği, wild herb, 09.04.2015, 1♂.

**Hosts:** Unknown.

**Distribution:** Western Palaearctic, Cyprus (New record).

**Subgenus: *Pendopius* Fischer, 1972**

**\**Opius (Pendopius) bajariae* Fischer, 1989**

**Material examined:** Northern Cyprus Türkmenköy, wild herb, 09.04.2015, 1♂.

**Hosts:** Unknown.

**Distribution:** Western Palaearctic, Cyprus (New record).

**Subfamily: Rogadinae**

**Genus: *Aleiodes* Wesmæl, 1838**

**Subgenus: *Aleiodes* Wesmæl, 1838**

**\**Aleiodes (Aleiodes) bicolor* (Spinola, 1808)**

**Material examined:** Northern Cyprus - Köprüküy, wheat, 30.05.2016, 1♀.

**Hosts:** Unknown.

**Distribution:** Palaearctic, Cyprus (New record).

**Subgenus: *Chelonorhogas* Enderlein, 1912**

**\**Aleiodes (Chelonorhogas) eurinus* (Telenga, 1941)**

**Material examined:** Northern Cyprus - İskele, wild herb, 07.04.2015, 1♂.

**Hosts: LEPIDOPTERA, Noctuidae: *Apamea anceps* (Denis & Schiffermüller, 1775); *E. ochrogaster* (Guenée, 1852).**

**Distribution:** Oriental, Palaearctic, Cyprus (New record).

**Subgenus: *Neorhogas* Szépligeti, 1906**

***Aleiodes (Neorhogas) ductor* (Thunberg, 1822)**

**Material examined:** Northern Cyprus - Beşparmak dağları, taşocağı, wild herb,

09.04.2015, 1♀, 4♂♂; Köprüküy, wheat, 30.05.2016, 1♀.

**Hosts: LEPIDOPTERA, Lasiocampidae:**

*Euthrix potatoria* (Linnaeus, 1758); *Anarta (Calocestra) trifolii* (Hufnagel 1766); *Autographa gamma* (Linnaeus, 1758); *Mamestra brassicae* (Linnaeus 1758);

**Nymphalidae:** *Brenthis ino* (Rottemburg 1775); **Sesiidae:** *Synanthedon scoliaeformis* (Borkhausen 1789).

**Distribution:** Palaearctic, Cyprus (Papp, 1998).

### Discussion

The study was conducted in 108 localities covering the north of Cyprus. A total of 42 species belonging to 14 genera of Agathidae, Braconinae, Cheloninae, Doryctinae, Euphorinae, Homolobinae, Opiinae and Rogadinae are reported from Northern Cyprus. Twenty-six species are recorded from Cyprus for the first time. The hosts of 15 species are known.

Due to its geographical location, Braconidae species of Cyprus are very common (Table. 1). Distribution of Braconid species depends to the host species. 84% of Cyprus fauna restricted to the Western Palaearctic, 9% in both Palaearctic and Oriental, 4% Holarctic, 2.4% Nearctic, Neotropical, Oriental and Palaearctic. A small number of species have been recorded from a large number of zoogeographic regions. The most common species in the zoogeographic regions was *Homolobus (Apatia) truncator*. It has been recorded from Afrotropical, Holarctic, Neotropical and Oriental region and has about 55 host species. While *Glyptomorpha (Glyptomorpha) pectoralis* and *Bracon (Rostrobracon) urinator* are recorded from Afrotropical, Oriental and Palaearctic regions. It was the most common species of north Cyprus with 12 locality records. The other most common species is *Meteorus rubens*. It has been recorded from Holarctic, Neotropical and Oriental regions.

**Table 1.** Distribution of species from Cyprus based on biogeographic regions.

Zoogeographical regions	Number of species
Afrotropical, Australasian, Oceanic, Oriental, Eastern Palaearctic.	1
Afrotropical, Australasian, Oriental, Holarctic.	1
Afrotropical, Australasian, Nearctic, Neotropical Oceanic, Oriental, Palaearctic.	1
Afrotropical, Oceanic, Western Palaearctic.	1
Afrotropical, Oceanic, Oriental, Palaearctic.	1
Afrotropical, Nearctic, Neotropical, Oceanic, Oriental, Palaearctic.	1
Afrotropical, Nearctic, Neotropical, Oriental, Palaearctic.	1
Afrotropical, Nearctic, Neotropical, Palaearctic.	1
Afrotropical, Oriental, Palaearctic.	2
Afrotropical, Palaearctic.	2
Afrotropical, Holarctic, Oceanic.	1
Australasian, Nearctic, Neotropical, Oceanic, Oriental, Palaearctic.	1
Australasian, Oceanic, Oriental, Palaearctic.	1
Australasian, Nearctic, Neotropical, Oceanic, Palaearctic.	1
Holarctic.	5
Holarctic, Oriental and introduced into USA (Oregon).	1
Nearctic, Eastern Palaearctic.	1
Nearctic, Neotropical, Oceanic, Oriental, Palaearctic.	1
Nearctic, Neotropical, Oceanic, Palaearctic.	1
Nearctic, Neotropical, Oriental, Palaearctic.	3
Neotropical, Western Palaearctic.	1
Oceanic, Palaearctic.	1
Oriental, Palaearctic.	11
Palaearctic.	64
Western Palaearctic.	19
<b>Total</b>	<b>124</b>

The geographical land nearest to Cyprus is Anatolia. The braconidae fauna of Cyprus is similar to the braconidae fauna of Anatolia in accordance with the island biogeography theory (Aydoğdu & Beyarslan, 2002, 2006a, 2006b, 2006c, 2009; Beyarslan, 1996, 2011, 2014, 2015a, 2015b, 2015c, 2016; Beyarslan & Aydogdu, 2012, 2013; Beyarslan & Çetin Erdoğan, 2011; Beyarslan & Fischer 2013; Beyarslan et al., 2013; Çetin Erdoğan & Beyarslan, 2016; Çetin Erdoğan et al., 2009, Fischer & Beyarslan, 2011, 2012, 2013).

#### Acknowledgments

We would like to thank the administrators of Bitlis Eren and Iğdır Universities.

#### Conflict of Interests

The authors declare that there is no conflict of interest regarding the publication of this paper.

#### References

- Achterberg, C. van (2003) The West Palaearctic Species of the genera *Gildoria* Hedqvist and *Platyspathius* Viereck, with Keys to the Species (Hymenoptera: Braconidae: Doryctinae). *Zoologische Mededelingen (Leiden)*, 77 (15), 267–290.
- Anonymous (2004) The Geological Structure of Cyprus. Available from: [http://www.cyprusgeology.org/english/EN/2\\_2\\_geology\\_en.htm](http://www.cyprusgeology.org/english/EN/2_2_geology_en.htm). [Accessed 10th September 2017].

- Anonymous (2015) The Geological Structure of Cyprus. Available from: [http://www.cyprusgeology.org/english/EN/2\\_2\\_geology\\_en.htm](http://www.cyprusgeology.org/english/EN/2_2_geology_en.htm). [Accessed 25th September 2017].
- Aydoğdu, M. & Beyarslan, A. (2002) The *Chelonus* Jürine (Hymenoptera: Braconidae: Cheloninae) Species of the Marmara Region. *Turkish Journal of Zoology*, 26 (1), 1–13.
- Aydogdu, M. & Beyarslan, A. (2006a) *Ascogaster disparilis* Tobias 1986 (Hymenoptera: Braconidae: Cheloninae), the second record for Palaearctic region from Turkey. *Linzer Biologische Beiträge*, 38 (1), 393–396.
- Aydogdu, M. & Beyarslan, A. (2006b) *Microchelonus Szepligeti* 1908 (Hymenoptera: Braconidae: Cheloninae) species from Marmara, Western and Blacksea regions of Turkey, *Linzer Biologische Beiträge*, 38 (1), 397–407.
- Aydogdu, M. & Beyarslan, A. (2006c) First records of *Aleiodes* Wesmael, 1838 species in East Marmara Region of Turkey (Hymenoptera: Braconidae: Rogadianae), *Acta Entomologica Slovenica*, 14 (1), 81–88.
- Aydoğdu, M. & Beyarslan, A. (2009) A review of the Phanerotomini (Hymenoptera: Braconidae: Cheloninae) in Turkey, with a new host record for *Phanerotoma (Bracotritoma) permixtella*. *Biologia*, 64 (4), 748–756.  
<https://doi.org/10.2478/s11756-009-0126-3>
- Beyarslan, A. (1996) Die *Orgilus*-Arten der Türkei (Hymenoptera, Braconidae, Orgilinae). *Entomofauna*, 17 (22), 353–360.
- Beyarslan, A. (2011) *Orgilus (Orgilus) elazigensis* sp n. from Eastern Turkey (Hymenoptera: Braconidae: Orgilinae). *Zoology in the Middle East*, 53, 95–98.  
<https://doi.org/10.1080/09397140.2011.10648866>
- Beyarslan, A. (2014) Checklist of Braconinae species of Turkey (Hymenoptera: Braconidae). *Zootaxa*, 3790 (2), 201–242.
- Beyarslan, A. (2015a) A faunal study of the subfamily Doryctinae in Turkey (Hymenoptera: Braconidae). *Turkish Journal of Zoology*, 39, 126–143.  
<https://doi.org/10.3906/zoo-1402-38>
- Beyarslan, A. (2015b) Taxonomical investigations on the fauna of Opiinae (Hymenoptera: Braconidae) in Ardahan, Erzurum, Iğdır, and Kars provinces of the Turkish northeastern Anatolian region. *Turkish Journal of Zoology*, 39, 820–831.  
<https://doi.org/10.3906/zoo-1411-33>
- Beyarslan, A. (2015c) Taxonomic survey on the Rogadinae Förster, 1862 (Hymenoptera, Braconidae) in the northeastern Anatolian region, Turkey. *Turkish Journal of Zoology*, 39, 811–819.  
<https://doi.org/10.3906/zoo-1407-35>
- Beyarslan, A. (2016) Taxonomic investigations on the Braconinae fauna (Hymenoptera, Braconidae) in north-eastern Anatolian region, Turkey, with the description of a new species. *Zootaxa*, 4079 (1), 001–033.  
<https://doi.org/10.11646/zootaxa.4079.1.1>
- Beyarslan, A. & Aydogdu, M. (2012) A Preliminary Study of the *Macrocentrus* Curtis, 1833 (Hymenoptera: Braconidae: Macrocentrinae) fauna of Turkey, with Zoogeographical Remarks. *Journal of the Entomological Research Society*, 14 (1), 83–90.
- Beyarslan, A. & Aydogdu, M. (2013) Additions to The rare species of Braconidae fauna (Hymenoptera: Braconidae) from Turkey. *Munis Entomology & Zoology*, 8 (1), 369–374.
- Beyarslan, A. & Cetin Erdoğan, Ö. (2011) A study of Orgilinae and Microtypinae (Hymenoptera: Braconidae) from Turkey with the description of a new species. *Biologia*, 63 (1), 121–129.  
<https://doi.org/10.2478/s11756-010-0148-x>
- Beyarslan, A. & Fischer, M. (2013) Checklist of Turkish Opiinae (Hymenoptera, Braconidae). *Zootaxa*, 3721 (5), 401–454.  
<https://doi.org/10.11646/zootaxa.3721.5.1>
- Beyarslan, A., Gözüaçık, C. & Özgen, I. (2013) A contribution on the subfamilies Helconinae, Homolobinae, Macrocentrinae, Meteorinae, and Orgilinae (Hymenoptera: Braconidae) of southeastern Anatolia with new records from other parts of Turkey. *Turkish Journal of Zoology*, 37, 501–505.  
<https://doi.org/10.3906/zoo-1301-10>
- Çetin Erdoğan, Ö., Achterberg, C. van & Beyarslan, A. (2009) On the zoogeographical Distribution of the genus *Agathis* Latreille, 1804 Hymenoptera: Braconidae: Agathidinae) in Turkey. *Journal of Entomological Research Society*, 11 (1), 17–25.

- Çetin Erdoğan, Ö. & Beyarslan, A. (2016) Faunistic survey on the Agathidinae (Hymenoptera: Braconidae) in the Central Anatolia Region with a new record for the Turkish fauna: *Bassus tegularis* (Thomson, 1895). *Turkish Journal of Zoology*, 40, 448–453. <https://doi.org/10.3906/zoo-1510-42>
- Fischer, M. (1999) Zur Evolution und zum System der Opiusverwandten Gattungen der Unterfamilie Opiinae mit einer erweiterten Aufteilung dieses Gattungs-Komplexes (Hymenoptera, Braconidae, Opiinae). *Linzer Biologische Beiträge*, 31, 277–336.
- Fischer, M. & Beyarslan, A. (2011) *Indiopijs yilmazae* (Hymenoptera: Braconidae: Opiinae), a new species from Turkey. *Arbeitsgemeinschaft Österreichischer Entomologen*, 63, 123–126.
- Fischer, M. & Beyarslan, A. (2012) New species of *Synaldis* Foerster and *Idiasta* Foerster, and further records of Turkish Alysini (Hymenoptera: Braconidae: Alysini). *Zoology in the Middle East*, 53, 41–60.
- Fischer, M. & Beyarslan, A. (2013) Additional contributions to the Opiinae fauna of Turkey (Hymenoptera: Braconidae). *Turkish Journal of Zoology*, 37, 525–538. <https://doi.org/10.3906/zoo-1204-9>
- Marsh, P.M. & Shaw S.R. (2003) Revision of North American *Aleiodes* Wesmael (Part 7): the compressor *Herrich-Schaeffer*, *ufei* (Walley), *gressetti* (Muesebeck), and *procerus* (Wesmael) species-groups (Hymenoptera: Braconidae: Rogadinae). *Proceedings of the Entomological Society of Washington*, 105, 698–707.
- Matthews, R.W. (1970) A revision of the genus *Spathius* in America north of Mexico (Hymenoptera: Braconidae). *Entomological Society of America*, 4, 1–86.
- Nixon, G.E.J. (1986) A revision of the European Agathidinae (Hymenoptera: Braconidae). *Bulletin of the British Museum (Natural History), Entomology series*, 52 (3), 183–242.
- Papp, J. (1982) Braconidae (Hymenoptera) from Greece, 1. *Annales Musei Goulandris*, 5, 183–193.
- Papp, J. (1990) Braconidae (Hymenoptera) from Greece, 3. *Annales Musei Goulandris*, 8, 269–290.
- Papp, J. (1998) Contributions to the braconid fauna of Cyprus (Hymenoptera, Braconidae: Braconinae). *Entomofauna*, 19 (14), 241–251.
- Papp, J. (1999) A revision of the *Bracon* species described by O. Schmiedeknecht (Insecta: Hymenoptera: Braconidae: Braconinae). *Entomologische Abhandlungen*, 58 (16), 89–308.
- Shaw, M.R. & Huddleston T. (1991) Classification and Biology of Braconid Wasps (Hymenoptera: Braconinae). Handbooks for the Identification of British Insects, Vol. 7, Part 11. *Royal Entomological Society of London, England*, 126 pp.
- Shenefelt, R.D. (1978) Braconidae 10. Hym. Cat. (n. ed.) pars 15, 1425–1872.
- Tobias, V.I. (1986) *Keys to the Insects of the European Part of the USSR*. Vol. 3. Hymenoptera. Part 4. [Opredelitel Nasekomych Evrospeiskoi Tsasti SSSR 3, Peredpontdatokrylye 4. Opr. Faune SSSR] *Nauka publisher, Leningrad*, 156–512 pp. [English translation. Lebanon, U.S.A.].
- Wang, Y.P., Chen, X.X., Wu, H. & He, J.H. (2010) Two genera of Braconinae (Hymenoptera, Braconidae) in China, with descriptions of four new species. *ZooKeys*, 61, 47–62. <https://doi.org/10.3897/zookeys.61.450>
- Yu, D. S., Achterberg C. van & Horstmann K. (2012) Interactive Catalogue of World Ichneumonoidea, Taxonomy, Biology, Morphology, and Distribution. Compact disc



مطالعه تاکسونومیک فون زنبورهای خانواده Braconidae (Hymenoptera: Ichneumonoidea) در شمال قبرس

احمد بایرسلان<sup>۱\*</sup>، سلالتین گوزواچیک<sup>۲</sup>، مصطفی گولو<sup>۳</sup> و آیدا کنوکسال<sup>۴</sup>

۱ گروه زیست‌شناسی، دانشکده هنر و علوم، دانشگاه ارن بیتلیس، ترکیه.

۲ گروه گیاهپزشکی، دانشکده کشاورزی، دانشگاه ایکدیر، ترکیه.

۳ موسسه تحقیقات کنترل بیولوژیک، آدانا، ترکیه.

۴ موسسه تحقیقات کشاورزی، لفکوسا، جمهوری ترکی شمال قبرس.

\* پست الکترونیکی نویسنده مسئول مکاتبه: [abeyars@gmail.com](mailto:abeyars@gmail.com)

تاریخ دریافت: ۲ مهر ۱۳۹۶، تاریخ پذیرش: ۲۹ مهر ۱۳۹۶، تاریخ انتشار: ۲ آبان ۱۳۹۶

**چکیده:** به منظور شناسایی فون زنبورهای براکونیده، نمونه‌های حشرات کامل این زنبورها از زیستگاه‌های مختلف در شمال قبرس بین سال‌های ۲۰۱۲ تا ۲۰۱۶ جمع‌آوری شد. نمونه‌برداری در هر دو گروه زیستگاه‌های طبیعی و مناطق کشاورزی با استفاده از روش تورجارو انجام شد. به طور کلی ۴۲ گونه متعلق به ۴ جنس از زیرخانواده‌های Agathidinae, Braconinae, Cheloninae, Doryctinae, Euphorinae, Homolobinae, Opiinae و Rogadinae از این منطقه گزارش شدند که از بین آنها ۲۶ گونه برای اولین بار گزارش می‌شوند.

**واژگان کلیدی:** کوه پنج‌دندانه، براکونیده، مجموعه جانوری، قبرس، پارازیتوئید