



Received:  
19 June, 2017

Accepted:  
23 August, 2017

Published:  
28 August, 2017

Subject Editor:  
Farzaneh Kazerani

## Study of the robber flies (Diptera: Asilidae) in East and West Azerbaijan provinces of Iran, with two new species records for the country

Rahman Mohammadi<sup>1\*</sup>, Samad Khaghaninia<sup>1</sup> and Dmitry Astakhov<sup>2</sup>

<sup>1</sup> Faculty of Agriculture, Department of Plant Protection, University of Tabriz, Tabriz 51664, Iran.

<sup>2</sup> Volgograd State University, 100 Universitetskiy pr., Volgograd 400062, Russia.

**ABSTRACT.** Based on specimens collected from East Azerbaijan and West Azerbaijan provinces during 2014–2016, fourteen species of the family Asilidae are collected and identified. Two of them, e.g. *Dysmachus transcaucasicus* Richter, 1962; *Saropogon megriensis* Richter, 1966 are recorded from Iran for the first time. All species are listed along with their geographic distributions. Diagnostic characters as well as supplementary illustrations of new records are provided.

**Key words:** Asilidae, East Azerbaijan, West Azerbaijan, Iran, New records.

**Citation:** Mohammadi, R., Khaghaninia, S. & Astakhov, D. (2017) Study of the robber flies (Diptera: Asilidae) in East and West Azerbaijan provinces of Iran, with two new species records for the country. *Journal of Insect Biodiversity and Systematics*, 3 (3), 247–255.

### Introduction

The robber flies (Diptera: Asilidae) is a large family with 7187 described species (Geller-Grimm et al., 2015) and 776 genera in the world (Dikow, 2016). The comprehensive study by Dikow (2009) proposes a revised, phylogenetic classification of Asilidae into 14 subfamily taxa. Robber flies are small to very large (3–50 mm) flies, often with conspicuous dense hair or almost bald. Eyes are large, dichoptic in both sexes, separated by markedly deepened vertex. All of them are very good flyers with long legs which serve for catching preys (Major, 1997). All robber flies are predators in larval and adult stages. Females in the subfamily Stenopogoninae and Dasypogoninae deposit their eggs into soil. While females of the genera *Eutolmus* and *Dysmachus* lays their

eggs on grass or other plant leaves. However, *Laphria* and *Choerades* females deposit their eggs in front of holes of xylophagus insects and cracks in bark (Lehr, 1988; Weinberg & Bachli, 1995). The majority of the larvae live in soil but those of the Laphriinae and Laphystiinae occur in decaying logs and stumps, where they feed on larvae and pupae of other insects (Geller-Grimm, 2002). The majority of species of Asilidae frequently observed in dry and sandy areas, a condition well shown by the greater numbers of species found in arid and semiarid regions, but even in desert or semi-desert countries the small drains of dry steam beds attract the greatest number, and sometimes the entire robber fly population of a region will be restricted to such places,

Corresponding author: Rahman Mohammadi, E-mail: [Rahmanmohamadi@gmail.com](mailto:Rahmanmohamadi@gmail.com)

**Copyright** © 2017 Mohammadi et al. This is an open access article distributed under the terms of the Creative Commons Attribution License (CC BY 4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

which also have the maximum vegetation and the greatest population of insects upon which the flies feed (Hull, 1962). Investigations on Asilidae in Iran are strongly restricted and have been conducted principally by non-Iranian researchers. Portschinsky (1873) described 2 new species from Iran. Bigot (1880), Hermann (1905), Becker and Stein (1913), Engel (1930), Oldroyd (1958), Janssens (1961), Abbassian-Lintzen (1964a, b), Tsacas (1968), and Theodor (1980) contributed to the knowledge of Iranian fauna. Timon-David (1955), Hradsky and Geller-Grimm (1998), Tomasovic (1999, 2002) were described some new species from Iran. So far 242 species of Asilidae were identified from Iran (Lehr et al., 2007; Hayat et al., 2008; Saghaei et al., 2008, and Mohammadi & Khaghaninia 2016a, 2016b).

#### Material and methods

Individual flies were collected using handy entomological net from the various regions of the East Azerbaijan and West Azerbaijan provinces of Iran during 2014–2016. To dissect the male genitalia, the end of the abdomen was removed and boiled in KOH solution, and then placed in acetic acid. The genitalia were washed and then stored in glycerine and examined under a stereo microscope. Distribution of species mostly follows Geller-Grimm et al., 2015, with minor additions and corrections. Morphological terminology follows that of Bei-Bienko G. (1988) and Geller-Grimm (2003, 2015). Collected specimens are deposited at the Collection of Dr. H. Maleki Milani, University of Tabriz, Iran (ICHMM) and the third author's personal collection.

#### Results

In this study, fourteen species belong to eleven genera were identified. Of them, two species *Dysmachus transcausicus* Richter, 1962 and *Saropogon megriensis* Richter, 1966

are being newly reported for the Iranian fauna which are marked with an asterisk.

#### The list of species

##### 1. *Anisopogon hermanni* (Engel, 1930)

**Material examined:** East Azerbaijan province, Kandovan, (37° 46.10' N, 46° 16.001' E) 2500m, 25.6.2015, 2♂♂, leg. S. Khaghaninia.

**Iranian Records:** Fars (Lehr et al., 2007).

**Distribution:** Israel, Kazakhstan, former South European territory, former Transcaucasus Republics, Turkey (Geller-Grimm et al., 2015).

##### 2. *Didysmachus picipes* (Meigen, 1820)

**Material examined:** East Azerbaijan province, Kandovan, (37° 44.200' N, 46° 18.001' E) 3090m, 20.5.2015, 1♂; West Azerbaijan province, Ziveh, (37°08' N, 44°52' E) 2630 m, 26.06.2015, 2♂♂, leg. S. Khaghaninia.

**Iranian Records:** Fars and Yazd (Lehr et al., 2007), West Azarbaijan (Shoeibi & Karimpour, 2010).

**Distribution** Albania, Armenia, Austria, Azerbaijan, Czech Republic, Denmark, Finland, France, Georgia, Germany, Hungary, Italy, Kazakhstan, Norway, Poland, Romania, Russia (North, Central and South European territory, West Siberia), Sweden, Switzerland, The Netherlands, Turkey, Ukraine, former Yugoslavia (Geller-Grimm et al., 2015).

##### 3. *Dysmachus bilobus* Loew, 1871

**Material examined:** East Azerbaijan province, Qaradagh forests, (38°55.601' N, 46°46.932' E) 1313 m, 20.6.2015, 2♂♂; Qurigol, (37° 55.305' N, 46° 41.542' E) 1928 m, 8.7.2015, 2♂♂, leg. S. Khaghaninia.

**Iranian Records:** Golestan (Lehr et al., 2007).

**Distribution:** Armenia, Azerbaijan, Bulgaria, Georgia, Kirgystan, Romania, Russia (South European territory), Turkey, Ukraine, Uzbekistan, former Yugoslavia (Geller-Grimm et al., 2015).

#### 4. *Dysmachus praemorsus* (Loew, 1854)

**Material examined:** East Azerbaijan province, Arasbaran, (46° 26.536'N, 44° 54.041' E) 1534m, 21.6.2015, 1♂, 1♀; Horand, (38° 53.838' N, 47° 16.988' E) 1367m, 20.6.2015, 1♂, 1♀; Chichekli, (38° 37.169' N, 46° 26.536' E) 1534m, 20.06.2015, 2♂♂; Kandovan, (37° 46.985' N, 46°15.686' E) 2341m, 24.07.2016, 2♂♂, 1♀; Kandovan, )37° 44.001' N, 46° 19.010'( 3000m, 24.07.2016, 1♂, leg. S. Khaghaninia.

**Iranian Records:** East Azarbaijan (Shoeibi and Karimpour, 2010), Fars (Lehr et al., 2007).

**Distribution:** Albania, Austria, Bulgaria, France, Germany, Hungary, Poland, Romania, Turkey, Ukraine, former Yugoslavia (Geller-Grimm et al., 2015).

#### 5. *Dysmachus transcaucasicus* Richter, 1962\*

**Material examined:** East Azerbaijan, Chichekli, (38° 37.169' N, 46° 26.536' E) 1534m, 21.6.2015, 1♂; Horand, (38° 53.838' N, 47° 16.988' E) 1367m, 20.6.2015, 2♂♂, leg. S. Khaghaninia.

**Distribution:** Armenia, Turkey (Geller-Grimm et al., 2015). Iran (**New record**).

**Diagnostic characters:** Mesopleuron without bristles or with inconspicuous weak ones only; dc and ac well developed, usually reaching level of humeral calli; Legs black (Fig. 1); fore femur on lower side with well developed bristle (Fig. 2); Abdomen and abdominal segments shortened and robust, apex of abdomen turned upward; Sternites with strong bristles at least on segments 2–4; Terminalia small; Two halves of epandrium without notch along middle margin on upper side; sternit 8 of female membranous along lower margin in apical half; Body 13 to 18 (Bei-Bienko, 1988 and Major, 2000) (Figs 4–8).

#### 6. *Laphria aurea* (Fabricius, 1794)

**Material examined:** East Azerbaijan province, Chichekli, (38° 37.169' N, 46° 26.536' E) 1534m, 1♂, leg. S. Khaghaninia.

**Iranian Records:** Mazandaran, Qom (Ghahari et al., 2007).

**Distribution:** Austria, Bulgaria, Czech Republic, France, Greece, Hungary, Israel, Italy, Romania, Russia (South European territory), Slovakia, Turkey, former Yugoslavia (Geller-Grimm et al., 2015).

#### 7. *Machimus annulipes* (Brullé, 1832)

**Material examined:** East Azerbaijan province, Horand, (38° 53.838' N, 47° 16.988' E) 1367m, 21.8.2015, 1♂; West Azerbaijan province, Urmieh, (36° 58.01' N, 45° 23.832' E), 26.6.2015, 3♀♀, leg. S. Khaghaninia.

**Iranian Records:** Fars (Saghaei et al., 2010), Kermanshah, Semnan (Hayat et al., 2008).

**Distribution:** Albania, Azerbaijan, Bosnia-Herzegovina, Bulgaria, Croatia, Great Britain, Greece, Hungary, Israel, Poland, Romania, Slovenia, Switzerland, Turkey (Geller-Grimm et al., 2015).

#### 8. *Philodicus ponticus* (Bigot, 1880)

**Material examined:** East Azerbaijan province, Arasbaran, (46° 26.536'N, 44° 54.041' E) 1534m, 21.6.2015, 2♂♂; West Azerbaijan province, Ziveh, (37° 08'N, 44° 52'E) 2630 m, 24.07.2015, 3♂♂, leg. S. Khaghaninia.

**Iranian Records:** Fars (Saghaei et al., 2008; Tomasovic & Saghaei, 2009), Golestan, Guilan (Hayat et al., 2008), Kerman, Khorasan (Becker & Stein, 1913), Khuzestan (Ghahari et al., 2007), Sistan & Baluchestan (Becker & Stein, 1913; Oldroyd, 1958), Iran (no locality cited) (Engel, 1930; Theodor, 1980).

**Distribution:** Afghanistan, Greece, Iraq, Israel, South European territory, Soviet Middle Asia, former Transcaucasus Republics (incl. Azerbaijan), Turkey (Geller-Grimm et al., 2015).


#### 9. *Philonicus albiceps* (Meigen, 1820)

**Material examined:** East Azerbaijan province, Qaradagh forests, (38° 55.601' N, 46° 46.932' E) 1313 m, 21.7.2015, 1♂, 1♀, leg. S. Khaghaninia.


**Iranian Records:** Fars (Tomasovic & Saghaei, 2009), Guilan, Northern Iran (Ricardo, 1920), Kordestan (Hayat et al., 2008), Mazandaran (Ghahari et al., 2007).

**Distribution:** Albania, Austria, Azerbaijan, Belgium, Bosnia-Herzegovina, Bulgaria, Croatia, Cyrus, Czech Republic, Denmark, Finland, France, Germany, Great Britain,

Greece, Hungary, Iceland, Ireland, Israel, Italy, Liechtenstein, Luxembourg, Malta, Mongolia, Norway, Poland, Portugal, Romania, Russia (East Siberia, Far East), Slovakia, Slovenia, Spain, Sweden, Switzerland, The Netherlands, Turkey, United Kingdom (Geller-Grimm et al., 2015).



**Figures 1-3.** *Dismachus transcausicus* Richter, 1962, male: 1. lateral view, 2. wing, 3. Fore femur.



**Figures 4–8.** *Dymachus transcaucasicus* Richter, 1962, male: **4.** epandrium, dorsal view, **5.** internal surface of gonocoxite, **6.** external surface of gonostylus, **7.** aedeagus, lateral view, **8.** hypandrium.

**10. *Polyphonus laevigatus* Loew, 1848**

**Material examined:** East Azerbaijan province, Kandovan, (37° 45.00' N, 46° 18.002' E) 2840, 16.6.2015, 2♀♀, leg. S. Khaghaninia.

**Iranian Records:** Khuzestan (Oldroyd, 1958), West Azarbaijan (Shoeibi & Karimpour, 2010).

**Distribution:** Albania, Azerbaijan, Greece, Palestine, Syria, Turkey (Geller-Grimm et al., 2015).

**11. *Polysarca violacea* Schiner, 1867\***

**Material examined:** East Azerbaijan province, Kandovan, (37° 44.200' N, 46° 18.001' E) 3090m, 25.7.2016, 1♂; Qaradagh forests, (38° 55.601' N, 46° 46.932' E) 1313 m, 21.7.2015, 3♂♂, leg. S. Khaghaninia.

**Iranian Records:** Guilan (Lehr, 1963).

**Distribution:** Algeria, Tunisia, Turkey (Geller-Grimm et al., 2015).



Figures 9–10. *Saropogon megriensis* Richter, 1966, female: 9. dorsal view, 10. head, lateral view.

### 12. *Promachus leoninus* Loew, 1848

**Material examined:** East Azerbaijan province, Chichekli, (38°37.169' N, 46°26.536' E) 1534m, 26.6.2015, 2♂♂, leg. S. Khaghaninia.

**Iranian Records:** Golestan, Kerman (Hayat et al., 2008), Iran (no locality cited) (Lehr, 1988).

**Distribution:** Azerbaijan, Bosnia-Herzegovina, Croatia, Greece, India, Israel, Myanmar (Burma), Romania, Russia, Slovenia, Turkey (Geller-Grimm et al., 2015).

### 13. *Saropogon megriensis* Richter, 1966\*

**Material examined:** East Azerbaijan province, Arasbaran, (38° 50.487' N, 46°54.189' E) 1656m, 26.6.2016, 1♀, leg. S. Khaghaninia.

**Distribution:** Armenia (Geller-Grimm et al., 2015), Iran (**New record**).

**Diagnostic characters:** Head and abdomen not drooping; Mystax covering only lower margin of face; ac present; above the epistoma no bristles or pile. anterior tibial process extremely stout and long, extends beyond the apex of tibia; hind femur

without stout bristles. fourth posterior cell closed and stalked; (Bei-Bienko, 1988; Major, 2000) (Figs 9–10).

### 14. *Saropogon platynotus* (Loew, 1847)

**Material examined:** East Azerbaijan province, Arasbaran (38° 51.077 N, 46° 59.932' E) 1367m, 26.6.2016, 1♂, leg. S. Khaghaninia.

**Iranian Records:** Golestan, Mazandaran (Ghahari et al., 2007).

**Distribution:** Turkey (Geller-Grimm et al., 2015).

### Discussion

During recent years a large amount of specimens of the family Asilidae have been collected from East and West Azerbaijan provinces. *Philodicus ponticus* (Bigot, 1880) is one of the most common robber fly species in our country, followed by *Machimus annulipes* (Brullé, 1832) and *Philonicus albiceps* (Meigen, 1820) (Mohammadi & Khaghaninia, 2016). The first species was the most common asilid in the study area, and the latter can be found

mostly at higher elevations. *Didymachus picipes* (Meigen, 1820) appeared to be comparatively common species. More interesting is the finding of *Saropogon platynotus* (Loew, 1847) of the subfamily Dasypogoninae, which is nearly rare in Iran (Ghahari et al., 2007). East and West Azerbaijan provinces are large region with various biogeographical areas, so it is expected that further species can be found in this area. As the fauna of the family Asilidae remain unknown, so study of this family in East and West Azerbaijan provinces can increase the number of species of this family in Iran.

### Acknowledgments

We would like to thank Dr. Geller Grimm (Museum Wiesbaden Natural History; Germany) for his help in obtaining invaluable data on Iranian Asilidae.

### Conflict of Interests

The author declare that there is no conflict of interest regarding the publication of this paper.

### References

- Abbassian-Lintzen, R. (1964a) Asilidae (Diptera) of Iran. I. Robber flies belonging to the subfamilies Laphriinae and Dasypogoninae (with description of new species). *Annals and Magazine of Natural History*, 13, 417–435.  
<https://doi.org/10.1080/00222936408651481>
- Abbassian-Lintzen, R. (1964b) Asilidae (Diptera) of Iran. II. Notes on the genus *Eremisca* Zin. and description of *E. schahgudiani* n. sp. *Annals and Magazine of Natural History*, 13, 547–552.  
<https://doi.org/10.1080/00222936408651497>
- Becker, Th. & Stein, P. (1913) Persische Dipteren von den Expeditionen des Herrn N. Zarudny 1898 und 1901. *Ezhegodnik Zoologicheskago Muzeya Imperatorskoi Akademi Nauk*, 17, 503–654.
- Bigot, J. (1880) Dipteres nouveaux ou peu connus. 13. partie (I). XX. Quelques Dipteres de Perse et du Caucase. *Annales de la Societe entomologique de France*, 10, 139–154.
- Dikow, T. (2016) Robber Flies (Asilidae), new catalogue for the genera, Available from: <https://asiloidflies.si.edu/asilidae-generic-classification-dikow2009> [Accessed 20 November, 2016].
- Dikow, T. (2009) Phylogeny of Asilidae Inferred from Morphological Characters of Imagines (Insecta: Diptera: Brachycera: Asiloidea). *Bulletin of the American Museum of Natural History*, 319, 1–175.  
<https://doi.org/10.1206/603.1>
- Ghahari, H., Lehr, P.A., Lavigne, R.J., Hayat, R. & Ostovan, H. (2007) New records of robber flies (Diptera, Asilidae) for the Iranian fauna with their prey records. *Far Eastern Entomologist*, 179, 1–9.
- Geller-Grimm, F., Dikow, T. & Lavigne, R.J. (2015) Robber Flies (Asilidae) Database, Species, Available from: <http://www.gellergrimm.de/catalog/species.htm> Accessed January 5, 2015].
- Geller-Grimm, F. (2003) A world catalogue of the genera of the family Asilidae (Diptera). *Studia dipterologica*, 10, 473–526.
- Geller-Grimm, F. (2002) Robber flies (Diptera: Asilidae) of the Socotra Archipelago, Yemen. *Fauna of Arabia*, 19, 467–489.
- Hayat, R., Ghahari, H., Lavigne, R., & Ostovan, H. (2008) Iranian Asilidae (Insecta: Diptera). *Turkish Journal of Zoology*, 32, 175–195.
- Hermann, F. (1905) Beitrag zur Kenntnis der Asiliden. *Berliner Entomologische Zeitschrift*, 50, 14–42.  
<https://doi.org/10.1002/mmnd.19050500106>
- Hradský, M. & Geller-Grimm, F. (1998) Eine neue Art der Gattung *Eutolmus* Loew, 1967 aus Vietnam (Diptera, Asilidae) [A new *Eutolmus* species from Vietnam (Diptera, Asilidae)]. *Studia dipterologica*, 5 (2), 249–252.
- Janssens, E. (1961) Sur quelques Dipteres (Asilidae) de l'Iraq. *Bulletin Institut Royal des Sciences naturelles de Belgique*, 37, 1–8.
- Lehr, P.A. (1988) Family Asilidae. In: Soos, A. & Papp, L. (Eds.), *Catalogue of Palaearctic Diptera*. Akadémiai Kiadó, Budapest, pp. 197–326.
- Lehr, P. A., Ghahari, H., & Ostovan H. (2007) A contribution to the Robber Flies of

- Subfamilies Stenopogoninae and Asilidae (Diptera: Asilidae) from Iran. *Far Eastern Entomologist*, 173, 1–14.
- Lehr, P.A. (1963) A Review of the Robber-Fly Genera *Polysarca* Schin., *Polysarcodes* Par., *Satanas* Jac., *Trypanoides* Beck., *Promachus* Lw., *Philodicus* Lw., and *Trichardopsis* Oldr. (Diptera, Asilidae) of the Fauna of the USSR, *Entomologicheskoe Obozrenie*, 42 (1), 195–210
- Mohammadi, R. & Khaghaninia, S. (2016a) Family Asilidae (Diptera: Brachycera: Asiloidea) in East Azerbaijan province, with two new records for Iranian Fauna. *Journal of Insect Biodiversity and Systematics*, 1 (2), 125–132.
- Mohammadi, R. & Khaghaninia, S. (2016b) Flies of the subfamily Asilinae Latreille, 1802 (Diptera: Asilidae) in East Azerbaijan province, with three new records for Iranian Fauna. *Biharean Biologist*, 10 (2), 79–81.
- Oldroyd, H. (1958) Some Asilidae from Iran. *Stuttgarter Beitrage zur Naturkunde*, 9: 1–10.
- Portschinsky, J. (1873) "Deux dipteres nouveaux de la Perse septentrionale." *Horae Societatis Entomologicae Rossicae*, 9, 292–293.
- Richter, V.A., Asilidae. In: Bei-Bienko, G. (1988) Keys to the Insect of the European Part of the USSR. Volume V. Diptera and Siphonoptera. Part II. Smithsonian Institution Libraries and The National Science Foundation Washington, D.C. pp. 779–819.
- Saghaei, N., Ostovan, H., Shojaei, M. & Hayat, R. (2008) Interduction to the Asilidae Fauna (Insecta: Diptera) of Fars Province, Iran. *Turkish Journal of Zoology*, 33, 187–200.
- Shoeibi, B. & Karimpour, Y. (2010) Contributions to the knowledge of Asilidae (Diptera: Brachycera) from Azarbaijan Province (Iran). *Munis Entomology and Zoology*, 5, 957–963.
- Theodor, O. (1980) *Diptera: Asilidae. Fauna Palaestina, Insecta*. Academy of Sciences and Humanities, Jerusalem, Israel, 453 pp.
- Timon-David, J. (1955) *Iranopogon brandti*, n. gen., n. sp., Asilidae xerophile d'Iran (Diptera). *Bulletin de la Societe entomologique de France*, 60, 102–104.
- Tomasovic, G. (2002) Etude sur materiaux typiques du complexe genital male de spet espece du genre *Erax* Scopoli, 1763 (Diptera: Asilidae) avec la description de trois especes nouvelles. *Notes faunistiques de Gembloux*, 46, 27–37.
- Tomasovic, G. (1999) Notes sur les Asilidae palearctiques (Diptera Brachycera) (10 et 11). Description et repartition geographique de 2 especes nouvelles de *Dasypogon* du groupe diadema (Fabricius, 1781). *Bulletin de la Societe royale Beige d'Entomologie*, 135, 216–221.
- Tomasovic, G. & Saghaei, N. (2009) Contribution to the knowledge of the Asilidae (Diptera: Brachycera) from Fars province (Iran). *Faunistic Entomology*, 62 (2), 45–56.
- Tsacas, L. (1968) Revision des especes du genere *Neomochtherus* Osten-Sacken (Dipteres: Asilidae). I. Region palearctique. [Revision of the species of the genus *Neomochtherus* Osten-Sacken (Diptera: Asilidae). I. Palearctic region]. *Bulletin of the British Museum, Natural History, Zoology*, 47, 129–328.
- Weinberg, M & Bächli, G. (1995) *Diptera: Asilidae*. Vol. 11, *Insecta Helvetica* (Fauna). Schweizerischen Entomologischen Gesellschaft, Geneva, 124 pp.



## مطالعه دزد مگس‌ها (Diptera: Asilidae) در استان‌های آذربایجان شرقی و غربی با گزارش دو گونه جدید برای کشور

رحمان محمدی<sup>۱\*</sup>، صمد خاقانی‌نیا<sup>۱</sup> و دیمیتری آستاخوف<sup>۲</sup>

۱ گروه گیاهپزشکی، دانشکده کشاورزی، دانشگاه تبریز، ۵۱۶۶۴، تبریز، ایران.

۲ دانشگاه ایالت وولگوگراد، ۴۰۰۶۲، وولگوگراد، روسیه.

\* پست الکترونیکی نویسنده مسئول مکاتبه: [Rahmanmohamadi90@gmail.com](mailto:Rahmanmohamadi90@gmail.com)

تاریخ دریافت: ۲۹ خرداد ۱۳۹۶، تاریخ پذیرش: ۱ شهریور ۱۳۹۶، تاریخ انتشار: ۷ شهریور ۱۳۹۶

**چکیده:** بر اساس نمونه‌های جمع‌آوری شده از استان‌های آذربایجان شرقی و غربی طی سال‌های ۱۳۹۳-۱۳۹۵، ۱۴ گونه از خانواده Asilidae جمع‌آوری و شناسایی شدند. دو گونه از آن‌ها، با نام‌های *Dysmachus transcausicus* Richter, 1962 و *Saropogon megriensis* Richter, 1966 برای اولین بار از ایران گزارش می‌شوند. لیست همه گونه‌ها همراه با پراکنش جغرافیایی آن‌ها آورده شده است. ویژگی‌های افتراقی و همچنین تصاویر تکمیلی گزارش‌های جدید ارائه شده است.

**واژگان کلیدی:** Asilidae، آذربایجان شرقی، آذربایجان غربی، ایران، گزارش جدید