A survey on the click beetles (Coleoptera, Elateridae) in the province of Kurdistan (Iran), with three new records for the Iranian fauna

Margarit Marjanyan¹, Hamed Ghobari^{2*} & Jamasb Nozari³

¹ Institute of Zoology, Scientific Centre of Zoology and Hydroecology of the National Academy of Sciences of Armenia, Yerevan, 0014, Armenia. mmarjanyan@yahoo.com

² Department of Plant Protection, College of Agriculture, University of Kurdistan, Sanandaj, Iran. h.ghobari@uok.ac.ir

³ Department of Plant Protection, University College of Agriculture & Natural Resources, University of Tehran, Karaj, Iran, PO Box 4111, Karaj, Iran. nozari@ut.ac.ir

Received:
25 July, 2020

Accepted:
25 November, 2020

Published:
07 December, 2020

Subject Editor:
Hiva Nasserzadeh

ABSTRACT. This study was carried out in the province of Kurdistan (Iran) during April to August 2015 in order to document the click beetle fauna (Coleoptera: Elateridae) of this region. The specimens were collected using sticky traps, window traps and pan traps. Fifteen species from six genera were identified, of which three species are new for the fauna of Iran: *Melanotus kravchenkoi* Platia, 2010, *Melanotus orbachorum* Platia, 2010 and *Melanotus chikatunovi* Platia, 2010. In addition, nine species are recorded from the province of Kurdistan for the first time. Brief diagnoses for the new record species from Iran and zoogeographical information for all species are provided.

Key words: Click beetles, fauna, Iran, Kurdistan province, zoogeography

Citation: Marjanyan, M., Ghobari, H. & Nozari, J. (2021) A survey on the click beetles (Coleoptera, Elateridae) in the province of Kurdistan (Iran), with three new records for the Iranian fauna. *Journal of Insect Biodiversity and Systematics*, 7 (2), 127–135.

Introduction

The Elateridae is one of the largest families of the Coleoptera with more than 10,000 described species worldwide (Platia & Ghahari, 2016). Various studies were carried out on Iranian fauna of the Elateridae in the past (e.g. Cate et al., 2002; Platia et al., 2002; Bagheri et al., 2004a, 2004b; Platia & Gudenzi, 2005, 2006; Mertlik & Dusanek, 2006; Nasserzadeh et al., 2008; Platia & Németh, 2011; Gholikhajeh et al., 2012; Farhangian et al., 2013; Mardjanian et al., 2013; Mertlik & Németh, 2014; Németh & Platia, 2014; Platia & Ghahari, 2016; Mardjanian & Barimani, 2017; Nasserzadeh et al., 2018; Kundrata et al., 2019). According to the latest research of the click-beetles of Iran, 246 species were recorded from Iran (Platia & Ghahari, 2016; Kundrata et al., 2019). In this paper, three species are added to known fauna of Iranian Elateridae.

Material and methods

The material was collected from different parts of Kurdistan Province (Iran) with different types of vegetation (rangelands, oak forests and gardens) by using different traps such as sticky traps,

Corresponding author: Hamed Ghobari, E-mail: h.ghobari@uok.ac.ir

Copyright © 2021, Marjanyan et al. This is an open access article distributed under the terms of the Creative Commons Attribution License (CC BY 4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

<https://doi.org/10.24238112.2021.7.2.3.5>

window traps and pan traps (Fig. 1). Data on the total distribution of beetles are given according to the Catalogue of Palaearctic Coleoptera (Cate, 2007), their original descriptions and reviews (Gurjeva, 1979; Marjanyan, 1987; Gurjeva, 1988; Platia, 2010). All specimens are deposited in the Collection of Invertebrates of the National Academy of Sciences of Armenia (CINASA).

Figure 1. Map of the sampling area of click beetles in Kurdistan. Red stars indicate the sampling locations.

Results

Fifteen species belong to six genera are listed as follows. Details on the collecting localities and the general distribution and distribution in Iran are given for all species.

Subfamily Agrypninae Candèze, 1857

Lacon punctatus (Herbst, 1779)

Material examined: IRAN, Kurdistan prov., Marivan-Sarvabad, (35°17'41" N, 46°22'25" E, 1160 m), 19.viii.2015, oak forest, 1♀, 2♂♂, leg. H. Ghobari (CINASA).

General distribution: Armenia, Azerbaijan, Bosnia Herzegovina, Bulgaria, Croatia, Russia: Central and South European Territory, Czech Republic, France, Germany, Georgia, Greece, Hungary, Italy, Malta, Portugal, Slovakia, Slovenia, Spain, Switzerland, Turkey, Ukraine, Serbia, Montenegro, Cyprus, Jordan, Syria, Morocco (incl. Western Sahara) (Cate, 2007), new record for Iran.

Drasterius bimaculatus (P. Rossi, 1790)

Material examined: IRAN, Kurdistan prov., Marivan-Garan, (35°32'02" N, 46°18'05" E, 1350 m), 18.vi.2015, oak forest, 3♀♀, 3♂♂, leg. H. Ghobari. (CINASA). IRAN, Kurdistan prov., Marivan-Sarvabad, (35°17'41" N / 46°22'25" E, 1160 m), oak forest, 11.vii.2015, 2♀♀, 4♂♂, leg.

H. Ghobari, (CINASA). IRAN, Kurdistan prov., Sanandaj-Noshor, (35°01'39" N, 46°56'05" E, 1460 m), garden, 04.vii, 01.viii, and 16.viii.2015, 4♀♀, 1♂, leg. H. Ghobari, (CINASA).

General distribution: Albania, Armenia, Austria, Azerbaijan, Belgium, Bosnia Herzegovina, Bulgaria, Croatia, Russia: Central and South European Territory, Czech Republic, France (incl. Corsica, Monaco), Germany, Georgia, Greece (incl. Crete), Hungary, Italy (incl. San Marino, Sardinia, Sicily), Malta, Macedonia, Moldavia, Poland, Portugal, Romania, Slovakia, Slovenia, Spain (incl. Gibraltar), Switzerland, Ukraine, Yugoslavia (Serbia, Montenegro), North Africa: Algeria, Canary Islands, (Lanzarote) Egypt, Libya, Morocco (incl. Western Sahara), Tunisia, Asia: Afghanistan, Cyprus, Iran, Iraq, Israel, Jordan, Kazakhstan, Kyrgyzstan, Syria, Turkey, Turkmenistan, Uzbekistan (Cate, 2007).

Distribution in Iran: Alborz, Ardabil, Golestan, Sistan & Baluchestan, West Azarbaijan, Chaharmahal & Bakhtiari, Qom, East Azarbaijan, Guilan, Mazandaran, Isfahan, Tehran, Fars, Golestan, Hamadan, Kermanshah, Northern Khorasan (Platia & Ghahari, 2016), new record for Kurdistan.

Subfamily Elaterinae Leach, 1815

Betarmon bisbimaculatus (Fabricius, 1803)

Material examined: IRAN, Kurdistan prov., Sanandaj-Noshor, (35°01'39" N, 46°56'05" E, 1460 m), garden, 04.vii.2015, 2♀♀, 2♂♂, leg. H. Ghobari, (CINASA).

General distribution: Armenia, Austria, Azerbaijan, Bulgaria, Croatia, Czech Republic, France (incl. Corsica, Monaco), Germany, Georgia, Hungary, Italy (incl. San Marino, Sardinia, Sicily), Poland, Romania, Slovakia, Slovenia, Spain (incl. Gibraltar), Switzerland, Ukraine, Yugoslavia (Serbia, Montenegro), Asia: Iran, Turkey (Cate, 2007).

Distribution in Iran: Guilan, Sistan & Baluchestan, Mazandaran, Golestan (Platia & Ghahari, 2016), new record for Kurdistan.

Synaptus filiformis (Fabricius, 1781)

Material examined: IRAN, Kurdistan prov., Sanandaj-Noshor, (35°01'39" N, 46°56'05" E, 1460 m), garden, 11.vii, 01.viii, and 16.viii.2015, 1♀, 1♂, leg. H. Ghobari, (CINASA).

General distribution: Albania, Armenia, Austria, Azerbaijan, Belgium, Bosnia Herzegovina, Bulgaria, Belarus, Croatia, Russia: Central and South European Territory, Czech Republic, Estonia, France (incl. Corsica, Monaco), Germany, Georgia, Great Britain (incl. Channel Islands), Greece (incl. Crete), Hungary, Italy (incl. San Marino Sardinia, Sicily), Latvia, Liechtenstein, Lithuania, Luxembourg, Macedonia, Moldavia, The Netherlands, Poland, Portugal, Romania, Slovakia Slovenia, Spain (incl. Gibraltar), Switzerland, Ukraine, Yugoslavia (Serbia, Montenegro), Asia: Iran, Kazakhstan, Syria, Turkey (Cate, 2007), Lebanon, Tadzhikistan (Platia, 2010).

Distribution in Iran: Guilan, Isfahan, Mazandaran (Platia & Ghahari, 2016), new record for Kurdistan.

Adrastus dolini Wellschmied, 1978

Material examined: IRAN, Kurdistan prov., Sanandaj-Noshor, (35°01'39" N, 46°56'05" E, 1460 m), garden, 14.viii.2015, 1♀, 2♂♂, leg. H. Ghobari, (CINASA).

General distribution: Azerbaijan, Georgia, Russia: South European Territory, Asia: Iran, Turkey (Cate, 2007).

Distribution in Iran: Mazandaran (Platia & Ghahari, 2016), new record for Kurdistan.

Melanotus castanipes (Paykull, 1800)

Material examined: IRAN, Kurdistan prov., Sanandaj-Saral station, (35°40'18" N, 47°07'07" E, 2070 m), rangeland, 20.vi.2015, 2♀♀, 2♂♂, leg. H. Ghobari, (CINASA).

General distribution: Austria, Belgium, Bosnia Herzegovina, Bulgaria, Belarus, Croatia, Russia: North and Central, South European Territory, Czech Republic, Denmark, Estonia, Finland, France (incl. Corsica, Monaco), Great Britain (incl. Channel Islands), Germany, Georgia, Greece (incl. Crete), Hungary, Italy (incl. San Marino, Sardinia, Sicily), Latvia, Liechtenstein, Lithuania, Luxembourg, Norway, Poland, Romania, Slovakia, Slovenia, Spain (incl. Gibraltar), Sweden, Switzerland, Ukraine, Yugoslavia (Serbia, Montenegro) Asia: China [Sichuan (Szechwan), Shanxi (Shansi), Yunnan], India: Uttaranchal, Iran, Jilin (Kirin), Pakistan, Turkey, , Uttar, Russia: west Siberia, (Cate, 2007).

Distribution in Iran: Guilan, Mazandaran (Platia & Ghahari, 2016), new record for Kurdistan.

Melanotus monticola Menetries, 1832

Material examined: IRAN, Kurdistan prov., Sanandaj-Saral station, (35°40'18" N, 47°07'07" E, 2070 m), rangeland, 19.viii.2015, 3♀♀, 1♂, leg. H. Ghobari, (CINASA).

General distribution: Armenia, Azerbaijan, Bulgaria, Georgia, Russia: South European Territory, Asia: Iran, Turkey (Cate, 2007).

Distribution in Iran: East Azarbaijan, Mazandaran (Platia & Ghahari, 2016), new record for Kurdistan.

Melanotus fulvus Reitter, 1891

Material examined: IRAN, Kurdistan prov., Sanandaj-Saral station, (35°40'18" N, 47°07'07" E, 2070 m), rangeland, 11.vii.2015, 1♀, 1♂, leg. H. Ghobari, (CINASA).

General distribution: Iran, Iraq, Turkmenistan (Cate, 2007).

Distribution in Iran: Alborz, Northern Khorasan, Razavi Khorasan, Fars, Kuhgiluyeh & Boyerahmad, Razavi Khorasan (Platia & Ghahari, 2016), new record for Kurdistan.

Melanotus chikatunovi Platia, 2010

Material examined: IRAN, Kurdistan prov., Sanandaj-Saral station, (35°40'18" N, 47°07'07" E, 2070 m), rangeland, 19.viii.2015, 2♂♂, leg. H. Ghobari, (CINASA); Kurdistan prov., Sanandaj-Noshor, (35°01'39" N, 46°56'05" E, 1460 m), garden, 19.viii.2015, 1♂, leg. H. Ghobari, (CINASA).

General distribution: Israel (Platia, 2010), New record for Iran.

Diagnostic characters: The body is reddish brown entirely and covered with dense and long yellow vestiture that is erect at sides of elytra partially. The width of the pronotum is greater than its length. Scutellum is plate-shaped and dotted, elytra have the same width of pronotum. The species is similar to *M. fulvus* Reitter in color and overall shape of body

However; it can be distinguished by its larger body and longer antennae than *M. fulvus* Reitter the species (Platia, 2010). Body length: 12.5-13 mm.

***Melanotus kravchenkoi* Platia, 2010**

Material examined: IRAN, Kurdistan prov., Sanandaj-Saral station, (35°40'18" N, 47°07'07" E, 2070 m), rangeland, 19.viii.2015, 2 ♂♂, leg. H. Ghobari, (CINASA).

General distribution: Palestine, Israel (Platia, 2010), new record for Iran.

Diagnostic characters: very similar to *M. chikatunovi* the body is bright fairly and entirely dark brown, Like *M. chikatunovi* Platia, the body is covered with dense and long yellow vestiture (pubescens) that is erect at sides of elytra partially but can be distinguished by darker color and more elevated carina of posterior angles of pronotum (Platia, 2010). Body length: 11-12 mm.

***Melanotus dichroides* Platia & Gudenzi, 1999**

Material examined: IRAN, Kurdistan prov., Sanandaj-Noshor, (35°01'39" N, 46°56'05" E, 1460 m), garden, 19.viii.2015, 1♀, 1♂, leg. H. Ghobari, (CINASA).

General distribution: Iran (Cate, 2007; current study).

Distribution in Iran: Guilan (Platia & Ghahari, 2016), new record for Kurdistan.

***Melanotus orbachorum* Platia, 2010**

Material examined: IRAN, Kurdistan prov., Sanandaj-Saral station, (35°40'18" N, 47°07'07" E, 2070 m), rangeland, 11.vii.2015, 1♂, leg. H. Ghobari, (CINASA); Kurdistan prov., Sanandaj-Noshor, (35°01'39" N, 46°56'05" E, 1460 m), garden, 19.viii.2015, 1♀, 1♂, leg. H. Ghobari, (CINASA).

General distribution: Israel (Platia, 2010), New record for Iran.

Diagnostic characters: The body is black completely antennae and legs is dark brown and covered with fine pubescence and erected on head and pronotum. The width of the pronotum is greater than its length, Pronotum 1.2x wider than long. Scutellum is plate-shaped and dotted, elytra are longer than pronotum but they are the same width. Scutellum is shaped like a tongue. Elytra are punctured and have distinct striae. body covered with dense, yellowish vestiture (Platia, 2010). Body length: 14-17.5 mm.

***Melanotus fusciceps* (Gyllenhal, 1817)**

Material examined: IRAN, Kurdistan prov., Sanandaj-Saral station, (35°40'18" N, 47°07'07" E, 2070 m), rangeland, 27.vii.2015, 1♂♂, leg. H. Ghobari, (CINASA).

General distribution: Azerbaijan, Armenia, Azores, Bulgaria, Croatia, Russia: Central European Territory, Georgia, Macedonia, Moldavia, Romania, Russia: South European Territory, Ukraine, Cyprus, Kazakhstan, Iran, Iraq, Israel, Lebanon, Syria, Turkey (Cate, 2007).

Distribution in Iran: Ardabil, Chaharmahal & Bakhtiari, Isfahan, Markazi, Bushehr, Fars, Qazvin, West Azarbaijan, Zanzan, Golestan, Fars (Platia & Ghahari, 2016), new record for Kurdistan.

Subfamily Cardiophorinae (Candeze, 1860)

Cardiophorus nigratissimus Buysson, 1891

Material examined: IRAN, Kurdistan prov., Sanandaj-Saral station, (35°40'18" N, 47°07'07" E, 2070 m), rangeland, 20.06., 11.07. and 27.vii.2015, 2♀♀, 2♂♂, leg. H. Ghobari, (CINASA); Kurdistan prov., Sanandaj-Noshor, 35°01'39" N, 46°56'05" E, 1460 m, garden, 19.viii.2015, 2♀♀, 2♂♂, leg. H. Ghobari, (CINASA).

General distribution: Azerbaijan, Armenia, Georgia Iran, Iraq, Syria, Turkmenistan, Turkey (Cate, 2007).

Distribution in Iran: Bushehr, East Azarbaijan, Khuzestan, Zanjan, Chaharmahal & Bakhtiari, Fars, Qazvin, Tehran, Golestan, Fars, Guilan, Hamadan, Kurdistan, Lorestan, West Azarbaijan, Zanjan (Platia & Ghahari, 2016).

Cardiophorus magnanii Platia, Furlan & Gudenzi, 2002

Material examined: IRAN, Kurdistan prov., Sanandaj-Noshor, 35°01'39" N, 46°56'05" E, 1460 m, garden, 14.viii.2015, 3♀♀, 3♂♂, leg. H. Ghobari, (CINASA).

Distribution: Iran (Cate, 2007; current study).

Discussion

This study provides additional zoogeographical data for the click beetle fauna of Kurdistan from the field. With three new records that have been documented in this study, the number of known click beetle species in Iran has increased to 249.

The heterogeneity of geography and microclimates in Iran has furnished a unique diversity of flora and vegetation cover across the country. This in turn seems to enhance the trends of diversity for many insects' families such as Elateridae. The results of this study demonstrate the urgent need for a more extensive and a long-term collection of click beetles across Iran. It also highlights the importance of involving regional taxonomic expertise in these studies.

Acknowledgments

This study was funded by University of Kurdistan, Iran under grant number 4/7069.

Conflict of Interests

The authors declare that there is no conflict of interest regarding the publication of this paper.

ORCID

Hamed Ghobari: <https://orcid.org/0000-0002-7832-4879>

Jamasb Nozari: <https://orcid.org/0000-0002-2649-7751>

References

- Bagheri, M.R., Rajabi, Gh.R., Ardebili, Zh., Haghshenas, A.R., Hosseini Nia, A. & Poorghaz, A. (2004a) Faunistic study of click beetles (Col.: Elateridae) in Iran. In: *Proceedings of the 16th Iranian Plant Protection Congress, 2004, 28 August – 01 September, University of Tabriz, Tabriz, Iran*. Ministry of Agriculture, Tehran, p. 129.

- Bagheri, M., Poorghaz, A. & Furlan, L. (2004b) Efficiency of sexual pheromone of seven species of *Agriotes* (Col.: Elateridae) on attraction of many adults in different regions of Iran. *Proceedings of 3rd National Conference on the Development in the application of Biological products & optimum utilization of chemical fertilizers & pesticides in Agriculture*, 2004, 21–23 February, Karaj, Iran. Agricultural Research, Education and Extension Organization, p. 515.
- Cate, P. C. (2007) Family Elateridae. In: Loeb I. & Smetana A. (eds.) *Catalogue of Palearctic Coleoptera. Vol. 4., Elateroidea - Derodontoidea - Bostrichoidea - Lymexyloidea - Cleroidea - Cucujidae*. Apollo Books, Stenstrup, 935 pp.
- Cate, P.C., Platia, G. & Schimmel, R. (2002) New species and records of click beetles (Coleoptera: Elateridae) from Iran, with a checklist of known species. *Folia Heyrovskyana*, 10 (1), 25–68.
- Dolin, W.G. & Atamuradov, Kh.U. (1993) *Click – beetles (Elateridae) of Turkmenistan*. Kiev. Institute of Zoology Acad. Sci. of Ukraina. 175 pp.
- Farhangian, S., Sakenin Chelav, H. & Monem, R. (2013) A faunistic study on Elateridae (Coleoptera: Elateroidea) from Isfahan, Iran. *Calodema*, 264, 1–3.
- Gholikhajeh, N., Nozari, J. & Hosseininaveh, V. (2012) Identification of damaging species of wireworms of genus *Agriotes* (Col.: Elateridae) in the wheat fields of the east of Golestan province. In: Sarafrazi, A. (ed.) *Proceedings of the 20th Iranian Plant Protection Congress, 2012, 25–28 August, University of Shiraz, Shiraz, Iran*. Ministry of Agriculture, Tehran, p. 149.
- Gurjeva, E.L. (1979) The click beetles (Elateridae). Subfamily Elaterinae. Tribes Megapenthini, Physorhinini, Ampedini, Elaterini, Pomachilini. In: Strelkov, A.A. & Medvedev, G.S. (eds.) *Fauna of the USSR. Beetles*. Vol. 12, Issue 4, Nauka, Leningrad, pp. 1–452. (In Russian).
- Gurjeva, E.L. (1988) New species of click-beetles of genus *Melanotus* Esch. (Coleoptera, Elateridae) from Middle Asia. *Entomological Review*, 72 (2), 338–345. (In Russian).
- Kundrata, R., Nemeth, T. & Jarzabek-Müller, A. (2019) Description of *Lacon mertliki* sp. nov. (Coleoptera: Elateridae: Agrypninae) from the Hyrcanian forest ecoregion, with a key to the *Lacon* species of Iran. *European Journal of Taxonomy*, 535, 1–13.
<https://doi.org/10.5852/ejt.2019.535>
- Marjanyan, M.A. (1987) *The Fauna of the Armenian SSR. Click Beetles (Elateridae)*. Acad. Sci. of the Armenian SSR, Yerevan, pp. 1–207. (In Russian)
- Marjanyan, M.A. & Barimani Varandi, H. (2014) New notes on fauna of click-beetles (Coleoptera, Elateridae) of Mazandaran province (Iran). *Biological Journal of Armenia*, 66(3), 81–83. (In Russian).
- Marjanyan, M.A. & Barimani Varandi, H. (2017) New species of click – beetles of genus *Cardiophorus* Eschscholtz, 1829 (Coleoptera, Elateridae, Cardiophorinae) from Mazandaran (Iran). *Reports*, 111 (4), 396–403. (In Russian).
- Marjanyan, M.A., Barimani Varandi, H. & Kalashian M.Y. (2008) Contribution on the knowledge of the fauna click-beetles (Coleoptera, Elateridae) of Iran. *Biological Journal of Armenia*, 60(3), 68–70. (In Russian).
- Marjanyan, M.A., Khaje, H., Nozari J. & Ghobari H. (2013) Contribution to the fauna of click-beetles (Coleoptera, Elateridae) of province Golestan (Iran). *Biological Journal of Armenia*, 65 (4), 130–132. (In Russian).
- Mertlik, J. & Dusanek, V. (2006) Description of five new species of click-beetles (Coleoptera, Elateridae) from the Palearctic region with remarks about the distributions 22 additional species. *Folia Heyrovskyana, Serie A*, 13(4), 145–162.
- Mertlik, J. & Németh, T. (2014) Distributional notes on *Lacon nadaii* and *L. unicolor* (Coleoptera: Elateridae). *Elateridarium*, 8, 61–66.

- Nasserzadeh, H., Platia, G. & Barimani, H. (2008) Taxonomic and faunistic notes on *Pleurothous* Reitter, and description of the male characters of *Calais brandti* Platia & Gudenzi (Coleoptera: Elateridae). *Koleopterologische Rundschau*, 78, 291–296.
- Nasserzadeh, H., Platia G., Parsi, F. & Serri, S. (2018) Three new species, *Elathous zagrosensis* sp. n., *Zorochros dilatatus* sp. n., and *Dicronychus iranicus* sp. n., and noteworthy records of click beetles from western Iran (Coleoptera: Elateridae). *Zootaxa*, 4394 (4), 517–536.
<https://doi.org/10.11646/zootaxa.4394.4.3>
- Németh, T. & Platia, G. (2014) On some Palaearctic click beetles deposited in the Hungarian Natural History Museum, 2 (Coleoptera: Elateridae), *Zootaxa*, 3841 (4), 451–490.
<https://doi.org/10.11646/zootaxa.3841.4>
- Platia, G. (2010) New species and chorological notes of click beetles from the Palaearctic Region, especially from the Middle East (Coleoptera, Elateridae). *Boletín de la Sociedad Entomológica Aragonesa*, 46, 23–49.
- Platia, G. (2013) Click-Beetles collected in Iran by the expeditions of the Naturkundemuseum Erfurt with description of two new species (Insecta: Coleoptera: Elateridae). *Vernate*, 32, 391–402,
- Platia, G., Furlan, L. & Gudenzi I. (2002) Descrizione di sei nuove specie di Elateridi dell'Iran (Insecta, Coleoptera, Elateridae). *Naturalista Valtellinese - Quaderno di Studi e Notizie di Storia naturale della Romagna*, 13, 65–77.
- Platia, G. & Ghahari, H. (2016) An annotated checklist of click-beetles (Coleoptera, Elateridae) from Iran. *Zootaxa*, 4137 (2), 239–275. <https://doi.org/10.11646/zootaxa.4137.2.6>
- Platia, G. & Gudenzi, I. (2005) Description of eleven new species of click-beetles of the palearctic region, a case of legs teratology and new records of some species of the Italian fauna. *Quaderno di Studi e Notizie di Storia Naturale della Romagna*, 21, 109–127.
- Platia, G. & Gudenzi, I. (2006) Click-beetles genera, species and records new to Palearctic and Indomalayan regions (Insecta Coleoptera Elateridae). *Quaderno di Studi e Notizie di Storia naturale della Romagna*, 23, 131–156.
- Platia, G. & Németh, T. (2011) On some Palaearctic click beetles deposited in the Hungarian Natural History Museum (Coleoptera: Elateridae). *Annales Historico-Naturales Musei Nationalis Hungarici*, 103, 65–106.
- Platia G. & Schimmel, R. (1992) Descrizione di sei nuove specie di Elateridi di Grecia e Turchia (Coleoptera, Elateridae). *Lavori-Società Veneziana di Scienze Naturali*, 17, 87–95.

مطالعه سخت بالپوشان خانواده Elateridae Leach, 1815 در استان کردستان و سه گزارش جدید برای فون ایران

مارگارت مرجانیان^۱، حامد غباری^{۲*} و جاماسب نوذری^۳

۱ مؤسسه جانورشناسی، مرکز علمی جانورشناسی و هیدرواکولوژی اکادمی ملی علوم ارمنستان، ایروان، ارمنستان.

۲ گروه گیاهپزشکی، دانشکده کشاورزی، دانشگاه کردستان، سنندج، ایران.

۳ گروه گیاهپزشکی، پردیس کشاورزی و منابع طبیعی دانشگاه تهران، کرج، ایران.

* پست الکترونیکی نویسنده مسئول مکاتبه: h.ghobari@uok.ac.ir

| تاریخ دریافت: ۰۴ مرداد ۱۳۹۹ | تاریخ پذیرش: ۰۵ آذر ۱۳۹۹ | تاریخ انتشار: ۱۷ آذر ۱۳۹۹ |

چکیده: این مطالعه بر اساس بررسی نمونه های جمع آوری شده متعلق به خانواده سخت بالپوشان خانواده Elateridae از عرصه های طبیعی استان کردستان در طول ماه های اردیبهشت تا شهریور ماه سال ۱۳۹۴ انجام گرفته است. در طی این تحقیق از روش های مختلف شامل تله پنجره ای، تله های سطلی و چسبنده رنگی برای جمع آوری نمونه ها استفاده شد. بر اساس نتایج بدست آمده ۱۵ گونه از ۶ جنس شناسایی شدند، که از این تعداد ۳ گونه از آنها که شامل گونه های *Melanotus chikatunovi* Platia, 2010، *M. kravchenkoi* Platia, 2010 و *Melanotus orbachorum* Platia, 2010 برای فون ایران جدید تشخیص داده شدند. علاوه بر این ۹ گونه نیز برای اولین بار از استان کردستان گزارش شد. در این تحقیق خلاصه ای از خصوصیات مهم تشخیصی هر کدام از گونه های جدید برای فون ایران و همچنین پراکنش جغرافیایی هر کدام از گونه های جمع آوری شده، ارائه شده است.

واژگان کلیدی: سوسک های پشتک زن، فون، ایران، استان کردستان، جغرافیای جانوری