



ESI

Received:
05 November, 2017

Accepted:
16 January, 2018

Published:
18 January, 2018

Subject Editor:
Ali Asghar Talebi

Xenylla mediterranea da Gama, 1964 (Collembola: Poduromorpha: Hypogastruride): A new record for Iranian fauna (Mazandaran province)

Leila Mohammadi Nodeheki^{1*}, Masoumeh Shayanmehr¹ and Mohsen Yazdanian²

¹ Department of Plant Protection, Sari Agricultural Science and Natural Resources University, Po. Box. 578, Sari, Mazandaran, Iran.

² Department of Plant Protection, Gorgan Agricultural Sciences and Natural Resources University, Iran.

ABSTRACT. During the collecting of Hypogastruridae (Collembola) in Mazandaran province, *Xenylla mediterranea* da Gama, 1964 was reported for the first time from Iran. With the new record in this study, the number of *Xynella* species known from Iran increased to four. Information for specie including material examined, description, ecology, distribution, and illustrations are given.

Key words: *Xenylla mediterranea*, Hypogastruridae, Collembola, Iran, New record

Citation: Mohammadi Nodeheki, L., Shayanmehr, M. & Yazdanian, M. (2018) *Xenylla mediterranea* da Gama, 1964 (Collembola: Poduromorpha: Hypogastruride): A new record for Iranian fauna (Mazandaran province). *Journal of Insect Biodiversity and Systematics*, 4 (1), 25–29.

Introduction

The family of Hypogastruridae Börner, 1906 is common and wide-spread and sprigtails with cosmopolitan distribution (Park & Park, 2006). This family consists of three main lineage of Ceratophysellan, Hypogastruran and Xenyllan (Thibaude *et al.*, 2004). *Xenylla* Tullberg, 1869 has 134 species with the highest number of species in the Xenillan lineage (Bellinger *et al.*, 2017). Members of this genus are usually dark and small or medium sized animals resembling *Hypogastrura*. They differ from other Hypogastruridae by the absence of Post-antennal Organ. Fourth antennal segment with simple apical bulb and six curved sensilla. Thoracic terga II and III without lateral microsensilla, chaeta m_2

absent. Retinaculum in various stages of reduction, never more than two setae on dens. Anal spines present, always shorter than claws (Fjellberg, 1998). Cox (1982) reported *Xenylla humicola* (Fabricius, 1780) and *Xenylla maritima* Tullberg, 1869 for the first time from Gilan, East and west Azarbaijan provinces for Iran fauna. Yahyapour (2012) reported the species *Xenylla welchi* Folsom, 1916 for the first time in Iran from Mazandaran and Falahati *et al.* (2012) from Kohgiluyeh and Boyer Ahmed. Shayanmehr *et al.* (2013) published a checklist of Iranian springtails which includes three species of *Xenylla*, including, *X. humicola* (Fabricius, 1780), *X. maritima* Tullberg, 1869 and *X. welchi*

Corresponding author: Leila Mohammadi Nodeheki, E-mail: mohammadi.leila991@gmail.com

Copyright © 2018, Mohammadi Nodeheki et al. This is an open access article distributed under the terms of the Creative Commons Attribution License (CC BY NC 4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Folsom, 1916. Yoosefi Lafooraki & Shayanmehr (2015) reported the *Xenylla maritima* Tullberg, 1869 for the first time in Mazandaran province. *Xenylla boernerii* Axelson, 1905 is recognized as a saffron bulb pest in South Khorasan (Tabadkani, 2016), but this has not been reported for the Iranian Collembola fauna. Özata *et al.* (2017) reported *X. mediterranea* da Gama, 1964 for the first time from Turkey.

Material and methods

The study area is located in the Amol Halomsar forest park, and the Babol Bezchaft forest park. Collembola specimens were collected from mosses on the tree in the Halomsar forest park (19 May, 2017) and mosses on the rock in the Bezchaft forest park (01 June, 2017). The species were extracted by Berlese Funnel and stored in 85% ethanol. After clearing, they fixed on the Hoyer medium for preparing microscopic slides and was identified by valid identification keys. Confirmation of genus and species was done by Dr. Dariusz Skarżyński (Poland).

Results

Material examined: 47 specimens, Halomsar forest park (N36°23'E52°20'), moss, 19.V.2017; 68 specimens, Bezchaft forest park (N36°22'E52°46'), moss, 01.VI.2017.

Description: Size 1 mm, colour bluish-grey (Fig 1A). 5+5 ommatidia (Fig 1B). Body hairs slightly longer and coarser, with relatively shorter lateral sensilla on thoracic and abdominal terga. Mandibles is developed (Fig 2A). Ventral tube with 4+4 setae (Fig 2B). Retinaculum with 2+2 teeth. Mucro smaller, not clearly set off from dens. Mucrodens significantly longer (Fig 3A). Dens with twosetae. Two anal spines (Fig 3B).

Distribution: Canary Islands, Corsica, Croatia, Greece, Italy, Morocco Sicily, Portugal, Scandinavia, Spain and Ukraine, (Fjellberg, 1998; Thibaud *et al.*, 2004) – New to Iran (Mazandaran: Amol, Babol) (this study). The specimens were collected on mosses and rock vegetation.




Figure 1. *Xenylla mediterranea* da Gama, 1964: **A.** General habitus, dorso-lateral view (Original); **B.** Ommatidia, 40x.




Figure 2. *Xenylla mediterranea* da Gama, 1964: **A.** Mandibles, 40x (Original); **B.** Ventral tube, 40x.




Figure 3. *Xenylla mediterranea* da Gama, 1964: **A.** Furca and retinaculum, 40x (Original); **B.** anal spines, 40x.

Discussion

The species was originally described as a subspecies of *X. brevisimilis* Stach, 1949. It is distinguished from it by the fact that the number of teeth in the retinaculum is 2+2 instead of 3+3 (Fjellberg, 1998). Babenko *et al.* (1994) found that *X. mediterranea* da Gama, 1964 has sublobal hair and seta m_5 present on Abdominal tergum IV, and ranked it as a full species. With our new records in this study, the number of *Xenylla* species known from Iran increased to a total of four. So far, the species *X. humicola*, *X. maritima*, *X. welchi* and *X. mediterranea* have been reported for Iran.

Acknowledgments

We thank the Sari Agricultural Sciences and Natural Resources University (SANRU), for providing the laboratory facilities for this study. We also thank Dr. Dariusz Szarzynski Zoological Institute of Wroclaw University to confirm identification.

Conflict of Interests

The authors declare that there is no conflict of interest regarding the publication of this paper.

References

- Babenko, A.B. Chernova, N.M., Potapov, M.B. & Stebaeva, S.K. (1994) *Collembola of Russia and adjacent countries: Family Hypogastruridae*. Moscow, Nauka, 336 pp.
- Bellinger, P.F., Christansen, K.A. & Janssens, F. (2017) *Checklist of the Collembola of the world*. Available from: <http://www.collembola.org> [Accessed 13 October, 2017].
- Cox, P. (1982) Collembola fauna of north and north western Iran. *Entomologists' monthly Magazine*, 118, 39–43.
- Falahati Hossein Abad, A., Skarzynski, D. & Alizadeh, S. (2012) The report of six new species from Hypogastruridae (Apterygota: Collembola) for Iran fauna. *Proceeding of the Twentieth Iranian plant protection Congress, 2012, 25-28 August, Shiraz University, Fars, Iran*, 1, 141–141.
- Fjellberg, A. (1998) *The Collembola of Fennoscandia and Denmark: Part I. Poduromorpha*. Brill, Leiden, Boston. pp. 50–56.
- Özata, M.A., Sevgili, H. & Kaprus, I.J. (2017) Poduromorpha fauna of Ordu province in Turkey (Hexapoda: Collembola). *Dış Kapak*, 57 (2), 231–250. <https://doi.org/10.16955/bitkorb.322379>
- Park, K.H. & Park, N.Y. (2006) Two new species of *Ceratophysella* (Collembola: Hypogastruridae) from Korea. *Florida Entomologist*, 89 (4), 489–496. [https://doi.org/10.1653/0015-4040\(2006\)89\[489:TNSOCC\]2.0.CO;2](https://doi.org/10.1653/0015-4040(2006)89[489:TNSOCC]2.0.CO;2)
- Shayanmehr, M., Yahyapour, E., Kahrarian, M. & Yoosefi Lafooraki, E. (2013) An introduction to Iranian Collembola (Hexapoda): an update to the species list. *ZooKeys*, 335, 69–83. <https://doi.org/10.3897/zookeys.335.5491>
- Tabadkani, S.M. (2016) *The Hexapoda (A color guide to insect identification)*. Tehran, Iranian student book agency. 380 pp. + 59pl.
- Thibaud, J.M., Schulz, H.J. & da Gama Assalino, M.M. (2004) *Synopses on Palaearctic Collembola, Vol. 4. Hypogastruridae*. Abhandlungen und Berichte des Naturkunde museums Görlitz, Germany.
- Yahyapour, E. (2012) *Faunistic Study on Collembola (Insecta: Apterygota) in Sari Regions*. MSc Dissertation, Department of plant protection, Sari University of Agricultural Sciences and Natural Resources (SANRU), Sari, 96 pp.
- Yoosefi Lafooraki, E. & Shayanmehr, M. (2015) First record of the genus *Schoettella* and three new records of the family Hypogastruridae (Collembola, Hexapoda) for Iranian fauna with an identification key for Mazandaran province. *Taxonomy and Biosystematics Journal*, 7 (23), 1–12.

گونه‌*Xenylla mediterranea* da Gama, 1964 (Collembola: Poduromorpha: Haypogastruride)، گزارش جدید برای فون ایران از استان مازنداران

لیلی محمدی نودهکی^{۱*}، معصومه شایان‌مهر^۱ و محسن یزدانیان^۲

۱ گروه گیاهپزشکی، دانشکده علوم زراعی، دانشگاه علوم کشاورزی و منابع طبیعی ساری، ساری، ایران.

۲ گروه گیاهپزشکی، دانشکده علوم زراعی، دانشگاه علوم کشاورزی و منابع طبیعی گرگان، گرگان، ایران.

* پست الکترونیکی نویسنده مسئول مکاتبه: mohammadi.leila991@gmail.com

تاریخ دریافت: ۱۴ آبان ۱۳۹۶، تاریخ پذیرش: ۲۶ دی ۱۳۹۶، تاریخ انتشار: ۲۸ دی ۱۳۹۶

چکیده: طی نمونه‌برداری از خانواده Hypogastruridae (Collembolla) در استان مازندران، گونه *Xenylla mediterranea* da Gama, 1964 برای اولین بار از ایران گزارش می‌شود. با ثبت این گونه، تعداد گونه‌های جنس *Xenylla* در ایران به چهار مورد افزایش یافت. اطلاعات مربوط به گونه، شامل مواد بررسی شده، توصیف، اکولوژی، پراکندگی و تصاویر خصوصیات مرفولوژیک ارائه شد.

واژگان کلیدی: *Xenylla mediterranea*, Hypogastruridae، پادمان، ایران، گزارش جدید