



ESI

Presence of ephedra gall midge, *Ephedromyia debilopalpis* Marikovskij (Diptera: Cecidomyiidae) in Iran

Omid Hadi¹, Hossein Lotfalizadeh^{2*}, Mohammad-Hossein Kazemi¹, and Marcela Skuhrová³

¹ Department of Plant Protection, College of Agriculture, Tabriz Branch, Islamic Azad University, Tabriz, Iran.

² Department of Plant Protection, East-Azarbaijan Agricultural and Natural Resources Research and Education Center, Agricultural Research, Education and Extension Organization (AREEO), Tabriz, Iran.

³ Bitovská 1227/9, CZ-140 00, Praha 4, Czech Republic.

Received:
17 September, 2018

Accepted:
28 October, 2018

Published:
10 November, 2018

Subject Editor:
Ebrahim Gilasian

ABSTRACT. We reared *Ephedromyia debilopalpis* Marikovskij (Dipt.: Cecidomyiidae) for the first time on *Ephedra major* (Ephedraceae) in the northwest of Iran in 2016. This is the new record of the genus and species for Iran. With including this new record, the known gall midge fauna of Iran reached 62 species and 34 genera.

Key words: Diptera, *Ephedromyia*, Iran, distribution, new record, Ephedraceae

Citation: Hadi, O., Lotfalizadeh, H., Kazemi, M.-H. & Skuhrova, M. (2018) Presence of ephedra gall midge, *Ephedromyia debilopalpis* Marikovskij (Diptera: Cecidomyiidae) in Iran. *Journal of Insect Biodiversity and Systematics*, 4 (2), 135–139.

Introduction

The family Ephedraceae includes only one genus, *Ephedra*, with 67 species which occur in Europe, Asia, North and South America. These plants inhabit both warm and cold biotops and may grow in deserts of Middle Asia and also at the seaside of the Black-Sea in Bulgaria. *Ephedra* species are spread far to the North and occur also at high altitudes in the mountain with 5400 m a.s.l. in the Himalaya Mts. in Asia and in the Andes in South America with 4700 m a.s.l. *Ephedra* species are often important components of sagebrush steppes which are formed of various *Artemisia* species. In the region Palaearctic, different species of *Ephedra* are widely distributed in the Mediterranean basin and in some part of

Asia (Ghahraman, 2006; Ghahraman & Attar, 1999; The Plant List, 2013).

Different herbivorous insects attacks *Ephedra*, that the gall midges (Dipt.: Cecidomyiidae) are one of them. The Cecidomyiidae with 6590 species in 812 genera in the world are one of the most specious families of Diptera (Gagné & Jaschhof, 2017). The known gall midge fauna of Iran is composed of 61 species in 33 genera which are associated with 50 plant species belonging to fifteen plant families (Skuhrová et al., 2014). Two gall midge species of the genus *Xerephedromyia* associated with *Ephedra major* were recorded in the Kerman province in Iran (Moeinadini et al., 2017).

Corresponding author: Hossein Lotfalizadeh, E-mail: hlotfalizadeh@gmail.com

Copyright © 2018, Hadi et al. This is an open access article distributed under the terms of the Creative Commons Attribution License (CC BY NC 4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Eleven species of gall midges are known to be associated with the host plants of the genus *Ephedra* in the world. Nine species of two genera (*Ephedromyia* and *Xerephedromyia*) which cause stem galls of various shape on *Ephedra* species in Asia, and two gall midge species of the genus *Lasioptera* which cause stem galls on *Ephedra* species in North America (Skuhrová, 1986; Gagné, 1989).

Material and methods

During our field collection of insect fauna associated with *Ephedra* in the northwest of Iran in 2016 (Hadi et al., 2017), we discovered stem galls of gall midge species belonging to the genus, *Ephedromyia*, which causes galls on stems of *Ephedra major* (Fig. 1B). Globular galls on stems of *E. major* were found at the locality Kordasht (geographic coordinates: 46°14'45.0"E & 38°52'11.0"N; at altitude of 551m a. s. l.) in the East-Azarbaijan province (Figs 1A & 2). Galls were collected on 19 April 2016 and transferred in polyethylen bag to the Entomology Laboratory of the Plant Protection Department, East-Azarbaijan Agricultural and Natural Resources Research Center, Tabriz.

Reared adults of gall midges were identified to the genus level using the key of Skuhrová (1997) and to the species level using the key of Fedotova (2000). The galls and specimens examined in this study are deposited in the insect collection of the Department of Plant Protection, East-Azarbaijan Research Center for Agriculture and Natural Resources, Tabriz, Iran, and in the gall midge collection of Marcela Skuhrová, Praha, Czech Republic.

Results

Galls on stems of *Ephedra major* were rounded, 9–13 mm in diameter, with brown unregular surface (Fig. 1D). Several adults emerged from galls in the polyethylen bag during May 2016.

Our finding of *Ephedromyia debilopalpis* (Fig. 1C) in this research is the first record of this species and genus in Iran. We enriched the gall midge fauna of Iran for one species and one genus. The present fauna of gall midges of Iran is composed of 62 species and 34 genera.

Discussion

Marikovskij (1953) found stem galls on *Ephedra* sp. in the mountains of Kazakhstan (Mount Chulak, Dzhungarskii Alatau) and described species as *Ephedromyia debilopalpis*. He described a male, female, gall and biology in detail. Only one generation develop per year. Adults emerge from galls in the spring and females lay eggs on young branches of *Ephedra*. Larvae develop inside stem and cause globular galls. They hibernate inside galls where they pupate in the spring of the following year. Fedotova (2000) gave as the host plant species *Ephedra lomatolepis* Schrenk and Kazakhstan as the distribution area; therefor, *Ephedra major* is a new host plant for this gall midge. Gagné and Jaschhof (2017) in their catalog gave that *E. debilopalpis* occurs also in the European part of Russia and in Turkmenistan but they do not give any reference supporting it. But Raymond Gagné in his letter of 12 September 2017 answered M.S. that he cannot find the reference for the geographic extensions of *Ephedromyia debilopalpis* in European part of Russia and Turkmenistan and that he will remove this information from the catalog.

Acknowledgments

We thank Department of Plant Protection, East-Azarbaijan Agricultural and Natural Resources Research and Education Center, for support of this research.

Conflict of Interests

The authors declare that there is no conflict of interest regarding the publication of this paper.




Figure 1. A) Collection locality in the northwest of Iran, B) *Ephedra major*, C) Adult of *Ephedromyia debilopalpis*, D) Two galls of *E. debilopalpis*, on stem of *E. major* in Iran.



Figure 2. Map of Iran including collection site of *Ephedromyia debilopalpis* in the northeast.

References

- Fedotova, Z.A. (2000) *Plant-Feeding Gall Midges (Diptera, Cecidomyiidae) of the Deserts and Mountains of Kazakhstan: Morphology, Biology, Distribution, Phylogeny, and Systematics*. Samara, 803 pp.
- Gagné, R.J. (1989) *The Plant-feeding Gall Midges of North America*. Ithaca and London: Comstock Publishing Associates, Cornell University Press, 356 pp.
- Gagné, R.J. & Jaschhof, M. (2017) *A Catalog of the Cecidomyiidae (Diptera) of the World*. Fourth Edition. Digital version. 762 pp.
- Ghahraman, A. (2006) *Flora of Iran. Flore de l'Iran en couleurs naturelles*. Vol. 25. Tehran: Institute des Recherches des Forests et des Paturage Département Botanique, 1176 pp.
- Ghahraman, A. & Attar, F. (1999) *Biodiversity of Plant Species in Iran*. Volume 1. Tehran: Tehran University Publications, 1176 pp.
- Hadi, O., Lotfalizadeh, H. & Kazemi, M.-H. (2017) New record of the genus and species of *Ksenoplata quadrata* Bouček (Hym.: Pteromalidae) on *Ephedra major* in Iran. *Applied Entomology and Phytopathology*, 84 (2), 359–361.
- Marikovskij, P.I. (1953) New genera and species of gall midges (Diptera, Cecidomyiidae) of southeastern Kazakhstan. *Izvestiya Akademia Nauk Kazakhskoi SSR, Seriya Biologicheskaya*, 8, 128–139.
- Moeinadini, A., Madjdzadeh, S.M. & Skuhrav, M. (2017) First records of two species of gall midges (Diptera: Cecidomyiidae) from Iran. *Journal of Insect Biodiversity and Systematics*, 3 (4), 309–318.
- Skuhravá, M. (1986) Family: Cecidomyiidae. In: Soós, Á. & Papp, L., (eds.) *Catalogue of Palaearctic Diptera*. Vol. 4. Budapest: Hungarian Academy of Sciences, Akadémiai Kiadó & Amsterdam: Elsevier, pp. 72–297.
- Skuhravá, M. (1997) Family Cecidomyiidae. In: Papp, L., Darvas, B., (eds.) *Contributions to a Manual of Palaearctic Diptera (with Special Reference to Flies of Economic Importance)*. Vol. 2. Nematocera and Lower Brachycera. Budapest: Science Herald, pp. 71–204.
- Skuhravá, M., Karimpour, Y., Sadeghi, H., Gol, A. & Joghataie, M. (2014) Gall midges (Dip.: Cecidomyiidae) of Iran: annotated list and zoogeographical analysis. *Acta Societatis Zoologicae Bohemicae*, 78, 269–301.
- The Plant List (2013) Version 1.1. Published on the Internet: <http://www.theplantlist.org/> (accessed 1st January).

حضور پشه گالزای افدرا *Ephedromyia debilopalpis* Marikovskij (Diptera: Cecidomyiidae) در ایران

امید هادی^۱، حسین لطفعلی زاده^{۲*}، محمدحسین کاظمی^۱ و مارسلا سکوراوا^۳

۱ گروه گیاه پزشکی، دانشکده کشاورزی، دانشگاه آزاد اسلامی تبریز، تبریز، ایران.
 ۲ بخش تحقیقات گیاه پزشکی، مرکز تحقیقات و آموزش کشاورزی و منابع طبیعی استان آذربایجان شرقی، سازمان تحقیقات، آموزش و ترویج کشاورزی، تبریز، ایران.
 ۳ بیئوسکا ۱۲۲۷، پراگ، جمهوری چک.

* پست الکترونیکی نویسنده مسئول مکاتبه: hlotfalizadeh@gmail.com

تاریخ دریافت: ۲۶ شهریور ۱۳۹۷، تاریخ پذیرش: ۰۶ آبان ۱۳۹۷، تاریخ انتشار: ۱۹ آبان ۱۳۹۷

چکیده: برای نخستین بار پشه گالزای افدرا *Ephedromyia debilopalpis* Marikovskij (Dipt.: Cecidomyiidae) از روی گونه *Ephedra major* متعلق به تیره Ephedraceae در شمال غرب ایران طی سال ۱۳۹۵ پرورش داده شد. این نخستین گزارش جنس و گونه از ایران می باشد. با افزوده شدن این گونه به فون ایران، تعداد پشه های گالزا در ایران به ۶۲ گونه و ۳۴ جنس می رسد.

واژگان کلیدی: دوبالان، *Ephedromyia*، ایران، پراکنش، گزارش جدید، Ephedraceae