



New records of the family Ichneumonidae (Hymenoptera, Ichneumonoidea) to the fauna of Iran

Abbas Mohammadi-Khoramabadi^{1*}, Hossein Lotfalizadeh² & Martin Schwarz³

¹ Department of Plant Production, Darab College of Agriculture and Natural Resources, Shiraz University, Darab, 74591-17666, I.R. Iran.

² Department of Plant Protection, East-Azarbaijan Agricultural and Natural Resources Research and Education Center, Agricultural Research, Education and Extension Organization (AREEO), Tabriz, Iran.

³ Eben 21, A-4202 Kirchschlag, Austria.

Received:
27 April, 2020

Accepted:
12 June, 2020

Published:
18 June, 2020

Subject Editor:
Ehsan Rakhshani

ABSTRACT. This study has been carried out to collect and identify Ichneumonidae species from northwestern Iran using Malaise traps during 2014. In total, fifteen species into 6 subfamilies are identified. Four species are recorded for the first time from Iran, i.e. *Hoplocryptus melanocephalus* (Gravenhorst, 1829), *Zoophthorus palpator* (Müller, 1776), *Exochus erythronotus* (Gravenhorst, 1820) and *Enicospilus cerebrator* Aubert, 1966. The geographical distribution data of the recorded species are provided.

Key words: Ichneumonidae, parasitoid, Iran, new record

Citation: Mohammadi-Khoramabadi, A., Lotfalizadeh, H. & Schwarz, M. (2020) New records of the family Ichneumonidae (Hymenoptera, Ichneumonoidea) to the fauna of Iran. *Journal of Insect Biodiversity and Systematics*, 6 (3), 229–237.

Introduction

Ichneumonidae (Hymenoptera: Ichneumonoidea) is the largest family of insects with more than 25285 described species in the world (Yu et al., 2016). In order to increase and complete knowledge on these abundant and important parasitoids of pests, researches on the their taxonomy, distribution and biology are ongoing throughout the world, even in some better studied areas such as Europe (Klopfstein et al., 2019; Vas, 2020).

Many studies in recent years have increased our knowledge on the fauna, distribution (Amiri et al., 2015a, 2015b, 2016a, 2016b; Hooshyar et al., 2018; Jahan et al., 2016; Mahyabadi et al., 2016; Mohammadi-Khoramabadi & Talebi, 2013; Mohammadi-Khoramabadi et al., 2013a; 2013b; Mohammadi et al., 2020; Riedel et al., 2019a; Sarafi et al., 2015; Shirzadegan et al., 2017, 2018a, 2018b) and bioecology of these important and abundant parasitoid wasps in forest, range and agricultural ecosystems of Iran (Akbarzadeh-Shoukat, 2012; Fathi et al., 2012; Kamangar et al., 2017; Mehrnejad & Basirat, 2009; Mohammadi-Khoramabadi et al., 2016a, 2016b, 2017; Pourhaji et al., 2016; Shamszadeh et al., 2015). Northwest of Iran is of great importance both in terms of animal genetic resources, which contains five national park and protection areas (Hanson et al., 2009; Zehzad et al., 2002) and in terms of its

Corresponding author: Abbas Mohammadi-Khoramabadi, E-mail: mohamadk@shirazu.ac.ir

Copyright © 2020, Mohammadi-Khoramabadi et al. This is an open access article distributed under the terms of the Creative Commons Attribution License (CC BY NC 4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

potency of agricultural products (Ebadzadeh et al., 2018). This region borders by Turkey, Azerbaijan and Armenia countries. The fauna of some subfamilies of Ichneumonidae in northwest of Iran is already studied and resulted in distributional extension of several species and also describing some new species to the world (Ghahari & Jussila, 2016; Pourhaji et al., 2016; Riedel & Aghadokht, 2017; Riedel et al., 2019a). In an ongoing collection and identification project of Iranian Ichneumonidae, here we present new data on the distribution of some Ichneumonidae species from northwest of Iran.

Material and methods

This study was carried out in East and West Azerbaijan provinces at northwest of Iran during July-August 2014. Sampling was done using Malaise traps (Table 1). The captured ichneumonid specimens were pinned or card mounted and then identified using relevant keys and descriptions available on Anomaloninae (Schnee, 2014), Campopleginae (Dbar, 1984; 1985; Kasparyan, 1981), Diplazontinae (Klopfstein, 2014), Metopiinae (Tolkanitz, 2007), Ophioninae (Broad & Shaw, 2016). The Cryptinae specimens were identified by the third author, Dr. M. Schwarz (A specialist on this subfamily). The general distribution of the identified species is followed Yu et al. (2016). The voucher specimens are deposited in the private collection of Dr. Schwarz (PCS) and Insect collection of Darab College of Agriculture and Natural Resources, Shiraz University (DCS).

Table 1. Geographical coordinates of sampling sites in northwest of Iran during 2014.

Malaise trap	Locality, Province	Geographical coordinates	Elevation a.s.l. (m)
M1	Basmanj, East Azerbaijan	46°28'17" N, 37°59'38" E	1748
M2	Kahriz, West Azerbaijan	44°58'15" N, 37°52'18" E	1370

Results

From a total of 176 captured specimens, distributional data of 15 species into 12 genera and 6 subfamilies are given as following. Four species are recorded for the first time from Iran, which are marked with an asterisk (*).

Subfamily Anomaloninae Viereck, 1918

Anomalon chinense (Kokujev, 1915)

Material examined: M2, 3♂♂ 1♀, 6–23.VI.2014, Leg. H. Lotfalizadeh; (DCS).

Distribution in Iran: Kermanshah (Zardouei-Heydari et al., 2020) and West Azerbaijan provinces (current study).

General distribution: Palaearctic (Yu et al., 2016).

Anomalon cruentatum (Geoffroy, 1785)

Material examined: M2, 12♂♂ 29♀♀, 6–23.VI.2014, Leg. H. Lotfalizadeh; (DCS).

Distribution in Iran: Ardabil, East Azerbaijan, Sistan and Baluchestan, Yazd (Barahoei et al., 2012), Kermanshah, North Khorasan, South Khorasan, Khorasan-e-Razavi, Isfahan (Zardouei-Heydari et al., 2020) and West Azerbaijan provinces (Current study).

General distribution: Palaearctic (Yu et al., 2016).

Subfamily Campopleginae Förster, 1869*Cymodusa australis* (Smits Van Burgst, 1913)**Material examined:** M1, 1♀ 2♂♂, 25.VII.2014–5.VIII.2014, Leg. H. Lotfalizadeh; (DCS).**Distribution in Iran:** Fars (Amiri et al., 2017) and East Azerbaijan provinces (current study).**General distribution:** Western Palaearctic (Yu et al., 2016).*Cymodusa longiterebra* Dbar, 1985**Material examined:** M1, 2♀♀ 2♂♂, 25.VII.2014–5.VIII.2014, Leg. H. Lotfalizadeh; (DCS).**Distribution in Iran:** Fars (Amiri et al., 2017) and East Azerbaijan provinces (current study).**General distribution:** Western Palaearctic (Yu et al., 2016).*Diadegma maculatum* (Gravenhorst, 1829)**Material examined:** M1, 1♀, 25.VII.2014–5.VIII.2014, Leg. H. Lotfalizadeh; (DCS).**Distribution in Iran:** West Azerbaijan (Pourhaji et al., 2016), Khorasan-e-Razavi (Ghahari et al., 2014) and East Azerbaijan provinces (Current study).**General distribution:** Western Palaearctic (Yu et al., 2016).**Subfamily Diplazontinae** Viereck, 1918*Homotropus nigritarsus* (Gravenhorst, 1829)**Material examined:** M2, 2♀♀, 6–23.VI.2014, Leg. H. Lotfalizadeh; (DCS).**Distribution in Iran:** Guilan, Tehran, Alborz, Mazandaran, Qazvin, Yazd, Isfahan, Fars, Ardabil, Khuzestan, Markazi (Mohammadi-Khoramabadi et al., 2013b) and West Azerbaijan provinces (Current study).**General distribution:** Palaearctic, Nearctic, Neotropical (Yu et al., 2016).*Diplazon laetatorius* (Fabricius, 1781)**Material examined:** M2, 1♀, 6–23.VI.2014, Leg. H. Lotfalizadeh; (DCS).**Distribution in Iran:** Alborz, Guilan, Mazandaran, Qazvin, Tehran (Mohammadi-Khoramabadi et al., 2013b), Chaharmahal and Bakhtiari (Nourbakhsh et al., 2008), Isfahan (Barahoei et al., 2015; Rakhshani et al., 2010), Kerman (Bakhtiari et al., 2014), Sistan and Baluchistan (Barahoei et al., 2013), Yazd (Mohammadi-Khoramabadi et al., 2016b).**General distribution:** Cosmopolite (Yu et al., 2016).**Subfamily Cryptinae** Kirby, 1837*Mesostenus transfuga* Gravenhorst, 1829**Material examined:** M2, 3♀♀, 6–23.VI.2014; M1, 1♀, 25.VII.2014–5.VIII.2014, Leg. H. Lotfalizadeh; (PCS).**Distribution in Iran:** Fars, Guilan, Isfahan, Khorasan-e-Razavi, Mazandaran (Mahyabadi et al., 2016), West and East Azerbaijan provinces (current study).**General distribution:** Palaearctic (Yu et al., 2016).

Hoplocryptus melanocephalus (Gravenhorst, 1829)*

Material examined: M2, 3♀, 6–20.VI.2014, Leg. H. Lotfalizadeh; (PCS).

Distribution in Iran: West Azerbaijan province (current study).

General distribution: Western Palaearctic (Yu et al., 2016).

Trychosis priesneri Rossem, 1971

Material examined: M2, 1♀, 6–23.VI.2014, Leg. H. Lotfalizadeh; (PCS).

Distribution in Iran: Isfahan, Khorasan-e-Razavi, Sistan and Baluchistan (Mahyabadi et al., 2016) and West Azerbaijan provinces (current study).

General distribution: Western Palaearctic (Yu et al., 2016).

Gelis areator (Panzer, 1804)

Material examined: M2, 1♀, 6–23.VI.2014, Leg. H. Lotfalizadeh; (PCS).

Distribution in Iran: Isfahan (Mahyabadi et al., 2016) and West Azerbaijan provinces (current study).

General distribution: Afrotropical (introduced), Oceanic, Oriental, Palaearctic (Schwarz, 2016; Yu et al., 2016).

Zoophthorus palpator (Müller, 1776)*

Material examined: M1, 1♀, 25.VII.2014–5.VIII.2014, Leg. H. Lotfalizadeh; (PCS).

Distribution in Iran: East Azerbaijan province (current study).

General distribution: Palaearctic (Yu et al., 2016).

Subfamily Metopiinae Förster, 1869*Exochus erythronotus* (Gravenhorst, 1820)*

Material examined: M1, 2♀ 3♂, 25.VII.2014–5.VIII.2014, Leg. H. Lotfalizadeh; (DCS).

Distribution in Iran: East Azerbaijan province (current study).

General distribution: Western Palaearctic (Yu et al., 2016).

Exochus gravipes (Gravenhorst, 1820)

Material examined: M1, 1♀, 25.VII.2014–5.VIII.2014, Leg. H. Lotfalizadeh; (DCS).

Distribution in Iran: East Azerbaijan province (current study).

General distribution: Holarctic (Yu et al., 2016).

Subfamily Ophioninae Shuckard, 1840*Enicospilus cerebrator* Aubert, 1966*

Material examined: M2, 1♀, 6–23.VI.2014, Leg. H. Lotfalizadeh; (DCS).

Distribution in Iran: West Azerbaijan province (current study).

General distribution: Western Palaearctic (Yu et al., 2016).

Discussion

This study added four new records, increasing the total number of Iranian Anomaloninae to 16, Campopleginae to 65, Cryptinae to 137, Diplazontinae to 19, Metopiinae to 11 and Ophioninae to 27 species (Table 2) (Amiri et al., 2015b, 2016a; Hooshyar et al., 2018; Mahyabadi et al., 2016; Mohammadi-Khoramabadi et al., 2013b; Mohammadi et al., 2020; Riedel et al., 2019a, 2019b; Schwarz, 2016; Zardouei-Heydari et al., 2020). Comparing the number of recorded species of the aforementioned subfamilies in Iran including the northwest provinces with the adjacent countries (Yu et al., 2016) (Table 2) show that more investigations are needed to obtain better knowledge on this family and unveil distribution range of Iranian Ichneumonidae.

Table 2. Current number of known species of the studied subfamilies of Ichneumonidae in the northwest of Iran and neighboring countries.

Subfamily	West Azer. prov.	East Azer. prov.	Northwest Iran	Iran	Azerbaijan	Turkey	Armenia
Anomaloninae	3	1	2	16	15	37	3
Campopleginae	11	5	15	65	67	124	29
Cryptinae	27	11	41	137	119	194	16
Diplazontinae	3	0	3	19	15	29	6
Metopiinae	0	2	3	11	16	58	13
Ophioninae	1	0	1	27	9	24	3

Acknowledgments

This work was supported by Shiraz University [grant number 95GRS0M2228]. We would like to express our cordially thanks to Anna Nuzhna (Schmalhausen Institute of Zoology, NAS of Ukraine, Kyiv, Ukraine) and Heinz Schnee (Birkenweg 18, D-04416 Markkleeberg) for their help with the identification of Anomaloninae species.

Conflict of Interests

The authors declare that there is no conflict of interest regarding the publication of this paper.

References

- Akbarzadeh-Shoukat, G. (2012) Larval parasitoids of *Lobesia botrana* (Denis and Schiffermüller, 1775) (Lepidoptera: Tortricidae) in Orumieh Vineyards. *Journal of Agricultural Science and Technology*, 14, 267–274.
- Amiri, A., Talebi, A.A., Jussila, R., Rakhshani, E. & Hajiqanbar, H. (2015a) A study of the Iranian Cremastinae (Hymenoptera: Ichneumonidae). *Journal of Insect Biodiversity and Systematics*, 1 (2), 87–100.
- Amiri, A., Talebi, A.A., Riedel, M., Rakhshani, E. & Hajiqanbar, H. (2015b) A survey of Metopiinae (Hymenoptera: Ichneumonidae) in southern Iran, with three new records. *Journal of Crop Protection*, 4 (4), 519–531.
- Amiri, A., Talebi, A., Jussila, R., Rakhshani, E. & Hajiqanbar, H. (2016a) Study of the subfamily Ophioninae (Hymenoptera: Ichneumonidae) in southern Iran. *Journal of Entomological Society of Iran*, 35 (4), 53–67.

- Amiri, A., Talebi, A., Rey Del Castillo, C., Rakhshani, E. & Hajiqaanbar, H. (2016b) Study of the genus *Lissonota* (Hymenoptera: Ichneumonidae: Banchinae) in southern Iran. *Journal of Entomological Society of Iran*, 36 (2), 89–99.
- Amiri, A., Talebi, A.A., Rakhshani, E. & Hajiqaanbar, H. (2017) First report of the genus *Cymodusa* (Ichneumonidae: Campopleginae) from Iran. *Journal of Insect Biodiversity and Systematics*, 3 (2), 81–89.
- Bakhtiari, F., Khayrandish, M. & Mohammadi-Khoramabadi, A. (2014) First record of *Homotropus elegans* (Hym.: Ichneumonidae, Diplazontinae) from Iran. *Journal of Entomological Society of Iran*, 34 (2), 77.
- Barahoei, H., Nader, E. & Rakhshani, E. (2015) A survey on Ichneumonidae of Isfahan province, central Iran. *Journal of Crop Protection*, 4 (2), 157–166.
- Barahoei, H., Rakhshani, E. & Riedel, M. (2012) A checklist of Ichneumonidae (Hymenoptera: Ichneumonidae) from Iran. *Iranian Journal of Animal Biosystematics*, 8 (2), 83–132.
- Barahoei, H., Schwarz, M., Kasparyan, D.R., Riedel, M., Khajeh, N. & Rakhshani, E. (2013) Contribution on the knowledge of Ichneumonidae (Hymenoptera) in the northern part of Sistan and Baluchistan province, Iran. *Acta Zoologica Bulgarica*, 65 (1), 131–135.
- Broad, G.R. & Shaw, M.R. (2016) The British species of *Enicospilus* (Hymenoptera: Ichneumonidae: Ophioninae). *European Journal of Taxonomy*, (187), 1–31. <http://dx.doi.org/10.5852/ejt.2016.187>
- Dbar, R.S. (1984) Review of the Palearctic species of the genus *Cymodusa* Holmgren (Hymenoptera, Ichneumonidae). *Entomological Review*, 63 (4), 127–138.
- Dbar, R.S. (1985) Revision of the Palearctic species of the genus *Cymodusa* Holmgren (Hymenoptera, Ichneumonidae). II. *Entomologicheskoe Obozrenie*, 64 (3), 588–600.
- Ebadzadeh, H.R., Ahmadi, K., Mohammadnia Afrouzi, S., Abbas-Taghani, R., Abbasi, M. & Yari, S. (2018) *Iran agriculture statistic*, Ministry of Jihad-e-Agriculture, Tehran. 420 pp.
- Fathi, S.A.A., Bozorg-Amirkalaei, M., Sarfraz, R.M. & Rafiee-Dastjerdi, H. (2012) Parasitism and developmental parameters of the parasitoid *Diadegma majale* (Gravenhorst) in control of *Plutella xylostella* (L.) on selected cultivars of canola. *BioControl*, 57 (1), 49–59. <https://doi.org/10.1007/s10526-011-9373-7>
- Ghahari, H. & Jussila, R. (2016) Contribution to the knowledge of the fauna of Ichneumonidae (Hymenoptera: Ichneumonoidea) from Iran. *Beiträge zur Entomologie*, 66 (1), 119–124.
- Ghahari, H., Ostovari, H., Jussila, R. & Behnood, S. (2014) A study on Ichneumonidae (Hymenoptera: Ichneumonoidea) from some regions of Khorasan province, north-eastern Iran. *Calodema*, 296, 1–2.
- Hanson, T., Brooks, T.M., Da Fonseca, G.A., Hoffmann, M., Lamoreux, J.F., Machlis, G., Mittermeier, C.G., Mittermeier, R.A. & Pilgrim, J.D. (2009) Warfare in biodiversity hotspots. *Conservation Biology*, 23 (3), 578–587. <https://doi.org/10.1111/j.1523-1739.2009.01166.x>
- Hooshyar, H., Vafaei-Shoushtari, R., Mohammadi-Khoramabadi, A., Goldasteh, S., Sanatgar, E. & Jussila, R. (2018) Study of the subfamilies Cryptinae and Ichneumoninae (Hymenoptera: Ichneumonidae) from Mazandaran province, with record of five species new to Iran. *Journal of Entomological Research*, 11 (1), 11–32.
- Jahan, F., Moghbeli-Gharaei, A., Hasanshahi, G., Mohammadi-Khoramabadi, A., Abbasipour, H. & Papp, J. (2016) First report of the parasitoid wasp *Dolichogenidea turkmene* (Hym.: Braconidae, Microgastrinae) from Iran. *Applied Entomology and Phytopathology*, 38 (2), 275–276.
- Kamangar, S., Lotfalizadeh, H., Mohammadi-Khoramabadi, A. & Seyedi-Sahebari, F. (2017) Parasitoids of *Tortrix viridana* L. in Kurdistan Province. *Iranian Journal of Forest and Range Protection Research*, 15 (2), 176–186. <https://doi.org/10.22092/ijfrpr.2018.109981.1264>
- Kasparyan, D.R. (1981) 11. Ctenopelmatinae. 12. Phrudinae. 13. Tersilochinae. 14. Cremastinae. 15. Campopleginae. 16. Ophioninae. *A guide to the insects of the European part of the USSR. Opredeliteli Fauna SSSR*, pp. 316–431.

- Klopfstein, S. (2014) Revision of the western Palaearctic Diplazontinae (Hymenoptera, Ichneumonidae). *Zootaxa*, 3801 (1), 1–143. <https://doi.org/10.11646/zootaxa.3801.1.1>
- Klopfstein, S., Riedel, M. & Schwarz, M. (2019) Checklist of ichneumonid parasitoid wasps in Switzerland (Hymenoptera, Ichneumonidae): 470 species new for the country and an appraisal of the alpine diversity. *Alpine Entomology*, 3, 51–81. <https://doi.org/10.3897/alpento.3.31613>
- Mahyabadi, M., Khayrandish, M., Takaloozadeh, H. & Barahoei, H. (2016) A checklist of Iranian Cryptinae (Hymenoptera: Ichneumonidae). *Journal of Insect Biodiversity and Systematics*, 2 (4), 449–466.
- Mehrnejad, M.R. & Basirat, M. (2009) Parasitoid complex of the pistachio twig borer moth, *Kermania pistaciella*, in Iran. *Biocontrol Science and Technology*, 19 (5), 499–510. <https://doi.org/10.1080/09583150902887610>
- Mohammadi-Khoramabadi, A., Kamangar, S.-A. & Lotfalizadeh, H. (2016a) Ichneumonid parasitoids of *Tortrix viridana* (Lepidoptera, Tortricidae) in the west of Iran. *Linzer Biologische Beiträge*, 48 (1), 681–691.
- Mohammadi-Khoramabadi, A., Lotfalizadeh, H. & Gharali, B. (2016b) A study on parasitoids of the hoverflies (Dipt.: Syrphidae) and their natural effects on them in organic aphid infested lettuce farms of Yazd Province, Iran. *Entomologia Generalis*, 107–115. <https://doi.org/10.1127/entomologia/2016/0131>
- Mohammadi-Khoramabadi, A., Pourhaji, A., Lotfalizadeh, H. & Riedel, M. (2017) *Chromoplex picticollis* (Hym.: Ichneumonidae) associated with Asteraceae capitulum: a genus and species new to Iran. *Journal of Entomological Society of Iran*, 37 (2), 283–284.
- Mohammadi-Khoramabadi, A. & Talebi, A.A. (2013) A study of the genus *Orthocentrus* (Hymenoptera: Ichneumonidae, Orthocentrinae) in Gilan and Tehran provinces of Iran, with first records of seven species and one subspecies. *Applied Entomology and Phytopathology*, 80 (2), 29–39.
- Mohammadi-Khoramabadi, A., Talebi, A.A. & Zwakhals, K. (2013a) A study of the subfamily Pimplinae (Hymenoptera: Ichneumonidae) in the north of Iran, with eleven new species records. *Entomofauna*, 34 (2), 29–56.
- Mohammadi-Khoramabadi, A., Talebi, A.A. & Zwakhals, K. (2013b) Study on Diplazontinae (Hymenoptera: Ichneumonidae) in the north central of Iran. *Journal of Crop Protection*, 2 (3), 241–261.
- Mohammadi, K.A., Riedel, M., Ziaaddini, M. & Asadi, A. (2020) New data on the fauna of Ichneumonidae from Kerman province with first records of two species to Iran. *Journal of Entomological Society of Iran*, 39 (4), 447–457. <https://doi.org/10.22117/jesi.2019.126001.1306>
- Nourbakhsh, S.H., Soleymannejadian, E. & Nemati, A.R. (2008) Biology and population dynamics of *Scaeva albomaculata* (Diptera: Syrphidae) in almond orchards of Shahrekord, Iran. *Journal of Entomological Society of Iran*, 27 (2), 93–108.
- Pourhaji, A., Lotfalizadeh, H., Farshbaf-Pourabad, R., Gharali, B. & Mohammadi-Khoramabadi, A. (2016) Ichneumonid parasitoids (Hymenoptera: Ichneumonidae) of fruit flies (Diptera: Tephritidae) in the northwest of Iran. *Journal of Insect Biodiversity and Systematics*, 2 (2), 193–202.
- Rakhshani, H., Ebadi, R., Hatami, B., Rakhshani, E. & Gharali, B. (2010) A survey of alfalfa aphids and their natural enemies in Isfahan, Iran, and the effect of alfalfa strip-harvesting on their populations. *Journal of Entomological Society of Iran*, 30 (1), 13–28.
- Riedel, M. & Aghadokht, P. (2017) Contribution to the Ichneumoninae (Hymenoptera: Ichneumonidae) of Iran, with descriptions of three new species. *Zoology in the Middle East*, 63 (4), 348–355. <https://doi.org/10.1080/09397140.2017.1361190>
- Riedel, M., Ameri, A., Talebi, A.A. & Ebrahimi, E. (2019a) Contribution to the Ichneumonidae (Hymenoptera) of Iran, with descriptions of seven new species. *Linzer Biologische Beiträge*, 51 (1), 361–390.

- Riedel, M., Mohammadi-Khoramabadi, A. & Khayrandish, M. (2019b) Two new species of Campopleginae (Hymenoptera: Ichneumonidae) from Iran. *Zoology in the Middle East*, 65 (3), 1–5. <https://doi.org/10.1080/09397140.2019.1615751>
- Sarafi, T., Barahoei, H., Madjdzadeh, S.M. & Askari, M. (2015) A contribution to the knowledge of the Ichneumonidae (Hym.: Ichneumonoidea) from Neyriz county of Fars province, Iran. *Journal of Crop Protection*, 4, 643–654.
- Schnee, H. (2014) Anomaloninae aus Israel und Palästina mit Beschreibung einer neuen Anomalon-Art (Hymenoptera: Ichneumonidae). *Entomologische Zeitschrift*, 124, 97–107.
- Schwarz, M. (2016) Die Schlupfwespengattung *Gelis* (Hymenoptera, Ichneumonidae, Cryptinae) mit macropteren Weibchen in der Westpaläarktis. *Linzer biologische Beiträge*, 48 (2), 1677–1752.
- Shamszadeh, M., Jafary, A. & Bishe, G. (2015) Natural parasitism of *Pharetophora iranica* (Hym.: Ichneumonidae) on saxaul seed moths larvae in Yazd province. *Iranian Journal of Forest and Range Protection Research*, 12 (2), 160–162. <https://doi.org/http://dx.doi.org/10.22092/ijfrpr.2015.13320>
- Shirzadegan, F., Talebi, A.A., Riedel, M. & Hajiqanbar, H. (2017) Two newly recorded species of the tribe Platylabini (Hymenoptera: Ichneumonidae, Ichneumoninae) from Iran. *Journal of Crop Protection*, 6 (3), 401–408.
- Shirzadegan, F., Talebi, A.A., Riedel, M. & Hajiqanbar, H. (2018a) Study of the tribe Heresiarchini (Hymenoptera: Ichneumonidae, Ichneumoninae) in northern Iran, with a new record for the Middle East. *Journal of Insect Biodiversity and Systematics*, 4 (2), 113–122.
- Shirzadegan, F., Talebi, A.A., Riedel, M. & Hajiqanbar, H. (2018b) Study of the tribe Phaeogenini (Hymenoptera: Ichneumonidae, Ichneumoninae) in northern Iran, with two new genera and four new species records for the fauna of Iran. *Journal of Entomological Society of Iran*, 37 (4), 417–432. <https://doi.org/10.22117/jesi.2018.116290.1158>
- Tolkanitz, V.I. (2007) Ichneumon flies of the genus *Exochus* Gravenhorst (Hymenoptera: Ichneumonidae: Metopiinae) of the fauna of Palaearctic region. *Russian Entomological Journal*, 16 (3), 339–358.
- Vas, Z. (2020) New species and records of Palaearctic Campopleginae, Cryptinae, Hybrizontinae, and Tersilochinae (Hymenoptera: Ichneumonidae). *Opuscula Zoologica*, 51 (1), 3–13. <https://doi.org/10.18348/opzool.2020.1.3>
- Yu, D.S., Van Achterberg, K. & Horstmann, K. (2016) Taxapad 2016 – World Ichneumonoidea 2015. Taxonomy, Biology, Morphology and Distribution Ontario: Nepean, Canada: On USB Flash Drive.
- Zardouei-Heydari, M., Rakhshani, E., Mokhtari, A. & Schwarz, M. (2020) The genus *Anomalon* Panzer (Hymenoptera: Ichneumonidae: Anomaloninae) of Iran, with description of a new species. *European Journal of Taxonomy*, 662, 1–19. <https://doi.org/10.5852/ejt.2020.662>
- Zehzad, B., Kiabi, B.H. & Madjnoonian, H. (2002) The natural areas and landscape of Iran: an overview. *Zoology in the Middle East*, 26, 7–10. <https://doi.org/10.1080/09397140.2002.10637915>

گزارش‌های جدید از خانواده Ichneumonidae (Hymenoptera, Ichneumonoidea) برای فون ایران

عباس محمدی خرم آبادی^{۱*}، حسین لطفعلی زاده^۲ و مارتین شوآرز^۳

۱ بخش تولیدات گیاهی، دانشکده کشاورزی و منابع طبیعی داراب، دانشگاه شیراز، ایران.

۲ بخش تحقیقات گیاهپزشکی، مرکز تحقیقات و آموزش کشاورزی و منابع طبیعی آذربایجان شرقی، تبریز، ایران.

۳ ابن ۲۱، آ-۴۲۰۲، کرکسلاگ، اتریش.

* پست الکترونیکی نویسنده مسئول مکاتبه: mohamadk@shirazu.ac.ir

| تاریخ دریافت: ۸ اردیبهشت ۱۳۹۹ | تاریخ پذیرش: ۲۳ خرداد ۱۳۹۹ | تاریخ انتشار: ۲۹ خرداد ۱۳۹۹ |

چکیده: این مطالعه به جمع‌آوری و شناسایی گونه‌های خانواده Ichneumonidae از شمال غرب ایران با استفاده از تله مالیز در سال ۱۳۹۳ پرداخته است. در مجموع، پانزده گونه از شش زیرخانواده شناسایی شدند. چهار گونه شامل *Hoplocryptus Zoophthorus palpator* (Müller, 1776) *Exochus erythronotus melanocephalus* (Gravenhorst, 1829) برای اولین بار از ایران گزارش می‌شوند. داده‌های پراکنش جغرافیایی برای گونه‌های گزارش شده ارائه شد.

واژگان کلیدی: Ichneumonidae، پارازیتوئید، ایران، گزارش جدید