



ESI

Casinaria Holmgren and *Dusona* Cameron (Hymenoptera: Ichneumonidae) in Iran: distribution extension

Abbas Mohammadi-Khoramabadi^{1*} & Matthias Riedel²

¹ Department of Plant Production, Darab College of Agriculture and Natural Resources, Shiraz University, Darab, 74591-17666, I.R. Iran.

² Zoologische Staatssammlung München, Münchhausenstr. 21, D-81247 München, Germany.

ABSTRACT. This study is carried out to present new species records and new distributional data of the genera *Casinaria* Holmgren and *Dusona* Cameron (Hymenoptera: Ichneumonidae, Campopleginae) from Iran and also to provide the illustrated taxonomic notes on their diagnostic morphological characters. Specimens were collected using Malaise traps during 2013, 2015 and 2016 from Fars, Kerman and Mazandaran provinces. Four species present here as *Casinaria kriechebaumeri* (Costa, 1884); *Casinaria trochanterator* Aubert, 1960; *Dusona cultrator* (Gravenhorst, 1829) and *Dusona erythrogaster* (Förster, 1868). Two species i.e. *D. cultrator* and *D. erythrogaster* are newly reported from Iran. Data about distribution of *D. rugifer* and *D. stygia* in Iran as part of eastern Palaearctic fauna is updated to the western Palaearctic realm, based on collecting sites in Iran (*D. rugifer* from Golestan province and *D. stygia* from West Azerbaijan province). Available data on adult flight period of Iranian species of *Casinaria* and *Dusona* may indicate that five species are univoltine and the others are bivoltine.

Key words: Distribution, parasitoid, taxonomy, new record, fauna

Received:
17 May, 2020

Accepted:
12 June, 2020

Published:
21 June, 2020

Subject Editor:
Seyed Masoud Majdzadeh

Citation: Mohammadi-Khoramabadi, A. & Riedel, M. (2020) *Casinaria* Holmgren and *Dusona* Cameron (Hymenoptera: Ichneumonidae) in Iran: distribution extension. *Journal of Insect Biodiversity and Systematics*, 6 (3), 239–246.

Introduction

Campopleginae is the third largest subfamily of Ichneumonidae (Hymenoptera: Ichneumonoidea) with more than 2127 described species in the world (Yu et al., 2016). Members of this subfamily are koinobiont endoparasitoids of mostly Lepidoptera larvae, although species of a few genera attack Coleoptera, Symphyta (Hymenoptera) and Raphidioptera (Neuroptera) larvae. Because of this, Campoplegines are widely investigated and considered for both classical and conservation biological control of pests (Quicke, 2015). In Iran, the first data on this subfamily comes from surveys on the parasitoids of some important pests (Gonzalez et al., 1980; Herard et al., 1979) but over past two decades, the faunal, taxonomical and bioecological researches have been considerably accelerated our information on this subfamily as up to now, 64 campoplegine species have been reported and 14 host-parasitoid relations have been

Corresponding author: Abbas Mohammadi-Khoramabadi, E-mail: mohamadk@shirazu.ac.ir

Copyright © 2020, Mohammadi-Khoramabadi & Riedel. This is an open access article distributed under the terms of the Creative Commons Attribution License (CC BY NC 4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

documented (Amiri et al., 2017; Barahoei et al., 2012; Fathi et al., 2012; Mohammadi-Khoramabadi et al., 2017, 2018; Mohammadi-Khoramabadi & Ziaaddini, 2017; Mohebban et al., 2016; Riedel et al., 2019a, 2019b; Shaw et al., 2016; Talebi et al., 2006).

Casinaria Holmgren, 1859 is a moderately large genus with 100 described species globally (Yu et al., 2016). The first recorded Iranian species of this genus was *C. tenuiventris* (Gravenhorst, 1829) which was reported as a larval parasitoid of *Lymantria dispar* Linnaeus, 1758 (Lep.: Lymantriidae) (Herard et al., 1979). Two other species have been recently added to the Iranian *Casinaria* records, i.e. *C. krieckbaumeri* (Costa, 1884) and *C. trochanterator* Aubert, 1960 (Riedel, 2018; Riedel et al., 2019a). *Dusona* Cameron, 1901 is a very large genus with more than 442 described species in the world (Yu et al., 2016) but only three species and very few collected specimens have been found in Iran so far, i.e. *D. rugifer* (Förster, 1868) (Hinz & Horstmann, 2004), *D. stygia* (Förster, 1868) (Horstmann, 2009) and *D. rufigaster* Riedel, 2019 (Riedel et al., 2019a). The aim of this study is to present new distributional records of these genera from Iran and also to provide some taxonomic notes with some photographs of their morphological characters.

Material and methods

Specimens were collected by Malaise trap during 2013, 2015 and 2016 in different regions of Iran and identified by M. Riedel. Digital photographs were taken using a Canon EOS 600D on a SZ-ST Olympus stereomicroscope. Stacking of photographs was done by Adobe Photoshop. Terminology followed Hinz & Horstmann (2004). The identified species are deposited in the private collection of Dr. M. Riedel (Bad Fallinghostel/ Germany) and insect collection of Darab college of Agriculture and Natural Resources, Shiraz University.

Results

Casinaria krieckbaumeri (Costa, 1884) (Figs 1-3)

Material examined: Iran, Mazandaran province, Amol county (36°27'14" N, 53°06'01" E, Elevation: 180 m a.s.l.), 1♂1♀, 14.VI-9.VII.2016.


Distribution in Iran: Khuzestan, Markazi (Riedel et al., 2019a) and Mazandaran provinces.

Distribution: Austria, Bulgaria, Croatia, France, Greece, Jordan, Kazakhstan, Kirgizstan, Malta, Slovakia, Spain, Tunisia, Turkey, Turkmenistan, Uzbekistan (Riedel, 2018) and Iran (Riedel et al., 2019a).

Diagnosis: In the recent key of West Palaearctic *Casinaria* (Riedel, 2018), it runs to couplets 9 and 15. The main diagnostic morphological characters of the Iranian specimens (Fig. 1) are: Mandibles black (Fig. 2); temple strongly narrowed behind eyes (Fig. 3); prepectal carina not elevated; propodeum with short basal carinae, coarsely rugose, area petiolaris with rather deep and narrow depression, the depression with transverse rugae; hind femur red, metasoma stout, apical tergites not strongly compressed, 2nd tergite 1.8–2.1 × as long as wide, 3rd tergite 0.85–1.1 × as long as wide (Fig. 1).

Casinaria trochanterator Aubert, 1960 (Figs 4-6)

Material examined: Iran, Fars province, Darab (28°45'04" N, 54°26'51" E, Elevation: 1107 m a.s.l.), 1♀, 7-20.X.2013; Kerman province, Maskoon (28°51'41" N, 57°52'10" E, Elevation: 1665 m a.s.l.), 1♂, 9-23.VI.2015.



Figures 1–6. *Casinaria* spp. **1–3.** *Casinaria kriechbaumeri*: **1.** habitus of male, **2.** face, **3.** head in dorsal view; **4–6.** *Casinaria trochanterator*: **4.** female habitus, **5.** head in dorsal view, **6.** propodeum, basal carina (arrow a) and area petiolaris (arrow b).

Distribution in Iran: Khuzestan, Markazi (Riedel et al., 2019a) and Fars provinces.

Distribution: Austria, Bulgaria, Czech (Moravia), Egypt, France, Germany, Hungary, Iran, Italy, Norway, Serbia, Slovakia, Spain, United Kingdom (Riedel, 2018).

Diagnosis: *C. trochanterator* runs to couplet 8 in the key (Riedel, 2018). This species (Fig. 4) is mainly characterized by: mandibles black; temple slowly narrowed behind eyes (Fig. 5); hind femur red; propodeum with basal carinae (area basalis and 1/3-1/2 of area superomedia) (Fig. 6, arrow a), area superomedia wider than long, area petiolaris slightly and deeply depressed and with rugae (Fig. 6, arrow b); prepectal carina lamelliform and strongly elevated at ventral; metasoma stout, apical tergites not strongly compressed, 2nd tergite 1.8–2.1 × as long as wide; thyridia longer than wide or roundish, distance from basal margins 1.5 × their length; 3rd tergite 0.9–1.0 × as long as wide (Fig. 4).


***Dusona cultrator* (Gravenhorst, 1829)** (Figs 7–10)

Material examined: Iran, Mazandaran province, Amol county (36°27'14" N, 53°06'01" E, Elevation: 180 m a.s.l.), 1♀, 8–25.IV.2016, Haftkhal (36°17'18" N, 53°23'43" E, Elevation: 861 m a.s.l.). 3♀♀, 26.IV–16.V.2016.

Distribution in Iran: Mazandaran province.

Distribution: Eastern and western Palaearctic (Yu et al., 2016) and Iran.

Diagnosis: This species can be separated from other species of the genus by a combination of the following characters: body length 12–16 mm (Fig. 7); antenna with 52–59 flagellomeres, antennal carina narrow (Fig. 8); hind femur red; hind tibia yellow (Fig. 7); propodeum deeply depressed, the sulcus with distinct transverse striae (Fig. 9, arrow a), area basalis sessile (Fig. 9, arrow b), area superomedia not bordered by carina; glymma present; metasoma black except for posterior half of the 2nd tergite and anterior half of the 3rd tergite red (Fig. 10).



Figures 7–13. *Dusona* spp. 7–10. *Dusona culterator*; 7. habitus of female, 8. face, antennal carina arrowed, 9. propodeum, arrows show area petiolaris (a) and area basalis (b), 10. metasomal tergites; 11–13. *Dusona erythrogaster*; 11. female habitus, 12. head and mesosoma in lateral view, genal carina arrowed, 13. metasoma.

Dusona erythrogaster (Förster, 1868) (Figs 11–13)

Material examined: Iran, Mazandaran province, Amol county (36°13'14" N, 53°39'24" E, Elevation: 1624 m a.s.l.), 15–25.V.2016.

Distribution in Iran: Mazandaran province.

Distribution: Western Palaearctic (Yu et al., 2016) and Iran.

Diagnosis: body length 7–8 mm (Fig. 11); genal carina joining the oral carina at the base of mandible (Fig. 12), oral carina not distinctly raised; antenna with 35–39 flagellomeres, antennal carina not raised; ventral part of the prepectal carina merge with the transverse carina, pleural part of prepectal carina obliterated; hind tibia black at base and apex, brown medially; glymma present; metasoma with the 0.2–0.3 of the 2nd tergite onwards red (Fig. 13).

Discussion

Distribution of the Iranian recorded species of the genera *Casinaria* and *Dusona* are shown in Fig. 14. Hinz & Horstmann (2004) reported *D. rugifer* and *D. stygia* from Iran as part of eastern Palaearctic fauna, but based on collecting sites in Iran (*D. rugifer* from Golestan province and *D. stygia* from west Azerbaijan province) as well as other countries (Yu et al., 2016), they are actually belonging to the western Palaearctic realm (Fig. 14 g and h).




Figure 14. Distribution map in Iran. a. *Casinaria kriechebaumeri* b. *C. tenuiventris*; c. *C. trochanterator*; d. *Dusona cultrator*; e. *D. erythrogaster*; f. *D. rufigaster*; g. *D. rugifer*; h. *D. stygia*.

The flight period of Iranian species of *Casinaria* and *Dusona* occurred in one (*C. kriechbaumeri*, *C. tenuiventris*, *D. cultrator*, *D. erythrogaster*, *D. rufigaster* and *D. rugifer*) and two (*C. trochanterator*) periods, respectively (Table 1). Our data may indicate that *C. kriechbaumeri*, *C. tenuiventris*, *D. cultrator*, *D. erythrogaster* and *D. rugifer* are univoltine and the others bivoltine (Horstmann, 2011).

Table 1. The flight period of Iranian species of *Casinaria* and *Dusona*.

Species	April	May	June	July	August	September	October
<i>C. kriechbaumeri</i>		■					
<i>C. tenuiventris</i>		■					
<i>C. trochanterator</i>		■					■
<i>D. cultrator</i>	■	■					
<i>D. erythrogaster</i>		■					
<i>D. rufigaster</i>					■		
<i>D. rugifer</i>		■					

Acknowledgments

We wish to thank H. Hooshyar for her help with us in collecting specimens. This work was supported by Shiraz University [grant number 96GRS0M2228].

Conflict of Interests

The authors declare that there is no conflict of interest regarding the publication of this paper.

References

- Amiri, A., Talebi, A.A., Rakhshani, E. & Hajiqanbar, H. (2017) First report of the genus *Cymodusa* (Ichneumonidae: Campopleginae) from Iran. *Journal of Insect Biodiversity and Systematics*, 3 (2), 81–89.
- Barahoei, H., Rakhshani, E. & Riedel, M. (2012) A checklist of Ichneumonidae (Hymenoptera: Ichneumonidae) from Iran. *Iranian Journal of Animal Biosystematics*, 8 (2), 83–132.
- Fathi, S.A.A., Bozorg-Amirkalaei, M., Sarfraz, R.M. & Rafiee-Dastjerdi, H. (2012) Parasitism and developmental parameters of the parasitoid *Diadegma majale* (Gravenhorst) in control of *Plutella xylostella* (L.) on selected cultivars of canola. *BioControl*, 57 (1), 49–59.
<https://doi.org/10.1007/s10526-011-9373-7>
- Gonzalez, D., Etzal, L., Esmaili, M., El-Heneidy, A.H. & Kaddou, I. (1980) Distribution of *Bathyplectes curculionis* and *Bathyplectes anurus* (Hym.: Ichneumonidae) from *Hypera* (Col.: Curculionidae) on alfalfa in Eagypt, Iraq and Iran. *Entomophaga*, 25 (2), 111–121.
<https://doi.org/10.1007/BF02374313>
- Herard, F., Mercadier, G. & Abai, M. (1979) Situation de *Lymantria dispar* [Lep.: Lymantriidae] et de son complexe parasitaire en Iran, en 1976. *Entomophaga*, 24 (4), 371–384.
<https://doi.org/10.1007/BF02374176>
- Hinz, R. & Horstmann, K. (2004) Revision of the eastern Palearctic species of *Dusona* Cameron: (Insecta, Hymenoptera, Ichneumonidae, Campopleginae). *Spixiana*, Supplement 29, 1–188.
- Horstmann, K. (2009) Revision of the western Palearctic species of *Dusona* Cameron (Hymenoptera, Ichneumonidae, Campopleginae). *Spixiana*, 32 (1), 45–110.

- Horstmann, K. (2011) Verbreitung und Wirte der *Dusona*-Arten in der Westpalaearktis (Hymenoptera, Ichneumonidae, Campopleginae). *Linzer biologische Beiträage*, 43 (2), 1295–1330.
- Mohammadi-Khoramabadi, A., Hesami, S. & Shafiei, S. (2018) First report of *Campoletis scyticus* (Hym.: Ichneumonidae, Campopleginae) from Iran. *Applied Entomology and Phytopathology*, 85 (2), 277–278.
- Mohammadi-Khoramabadi, A., Pourhaji, A., Lotfalizadeh, H. & Riedel, M. (2017) *Chromoplex picticollis* (Hym.: Ichneumonidae) associated with Asteraceae capitulum: a genus and species new to Iran. *Journal of Entomological Society of Iran*, 37 (2), 283–284.
- Mohammadi-Khoramabadi, A. & Ziaaddini, M. (2017) Report of *Diadegma semiclausum* (Hym.: Ichneumonidae, Campopleginae) parasitoid of the pistachio fruit hull borer moth, *Arimania komaroffi* (Lep.: Pyralidae) from Rafsanjan, Iran. In: Golchini, H., Sabouri, H. & Taliei, F. (eds.) *The Second National Congress of Monitoring and Forecasting in Plant Protection. Gonbad Kavous, Golestan province, Iran: Gonbad Kavous University.*
- Mohebban, S., Barahoei, H., Takaloozadeh, H.M., Madjdzadeh, S.M. & Riedel, M. (2016) A survey of the Ichneumonidae (Hymenoptera, Ichneumonoidea) of Kerman province, south-east Iran. *Journal of Insect Biodiversity and Systematics*, 2 (4), 419–437.
- Quicke, D.L. (2015) *The Braconid and Ichneumonid Parasitoid Wasps: Biology, Systematics, Evolution and Ecology*. Wiley Blackwell, Chichester. 681 pp.
- Riedel, M. (2018) Revision of the Western Palaearctic species of the genus *Casinaria* Holmgren (Hymenoptera, Ichneumonidae, Campopleginae). *Linzer biologische Beiträge*, 50 (1), 687–716.
- Riedel, M., Ameri, A., Talebi, A.A. & Ebrahimi, E. (2019a) Contribution to the Ichneumonidae (Hymenoptera) of Iran, with descriptions of seven new species. *Linzer Biologische Beiträge*, 51 (1), 361–390.
- Riedel, M., Mohammadi-Khoramabadi, A. & Khayrandish, M. (2019b) Two new species of Campopleginae (Hymenoptera: Ichneumonidae) from Iran. *Zoology in the Middle East*, 65 (3), 1–5. <https://doi.org/https://doi.org/10.1080/09397140.2019.1615751>
- Shaw, M.R., Horstmann, K. & Whiffin, A.L. (2016) Two hundred and twenty-five species of reared western Palaearctic Campopleginae (Hymenoptera: Ichneumonidae) in the National Museums of Scotland, with descriptions of new species of *Campoplex* and *Diadegma*, and records of fifty-five species new to Britain. *Entomologist's Gazette*, 67 (3), 177–222.
- Talebi, A.A., Rakhshani, E., Daneshvar, S., Fathipour, Y., Moharramipour, S. & Horstman, K. (2006) Report of *Campoplex tumidulus* and *Itopectis tunetana* (Hym.: Ichneumonidae), parasitoids of *Yponomeuta malinellus* Zell. (Lep.: Yponomeutidae) from Iran. *Applied Entomology and Phytopathology*, 73 (1), 134.
- Yu, D.S., Van Achterberg, K. & Horstmann, K. (2016) *Taxapad 2016 – World Ichneumonoidea 2015. Taxonomy, Biology, Morphology and Distribution*. Ontario: Nepean, Canada. Database on Flash Drive.

جنس‌های *Casinaria* Holmgren و *Dusona* Cameron (Hymenoptera: Ichneumonidae) در ایران: گسترش پراکنش

عباس محمدی خرم آبادی^{۱*} و ماتیس ریدل^۲

۱ بخش تولیدات گیاهی، دانشکده کشاورزی و منابع طبیعی داراب، دانشگاه شیراز، ایران.

۲ موزه جانورشناسی مونیخ، ۲۱، د-۸۱۲۴۷، آلمان.

* پست الکترونیکی نویسنده مسئول مکاتبه: mohamadk@shirazu.ac.ir

| تاریخ دریافت: ۲۸ اردیبهشت ۱۳۹۹ | تاریخ پذیرش: ۲۳ خرداد ۱۳۹۹ | تاریخ انتشار: ۰۱ تیر ۱۳۹۹ |

چکیده: این مطالعه به منظور معرفی گزارش‌های جدید و نیز گزارش پراکنش جدید از گونه‌های جنس‌های *Casinaria* Holmgren و *Dusona* Cameron (Hymenoptera: Ichneumonidae, Campopleginae) از ایران و همچنین رایه توضیحات تاکسونومیک مصور از ویژگی‌های ریخت‌شناسی افتراقی آنها، انجام شد. نمونه‌ها بوسیله تله مالیز طی سال‌های ۱۳۹۲ و ۱۳۹۵ از استان‌های فارس، کرمان و مازندران جمع‌آوری شدند. چهار گونه شامل *Casinaria kriechebaumeri* (Costa, 1884)، *Dusona trochanterator* Aubert, 1960، *Dusona erythrogaster* (Förster, 1868) و *cultrator* (Gravenhorst, 1829) معرفی شدند. دو گونه شامل *D. erythrogaster* و *D. cultrator* برای اولین بار از ایران گزارش می‌شوند. پراکنش دو گونه *D. rugifer* و *D. stygia* گزارش شده از ایران به عنوان بخشی از فون شرق پالئارکتیک بر اساس محل جمع‌آوری آنها (*D. rugifer* از استان گلستان و *D. stygia* از استان آذربایجان غربی) به فهرست فون غرب پالئارکتیک تغییر کرد. داده‌های موجود از دوره پرواز حشرات بالغ گونه‌های ایرانی جنس‌های *Casinaria* و *Dusona* بیان‌گر این است که پنج گونه از جنس‌های مذکور دارای یک نسل و سایرین دو نسل در سال دارند.

واژگان کلیدی: پراکنش، پارازیتوئید، تاکسونومی، گزارش جدید، فون