



Subfamily Alysiinae Leach, 1815 (Hymenoptera: Braconidae) from Iran: new records and updated list

Elena Cortés¹, Ali Ameri², Ali Asghar Talebi², Ehsan Rakhshani³ and Francisco Javier Peris-Felipo^{4*}

¹ Fernando el Católico, 7, 1A, CP-03804, Alcoy, Alicante, Spain.

² Department of Entomology, Faculty of Agriculture, Tarbiat Modares University, P.O. Box: 14115-336, Tehran, Iran.

³ Department of Plant Protection, College of Agriculture, University of Zabol, P.O. Box: 98615-538, I.R. Iran.

⁴ Bleichestrasse 15, CH-4058 Basel, Switzerland.

ABSTRACT. The subfamily Alysiinae (Hymenoptera, Braconidae) is one of the most peculiar and diverse endoparasitoids of the family Braconidae. A sampling survey was done in different parts of Iran, mainly in South, Central and South Eastern provinces. Fifteen species are recorded for the first time from Iran. They were belonging to tribe Alysiini (*Dinotrema amparoae* Peris-Felipo, 2013; *D. borzhomii* Tobias, 2004; *D. dimorpha* (Fischer, 1976); *D. perlustrandum* (Fischer, 1973); *D. sinecarinum* (Fischer, 1993); *D. tauricum* (Telenga, 1935); *Orthostigma longicorne* Konigsmann, 1969; *Phaenocarpa bicolor* (Foerster, 1863); *P. carinthiaca* Fischer, 1975 and *Synaldis maxima* Fischer, 1972) and Dacnusini (*Chorebus melanophytiae* Griffiths, 1968; *C. thusa* (Nixon, 1937); *Dacnusa areolaris* (Nees, 1811); *D. aquilegiae* Marshall, 1896 and *D. metula* (Nixon, 1954)). List of Iranian Alysiinae including 108 species is also presented.

Key words: Alysiinae, updated list, new records, Iran.

Received:
02 November, 2016

Accepted:
12 December, 2016

Published:
14 December, 2016

Subject Editor:
Cornelis
van Achterberg

Citation: Cortés, E., Ameri, A., Talebi, A.A., Rakhshani, E. and Peris-Felipo, F.J. 2016. Subfamily Alysiinae Leach, 1815 (Hymenoptera: Braconidae) from Iran: new records and updated list. *Journal of Insect Biodiversity and Systematics*, 2(4): 411–418.

Introduction

The subfamily Alysiinae is one of the most peculiar and diverse endoparasitoid subfamilies of the family Braconidae (Yu *et al.* 2012). Morphologically, the two tribes are mainly separated by the presence (Alysiini) or absence (Dacnusini) of the fore wing vein r-m (Yu *et al.* 2012; Peris-Felipo *et al.* 2014). Alysiini are parasitoids of a wide range of Diptera-Cyclorrhapha while Dacnusini are exclusively specialized on leaf and stem miners of the families

Agromyzidae, Ephydriidae and Chloropidae (Belokobylskij 2005; Fischer & Beyarslan 2012; Peris-Felipo *et al.* 2014).

Recently, the number of Alysiinae known from Iran has reached to 93 species (Gadallah *et al.* 2015, Farahani *et al.* 2016, Peris-Felipo *et al.* 2016, and Yari *et al.* 2016). In the present paper, new records and an updated list of Alysiinae from Iran are provided.

Corresponding author: Francisco Javier Peris-Felipo, E-mail: peris.felipo@gmail.com

Copyright © 2016, Cortés *et al.* This is an open access article distributed under the terms of the Creative Commons Attribution License (CC BY 4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Material and methods

Specimens were collected using Malaise traps in different provinces of Iran during 2009–2013. They were dropped directly into ethyl alcohol (75%) for subsequent studies. The criteria indicated by Tobias (1986), Fischer (1990, 1993), van Achterberg (1993), Sharkey & Wharton (1997), and Peris-Felipo *et al.* (2014) were used for identification of new records. Classification and distribution follows Yu *et al.* (2012) and Peris-Felipo *et al.* 2014. New records of species are marked by an asterisk (*).

The material examined is deposited in the Department of Plant Protection at University of Zabol (Iran; DPPZ), Entomological Collection at University of Valencia (Spain; ENV), and Department of Entomology of Tarbiat Modares University (Tehran, Iran; TMUC). An updated list of all recorded Alysiinae species from different parts of Iran (Fig. 1) is presented.

Results

A total of 108 species belonging to 18 genera, *Adelurola* Strand, 1928, *Alloea* Haliday, 1833, *Alysia* Latreille, 1804, *Angelovia* Zaykov, 1980, *Aphaereta* Foerster, 1863, *Aristelix* Nixon, 1943, *Aspilota* Foerster, 1863, *Chorebus* Haliday, 1833, *Coelinidea* Viereck, 1913, *Coloneura* Foerster, 1863, *Dacnusa* Haliday, 1833, *Dinotrema* Foerster, 1863, *Idiasta* Foerster, 1863, *Orthostigma* Ratzeburg, 1844, *Phaenocarpa* Foerster, 1863, *Protodacnusa* Griffiths, 1964, *Pseudopezomachus* Mantero, 1905 and *Synaldis* Foerster, 1863 are recorded and catalogued for Iran. New records with material examined and distribution are provided below. Finally, a checklist of Iranian species of Alysiinae is given.

Class Hexapoda Blainville, 1816

Order Hymenoptera L., 1758

Family Braconidae Nees, 1811

Subfamily Alysiinae Leach, 1815

Tribe Alysiini Leach, 1815

**Dinotrema amparoae* Peris-Felipo, 2013

Material examined: 1 ♀, Iran, Hormozgan, Tezerj, 27°17'51.81"N 55°45'14.76"E, 867 m, 29.viii.2013 (A. Ameri leg.) (TMUC).

Distribution: Palaearctic.

**Dinotrema borzhomii* Tobias, 2004

Material examined: 1 ♀, Iran, Kerman, Lalezar, 29°31'52.57"N 56°47'49.93"E, 2701 m, 22.vi.2013 (A. Ameri leg.) (TMUC).

Distribution: Eastern Palaearctic.

**Dinotrema dimorpha* (Fischer, 1976)

Material examined: 2 ♂♂, Iran, Hormozgan, Tezerj, 27°17'51.81"N 55°45'14.76"E, 877 m, 25.i.2013 (A. Ameri leg.) (TMUC).

Distribution: Palaearctic.

**Dinotrema perlustrandum* (Fischer, 1973)

Material examined: 1 ♀, Iran, Sistan-& Baluchestan, Zahak, 30°54'07"N, 61°40'24"E, 491m, 23.viii.2009, swept on *Medicago sativa* L. (N. Khajeh leg) (ENV).

Distribution: Palaearctic.

**Dinotrema sinecarinum* (Fischer, 1993)

Material examined: 1 ♂, Iran, Kerman, Delfard, 28°57'59.23"N, 57°38'34.71"E, 2066 m, 19.iv.2013 (A. Ameri leg.) (TMUC).

Distribution: Palaearctic.

**Dinotrema tauricum* (Telenga, 1935)

Material examined: 1 ♀, Iran, Isfahan, Najafabad, 32°39'29.87"N 51°25'27.47"E, 1654 m, 6.vi.2012 (E. Nader leg.) (ENV).

Distribution: Palaearctic.

**Orthostigma longicorne* Königsmann, 1969

Material examined: 1 ♂, Iran, Kerman, Lalezar, 29°31'52.57"N 56°47'49.93"E, 2701 m, 22.vi.2013 (A. Ameri leg.) (TMUC).

Distribution: Palaearctic.

**Phaenocarpa* (*Phaenocarpa*) *bicolor*
(Foerster, 1863)

Material examined: 1 ♀, 1 ♂, Iran, Kerman, Bardsir, 29°57'23.83"N 56°34'33.223"E, 2026m, 16.iv.2013, swept on *Medicago sativa* L. (M. Azad leg.) (DPPZ).

Distribution: Palaearctic.

**Phaenocarpa* (*Phaenocarpa*) *carinthiaca*
Fischer, 1975

Material examined: 1 ♀, Iran, Kerman, Delfard, 28°57'59.23"N, 57°38'34.72"E, 2067 m, 30.vi.2013 (A. Ameri leg.) (TMUC).

Distribution: Palaearctic.

**Synaldis maxima* Fischer, 1972

Material examined: 1 ♂, Iran, Kerman, Lalezar, 29°31'52.57"N 56°47'49.93"E, 2701 m, 22.vi.2013 (Ameri leg.) (TMUC).

Distribution: Palaearctic.

Tribe *Dacnusini* Foerster, 1863

**Chorebus* (*Stiphocera*) *melanophytobiae*
Griffiths, 1968

Material examined: 1 ♀, Iran, Kerman, Delfard, 28°57'59.23"N, 57°38'34.71"E, 2068 m, 1.vii.2013 (A. Ameri leg.) (TMUC); 1 ♀, 1 ♂, Iran, Fars, Jahrom, 28°29'23.07"N 53°31'20.12"E, 1105 m, 4.vi.2013 (A. Ameri leg.) (TMUC); 1 ♂, same locality but, 28°34'01.62"N 53°27'29.53"E, 1540 m, 22.iv.2013 (A. Ameri leg.) (TMUC).

Distribution: Palaearctic.

**Chorebus* (*Chorebus*) *thusa* (Nixon, 1937)

Material examined: 1 ♂, Iran, Fars, Jahrom, 28°34'01.62"N 53°27'29.53"E, 1105 m, 10.i.2012 (A. Ameri leg.) (TMUC); 1 ♂, Iran, Hormozgan, Abmah, 27°46'1.76"N 56°43'30.87"E, 643 m, 12.iv.2013 (A. Ameri leg.) (TMUC); 1 ♂, Iran, Hormozgan, Bahne, 27°53'7.32"N 56°19'58.34"E, 1020 m, 4.xii.2012 (A. Ameri leg.) (TMUC); 1 ♀, 1 ♂, Iran, Hormozgan, Chelo, 27°10'30.39"N

57°1'9.79"E, 16 m, 23.xi.2012 & 22.i.2013 (A. Ameri leg.) (TMUC); 1 ♀, Iran, Hormozgan, Goleshvar, 27°58'30.57"N 56°59'53.55"E, 14 m, 29.iii.2011 (A. Ameri leg.) (TMUC); 2 ♂♂, Iran, Hormozgan, Rahbari, 27°27'2.67"N 57°8'54.22"E, 213 m, 25.iii.2011 & 1.iv.2013 (A. Ameri leg.) (TMUC); 1 ♂, Iran, Hormozgan, Ramkan, 26°52'25.27"N 56°1'7.33"E, 34 m, 17.iv.2013 (A. Ameri leg.) (TMUC); 1 ♂, Iran, Hormozgan, Tezerj, 27°17'51.81"N 55°45'14.76"E, 867 m, 8.iv.2013 (A. Ameri leg.) (TMUC); 2 ♂♂, Iran, Hormozgan, Zakin, 27°28'53"N 56°18'27"E, 632-687 m, 3.ii & 7.v.2013 (A. Ameri leg.) (TMUC); 1 ♂, Iran, Kerman, Lalezar, 29°31'52"N 56°47'49"E, 2701 m, 25.vi.2013 (A. Ameri leg.) (TMUC).

Distribution: Palaearctic.

**Dacnusa* (*Dacnusa*) *areolaris* (Nees, 1811)

Material examined: 3 ♀♀, 4 ♂♂, Iran, Hormozgan, Bahne, 27°53'7.32"N 56°19'58.34"E, 1021 m, 12.iv.2013 (A. Ameri leg.) (TMUC); 1 ♂, Iran, Hormozgan, Tezerj, 27°17'51.81"N 55°45'14.76"E, 879 m, 25.i.2012 (A. Ameri leg.) (TMUC); 1 ♀, Iran, Kerman, Delfard, 28°57'59.23"N 57°38'34.71"E, 2067 m, 1.vii.2013 (A. Ameri leg.) (TMUC); 1 ♀, 1 ♂, Iran, Kerman, Lalezar, 29°31'52.57"N, 56°47'49.93"E, 2700 m, 27.vi.2013 (A. Ameri leg.) (TMUC).

Distribution: Palaearctic.

**Dacnusa* (*Pachysema*) *aquilegiae* Marshall,
1896

Material examined: 1 ♀, Iran, Fars, Dejh, 30°55'42.46"N 52°43'46.85"E, 2100 m, 10.vii.2012 (A. Ameri leg.) (TMUC); 1 ♂, Iran, Kerman, Lalezar, 29°31'52.57"N 56°47'49.93"E, 2700 m, 27.vi.2013 (A. Ameri leg.) (TMUC); 2 ♀♀, 1 ♂, Iran, Hormozgan, Tezerj, 27°17'51.81"N 55°45'14.76"E, 880 m, 25.i.2012 (A. Ameri leg.) (TMUC); 5 ♀♀, 6 ♂♂, Iran, Hormozgan, Zakin, 27°28'53.23"N 56°18'27.03"E, 682 m, 11.v., 6.vii & 10.xi.2012 (A. Ameri leg.) (TMUC).

Distribution: Palaearctic.




Figure 1. Province distribution of alysiine records in Iran.

**Dacnusa (Pachysema) metula* (Nixon, 1954)

Material examined: 2 ♀♀, Iran, Hormozgan, Zakin, 27°28'53.23"N 56°18'27.03"E, 685 m, 24.v & 10.xi.2012 (A. Ameri leg.) (TMUC).

Distribution: Palaearctic.

**Updated list of Iranian Alysiinae species
Tribe Alysiini Leach, 1815**

1. *Adelurola amplidens* (Fischer, 1966)
2. *Alloea contracta* (Haliday, 1833)
3. *Alysia (Anarcha) rufidens* Nees, 1834
4. *Angelovia elipsocubitalis* Zaykov, 1980
5. *Aphaereta difficilis* Nixon, 1939
6. *Aphaereta minuta* (Nees, 1811)
7. *Aspilota alfalfae* Fischer, Lashkari Bod, Rakhshani & Talebi, 2011
8. *Aspilota delicata* Fischer, 1973
9. *Aspilota flagimilis* Fischer, 1996
10. *Aspilota insolita* (Tobias, 1962)
11. *Aspilota isfahanensis* Peris-Felipo, 2016
12. *Aspilota latitemporata* Fischer, 1976
13. *Aspilota nidicola* Hedqvist, 1972
14. *Dinotrema amoenidens* (Fischer, 1973)
15. *Dinotrema amparoae* Peris-Felipo, 2013*
16. *Dinotrema borzhomii* Tobias, 2004*
17. *Dinotrema concinnum* (Haliday, 1838)
18. *Dinotrema contracticorne* (Fischer, 1974)
19. *Dinotrema cratocera* (Thomson, 1895)
20. *Dinotrema dimorpha* (Fischer, 1976)*
21. *Dinotrema intermissum* (Fischer, 1974)
22. *Dinotrema perlustrandum* (Fischer, 1973)*
23. *Dinotrema significarium* (Fischer, 1973)
24. *Dinotrema sinecarinum* (Fischer, 1993)*
25. *Dinotrema tauricum* (Telenga, 1935)*
26. *Idiasta (Idiasta) picticornis* (Ruthe, 1854)
27. *Orthostigma beyarslani* Fischer, 1995
28. *Orthostigma laticeps* (Thomson, 1895)
29. *Orthostigma longicorne* Königsmann, 1969*
30. *Orthostigma maculipes* (Haliday, 1838)
31. *Phaenocarpa (Phaenocarpa) bicolor* (Foerster, 1863)*
32. *Phaenocarpa (Phaenocarpa) brevipalpis* (Thomson, 1895)
33. *Phaenocarpa (Phaenocarpa) carinthiaca* Fischer, 1975*

34. *Phaenocarpa (Phaenocarpa) ruficeps* (Nees, 1812)
35. *Pseudopezomachus cursitans* (Ferrière, 1930)
36. *Pseudopezomachus masii* Nixon, 1940
37. *Synaldis concolor* (Nees, 1812)
38. *Synaldis distracta* (Nees, 1834)
39. *Synaldis maxima* Fischer, 1972*
40. *Synaldis megastigma* Fischer, 1967
41. *Synaldis ultima* Fischer, 1970
- Tribe Dacnusini Foerster, 1863**
42. *Aristelix persica* Peris-Felipo, 2015
43. *Chorebus (Chorebus) affinis* (Nees, 1812)
44. *Chorebus (Stiphrocera) anasellus* (Stelfox, 1951)
45. *Chorebus (Stiphrocera) aphantus* (Marshall, 1895)
46. *Chorebus (Phaenolexis) ares* (Nixon, 1944)
47. *Chorebus (Stiphrocera) asphodeli* Griffiths, 1968
48. *Chorebus (Chorebus) axillaris* Fischer, Lashkari Bod, Rakhshani & Talebi, 2011
49. *Chorebus (Stiphrocera) baeticus* Griffiths, 1967
50. *Chorebus (Phaenolexis) bathyzonus* (Marshall, 1895)
51. *Chorebus (Phaenolexis) caesariatus* Griffiths, 1967
52. *Chorebus (Phaenolexis) compressiventris* (Telenga, 1935)
53. *Chorebus (Stiphrocera) cubocephalus* (Telenga, 1935)
54. *Chorebus (Stiphrocera) diremtus* (Nees, 1834)
55. *Chorebus (Phaenolexis) femoratus* (Tobias, 1962)
56. *Chorebus (Stiphrocera) flavipes* (Goureau, 1851)
57. *Chorebus (Phaenolexis) fuscipennis* (Nixon, 1937)
58. *Chorebus (Phaenolexis) gedanensis* (Ratzeburg, 1852)
59. *Chorebus (Chorebus) gracilipes* (Thomson, 1895)
60. *Chorebus (Chorebus) groschkei* Griffiths, 1967
61. *Chorebus (Phaenolexis) heringianus* Griffiths, 1967
62. *Chorebus (Phaenolexis) iridis* Griffiths, 1968
63. *Chorebus (Stiphrocera) lar* (Morley, 1924)
64. *Chorebus (Phaenolexis) leptogaster* (Haliday, 1839)
65. *Chorebus (Chorebus) longiarticulis* Fischer, Lashkari Bod, Rakhshani & Talebi, 2011
66. *Chorebus (Stiphrocera) melanophytobiae* Griffiths, 1968*
67. *Chorebus (Stiphrocera) merellus* (Nixon, 1937)
68. *Chorebus (Stiphrocera) misellus* (Marshall, 1895)
69. *Chorebus (Chorebus) mucronatus* (Telenga, 1935)
70. *Chorebus (Chorebus) nigridiremptus* Fischer, Lashkari Bod, Rakhshani & Talebi, 2011
71. *Chorebus (Chorebus) nigriscaposus* (Nixon, 1949)
72. *Chorebus (Chorebus) nixoni* Burghelle, 1959
73. *Chorebus (Phaenolexis) ornatus* (Telenga, 1935)
74. *Chorebus (Chorebus) parvungula* (Thomson, 1895)
75. *Chorebus (Phaenolexis) posticus* (Haliday, 1839)
76. *Chorebus (Chorebus) properesam* Fischer, Lashkari Bod, Rakhshani & Talebi, 2011
77. *Chorebus (Stiphrocera) pseudomisellus* Griffiths, 1968
78. *Chorebus (Chorebus) ruficollis* (Stelfox, 1957)
79. *Chorebus (Chorebus) scabiosae* Griffiths, 1967
80. *Chorebus (Chorebus) solstitialis* (Stelfox, 1951)
81. *Chorebus (Stiphrocera) spenceri* Griffiths, 1964
82. *Chorebus (Chorebus) stilifer* Griffiths, 1968
83. *Chorebus (Phaenolexis) tamisi* (Nixon, 1944)
84. *Chorebus (Chorebus) thusa* (Nixon, 1937)*
85. *Chorebus (Stiphrocera) tumidus* (Tobias, 1966)

86. *Chorebus (Chorebus) uliginosus* (Haliday, 1839)
87. *Chorebus (Stiphrocera) venustus* (Tobias, 1962)
88. *Chorebus (Chorebus) zarghanensis* Fischer, Lashkari Bod, Rakhshani & Talebi, 2011
89. *Coelinidea elegans* (Curtis, 1829)
90. *Coelinidea gracilis* (Curtis, 1829)
91. *Coloneura dice* (Nixon, 1943)
92. *Dacnusa (Aphanta) hospita* (Foerster, 1863)
93. *Dacnusa (Aphanta) sasakawai* Takada, 1977
94. *Dacnusa (Dacnusa) areolaris* (Nees, 1811)*
95. *Dacnusa (Dacnusa) confinis* Ruthe, 1859
96. *Dacnusa (Dacnusa) gentianae* Griffiths, 1967
97. *Dacnusa (Dacnusa) pubescens* (Curtis, 1826)
98. *Dacnusa (Pachysema) abdita* (Haliday, 1839)
99. *Dacnusa (Pachysema) alpestris* Griffiths, 1967
100. *Dacnusa (Pachysema) aquilegiae* Marshall, 1896*
101. *Dacnusa (Pachysema) aterrima* Thomson, 1895
102. *Dacnusa (Pachysema) clematidis* Griffiths, 1967
103. *Dacnusa (Pachysema) evadne* Nixon, 1937
104. *Dacnusa (Pachysema) metula* (Nixon, 1954)*
105. *Dacnusa (Pachysema) monticola* (Foerster, 1863)
106. *Dacnusa (Pachysema) sibirica* Telenga, 1935
107. *Protodacnusa aridula* (Thomson, 1895)
108. *Protodacnusa litoralis* Griffiths, 1964

Discussion

In the present study, a total of 108 Alysiinae species are catalogued from Iran (41 species belonging to Alysiini and 67 belonging to Dacnusini). Fifteen species are recorded for the first time: *Idiasta picticornis* (Ruthe, 1854), *Orthostigma latinerve* (Petersen, 1956), *Chorebus anasellus* (Stelfox, 1951), *Chorebus melanophytobiae* Griffiths, 1968, *Chorebus*

thusa (Nixon, 1937), *Dacnusa areolaris* (Nees, 1811), *Dacnusa aquilegiae* Marshall, 1896, *Dacnusa metula* (Nixon, 1954), *Dinotrema amparoae* Peris-Felipo, 2013, *Dinotrema borzhomii* Tobias, 2004, *Dinotrema dimorpha* (Fischer, 1976), *Dinotrema perlustrandum* (Fischer, 1973), *Dinotrema sinecarinum* (Fischer, 1993), *Dinotrema tauricum* (Telenga, 1935), *Orthostigma longorne* Königsmann, 1969, *Phaenocarpa bicolor* (Foerster, 1863), *Phaenocarpa carinthiaca* Fischer, 1975, and *Synaldis maxima* Fischer, 1972.

This number of species is higher than the number in adjacent countries such as Azerbaijan (107), Turkey (36), Armenia (26), Turkmenistan (10), Afghanistan (10), and Pakistan (1) (Yu *et al.* 2012). Comparison of the distribution of the new species recorded for Iran shows that Hormozgan with nine records was the province with the largest increase followed by Kerman and Fars with three (Fig. 1). Further investigations both on the fauna composition and the host associations of the Iranian Alysiinae are necessary to provide a basis for biological control of the dipterous pests in agricultural and urban ecosystems.

Acknowledgments

We would like to thank the Department of Entomology, Tarbiat Modares University for funding this research. Contribution by Ehsan Rakhshani was supported by the grant No. 89-9198, University of Zabol

References

- Belokobylskij, S.A. 2005. On the systematic and distribution of three rare alysiine genera (Hymenoptera: Braconidae: Alysiinae). *Genus*, 16(3): 431–444.
- Farahani, S., Talebi, A.A. and Rakhshani, E. 2016. Iranian Braconidae (Insecta: Hymenoptera: Ichneumonoidea): diversity, distribution and host association. *Journal of Insect Biodiversity and Systematics*, 2(1): 1–92.
- Fischer, M. 1990. Westpaläarktische *Phaenocarpa*-Arten: Vorläufiger Bestimmungsschlüssel,

- Deskriptionen und Redeskriptionen (Hymenoptera, Braconidae, Alysiinae). *Annalen des Naturhistorisches Museum Wien*, 91B: 105–135.
- Fischer, M. 1993. Zur Formenvielfalt der Kieferwespen der Alten Welt: Über die Gattungen *Synaldis* Förster, *Trisynaldis* Fischer und *Kritscheryia* Fischer gen. nov. (Hymenoptera, Braconidae, Alysiinae). *Annalen des Naturhistorisches Museum Wien*, 94/95 B: 451–490.
- Fischer, M. and Beyarslan, A. 2012. New species of *Synaldis* Förster and *Idiasta* Förster, and further records of Turkish Alysiini (Hymenoptera: Braconidae: Alysiinae). *Zoology in the Middle East*, 55: 55–64. doi: 10.1080/09397140.2012.10648918
- Gadallah, N.S., Ghahari, H., Peris-Felipo, F.J. and Fischer, M. 2015. An annotated catalogue of the Iranian Alysiinae (Hymenoptera: Braconidae). *Zootaxa*, 3974(1): 1–28. doi: 10.11646/zootaxa.3974.1.1c
- Peris-Felipo, F.J., Belokobylskij, S.A. and Jiménez-Peydró, R. 2014. Revision of the western Palaearctic species of the genus *Dinotrema* Förster, 1862 (Hymenoptera, Braconidae, Alysiinae). *Zootaxa*, 3885(1): 1–483. doi: 10.11646/zootaxa.3885.1.1
- Peris-Felipo, F.J., Ameri, A., Rakhshani, E. and Belokobylskij, S.A. 2016. The genus *Aspilota* Foerster (Hymenoptera: Braconidae: Alysiinae) in western Asia. *Journal of Insect Biodiversity and Systematics*, 2(2): 259–283.
- Sharkey, M.J. and Wharton, R.A. 1997. Morphology and terminology. In: Wharton, R.A., Marsh, P.M., Sharkey, M.J. (eds) *Manual of the New World Genera of the Family Braconidae (Hymenoptera)*. International Society Hymenopterists, Washington DC: pp. 19–37.
- Tobias, V.I. 1986. *Keys to the insects of the European part of the USSR*, Vol. III. Hymenoptera. Part V. Nauka, Leningrad, 231 pp.
- van Achterberg, C. 1993. Illustrated key to the subfamilies of the Braconidae (Hymenoptera: Ichneumonoidea). *Zoologische Verhandelingen*, Leiden 283: 1–189.
- Yari, Z., Cortés, E., Peris-Felipo, F.J. and Rakhshani, E. 2016. A faunistic survey on the genus *Chorebus* Haliday (Hymenoptera, Braconidae, Alysiinae, Dacnusini) in Eastern Iran. *Journal of Insect Biodiversity and Systematics*, 2(3): 355–366.
- Yu, D.S.K., van Achterberg, C. and Horstmann, K., 2012. Taxapad 2012, Ichneumonoidea 2011. Database on flash-drive, Ottawa, Ontario, Canada.

زیرخانواده گزارش‌های جدید و فهرست به روز شده گونه‌ها

النا کورتس^۱، علی عامری^۲، علی اصغر طالبی^۲، احسان رخشانی^۳ و فرانسیسکو پریس فلیپو^{۴*}

۱ فرناندو کاتولیکو ۷، آلیکانت، اسپانیا.

۲ گروه حشره‌شناسی، دانشکده کشاورزی، دانشگاه تربیت مدرس، تهران، ایران.

۳ گروه گیاه‌پژوهشکی، دانشکده کشاورزی، دانشگاه زابل، ایران.

۴ بلیچسترنس ۱۵، ch-۴۰۵۸، بازل، سوئیس.

* پست الکترونیکی نویسنده مسئول مکاتبه: peris.felipo@gmail.com

تاریخ دریافت: ۱۲ آبان ۱۳۹۵، تاریخ پذیرش: ۲۲ آذر ۱۳۹۵، تاریخ انتشار: ۱۴ آذر ۱۳۹۵

چکیده: زیرخانواده آلیزینه یکی از گروه‌های متنوع و ناشناخته از زنبورهای پارازیتویید داخلی متعلق به خانواده برآکونیده (Braconidae) محسوب می‌شود. نمونه‌برداری در مناطق مختلف ایران عمدها در نواحی جنوبی، جنوب شرق و مرکز انجام شد. پانزده گونه برای اولین بار از ایران گزارش شدند. این گونه‌های متعلق به هر دو قبیله Alysiini؛ *D. borzhomii* Tobias, 2004; *Dinotrema amparoae* Peris-Felipo, 2013; *D. perlustrandum* (Fischer, 1973); *D. dimorpha* (Fischer, 1976); *orthostigma*; *D. tauricum* (Telenga, 1935); *D. sinecarinum* (Fischer, 1993); *Phaenocarpa bicolor* (Foerster, 1863); *longicornis* Konigsman, 1969 و *Synaldis maxima* Fischer, 1972 و *P. carinthiaca* Fischer, 1975 و *C. thusa* (Nixon, 1937); *Chorebus melanophytobiae* Griffiths, 1968) Dacnusini و *D. aquileiae* Marshall, 1896; *Dacnusa areolaris* (Nees, 1811) ۱۹۳۷ و *D. metula* (Nixon, 1954) ایران بودند. فهرست کل گونه‌های زیرخانواده Alysiinae ایران مشتمل بر ۱۰۸ گونه نیز ارایه شده است.

واژگان کلیدی: Alysiinae، لیست به روز شده، گزارش جدید، ایران.