



Parasitoids (Hymenoptera, Braconidae, Aphidiinae) of *Aphis gossypii* (Hemi., Aphididae) in Bengaluru, India

Dayanandam Lokeshwari^{1*}, Nallur Kothandaraman Krishna Kumar² and Ehsan Rakhshani³

¹ Division of Entomology and Nematology, Indian Institute of Horticultural Research, Hessaraghatta Lake Post, Bengaluru-560 089, India.

² Division of Horticultural Science, Indian Council of Agriculture Research, Krishi Anusandhan Bhawan-II, New Delhi-110 012, India.

³ Department of plant protection, College of Agriculture, University of Zabol, I.R. Iran

Received:
13 March 2016

Accepted:
02 June 2016

Published:
06 June 2016

Subject Editor:
Vladimir Žikić

ABSTRACT. A survey was conducted in fields of cucumber and watermelon in Bengaluru, India during April, 2014 for determination of the common parasitoids (Hym.: Braconidae, Aphidiinae) attacking melon aphid, *Aphis gossypii* Glover (Hemi.: Aphididae). Three species of Aphidiinae, including *Aphidius platensis* Brethes, 1913; *Binodoxys acalephae* (Marshall, 1896) and *Lipolexis oregmae* (Gahan, 1931) were collected and identified, from which *Aphidius platensis* is newly recorded for India. The early evidence of *Aphidius colemani* Viereck, 1912 in India can also be contributed to *A. platensis*. Diagnostic characters of three species were compared with closely related species in brief.

Key words: *Aphidius platensis*, *Lipolexis oregmae*, *Binodoxys acalephae*, New record, biological control

Citation: Lokeshwari, D., Kumar, N.K. and Rakhshani, E. 2015. Parasitoids (Hymenoptera, Braconidae, Aphidiinae) of *Aphis gossypii* (Hemi., Aphididae) in Bengaluru, India. *Journal of Insect Biodiversity and Systematics*, 1 (2): 155–163

Introduction

The melon aphid, *Aphis gossypii* Glover (or cotton aphid) is highly polyphagous species, marked as a pest of various cultivated plants from the families Cucurbitaceae, Rutaceae and Malvaceae (Blackman and Eastop 2000). This aphid is a serious pest of cucurbits in temperate zones principally in fields and green houses (Leclant and Deguine 1994). The aphids feed on the underside of the leaves sucking nutrients from the host plant. When they are in dense populations, aphids cause the leaves to

become chlorotic, curled and distorted, hindering photosynthetic capacity of the plant (Shannag *et al.* 1998). A further problem for plant is excretion of honeydew by aphids, which provides a substrate for growth of sooty mold directly reducing the photosynthesis and eventually the quality of the fruits. The melon aphid effectively transmits important plant pathogenic viruses such as cucumber mosaic virus (CMV) and watermelon mosaic virus (WMV) (Castle *et al.* 1992).

Corresponding author: Dayanandam Lokeshwari, E-mail: lokibiotech@gmail.com

Copyright © 2015, Lokeshwari et al. This is an open access article distributed under the terms of the Creative Commons Attribution License (CC BY NC 4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

The control of aphid pests depends mainly on the frequent application of synthetic insecticides, which often results in many negative effects, such as development of insecticide resistance (Herron *et al.* 2001) and other associated environmental problems. The application of insecticides leads to the outbreaks of the melon aphid by destruction of natural enemies (Pimentel *et al.* 1992). Furthermore, resistance by melon aphid to all the usual organophosphate, carbamate and synthetic pyrethroid aphicides is also widely documented (Herron *et al.* 2001).

Many insects as natural enemies may be very effective against melon aphid, including ladybirds (Coleoptera: Coccinellidae), hoverflies (Diptera: Syrphidae) and braconid wasps (Hymenoptera: Braconidae). Under the influence of natural enemies the aphid populations increase only by fraction of their potential rates, in field experiments (Frazer 1988; Starý 1988). Aphid parasitoids such as Aphidiinae have been considered as useful agents for biocontrol of *A. gossypii*, both in greenhouses and in the fields (van Schelt *et al.* 1990; Vásquez *et al.* 2006). For successful biological control of aphids, it is necessary to search for effective natural enemies, particularly those that can be used to control aphids on crops (Rabasse and van Steenis 1987; Burgio *et al.* 1997).

The subfamily Aphidiinae is a group of solitary endoparasitoids that attack only aphids, playing a significant role in the structure of their populations (Starý 1970, 1988). Taxonomy and the diversity of plant-host-parasitoid associations concerning aphidiines in India have been investigated by several authors and the data are summarized within a recently published catalogue by Akhtar *et al.* (2011), since then, supplemented with few other contributions (e.g. Rakhshani *et al.* 2012).

Very few species of Aphidiinae have been recorded from Karnataka province (India), including *Aphidius absinthii*

(Marshall (Starý and Ghosh 1983), *Aphidius colemani* Viereck (Starý and Ghosh 1983), *Binodoxys acalephae* (Marshall) (Dharmadhikari and Ramaseshiah 1970), *Binodoxys brevicornis* (Haliday) (Starý and Ghosh 1983), *Binodoxys indicus* (Subba Rao and Sharma) (Ramaseshiah 1972), *Diaeretiella rapae* (M'Intosh) (Sethumadhavan and Dharmadhikari 1969), *Lipolexis oregmae* (Gahan) (Dharmadhikari and Ramaseshiah 1970), *Lysiphlebus testaceipes* (Cresson) (Ramaseshiah *et al.* 1968), *Praon abjectum* Haliday (Dharmadhikari and Ramaseshiah 1970), *Praon volucre* (Haliday) (Dharmadhikari and Ramaseshiah 1970) and *Toxares shigai* Takada (Dharmadhikari and Ramaseshiah 1970). Among the recorded species, only *Aphidius colemani*, *Binodoxys indicus*, *Diaeretiella rapae*, *Lipolexis oregmae* and *Praon volucre* have been recorded in the association with the melon aphid on various host plants (Akhtar *et al.* 2011). The objective of this research was to provide the background systematics information on the Aphidiinae parasitoids of melon aphid, *Aphis gossypii* in Bengaluru, India.

Material and methods

The survey was conducted in fields of cucumber and watermelon cultivated in the experimental plot maintained at the Indian Institute of Horticultural Research (IIHR), Bengaluru, India during April 2014. Bengaluru, known as Bangalore is situated in the southeast of the South Indian state of Karnataka, positioned at 12.97° N 77.56° E covering an area of 2.190 square kilometers. It has a tropical wet and dry savanna climate with a pronounced dry season in the low-sun months (Kottek *et al.* 2006). The district is endowed with various agricultural and horticultural crops including cotton, pumpkins, rice, groundnut, peppers, sugarcane, grapes and mulberry. Leaf samples infested with the colonies of melon aphids, were collected from field and carefully places into semi-transparent rearing

plastic boxes, immediately covered by mesh for ventilation. The reared material was transferred to the laboratory, where it was subsequently maintained for 2–3 weeks until parasitoids' emergence. Once they were detected, they were carefully picked up using an aspirator and dropped into the 75% ethanol kept for identification. The external morphology of parasitoids was studied using NIKON SMZ645 stereomicroscope and a NIKON Eclipse E200 microscope. Terminology for the selected morphological characters follows Sharkey and Wharton (1997). Distributional data of the recorded species followed Yu *et al.* (2012). The specimens were deposited at the Insect Collection of Zabol University, Zabol, Iran.

Results

As a result of our investigation of parasitoids of melon aphids three Aphidiinae species belonging to two genera listed below.


Aphidius platensis Brethes, 1913 (Figs. 1–3)

Material examined: 4♀♀2♂♂, reared from *A. gossypii*, on *Cucurbita pepo* L., April, 2014, IIHR, Bengaluru; 2♀♀, reared from *A. gossypii*, on *Gossypium hirsutum* L., April 2014, IIHR, Bengaluru, Leg. D. Lokeshwari.

Diagnosis: *Aphidius platensis* is closely related to *A. colemani* Viereck on the basis of wing venations (Fig. 1) and shape of ovipositor sheath (Fig. 3), from which it can be immediately differentiated by having a shorter R1 vein (0.77–0.89X as long as stigma in *A. platensis*, vis1.0–1.1X as long as stigma in *A. colemani*). Anterolateral area of petiole in *A. platensis* bearing 4–5 sharply prominent costae (Fig. 2), while in *A. colemani* 2–3 bluntly shallow costae occur.

Distribution in India: New record for India.

General distribution: Neotropical, Central Asia (Iran), Oriental (New record).



Figures 1–3. Diagnostic characters of *Aphidius platensis* Brethes, **1.** Forewing, **2.** Lateral aspect of petiole, **3.** ovipositor sheath.

***Binodoxys acalephae* (Marshall, 1896)**
(Figs. 4–6)


Material examined: 2♀♀ 2♂♂, reared from *A. gossypii* Glover, on, *Gossypium hirsutum* L., April 2014, IIHR, Bengaluru, Leg. D. Lokeshwari.

Diagnosis: *Binodoxys acalephae* is closely related to *B. angelicae* (Haliday) and *B. indicus* (Subba Rao and Sharma) from which it can be readily distinguish by the characters of the wing venation (Fig. 4), and petiole (Fig. 5). *Binodoxys acalephae* has

shorter R1 vein (0.42–0.48X as long as stigma in *B. acalephae*, vis 0.65–0.68X and 0.64–0.66X as long as stigma in *B. angelicae* and *B. indicus*, respectively). In petiole of *B. acalephae* the distance between the primary and secondary tubercles at spiracles is smaller than its width; while in two other species it is clearly larger.

Distribution in India: Jammu and Kashmir, Karnataka, Manipur (Akhtar *et al.* 2011).

General distribution: Holarctic.



Figures 4–6. Diagnostic characters of *Binodoxys acalephae* (Marshall), 4. Forewing, 5. Lateral aspect of petiol, 6. ovipositor sheath.

***Lipolexis oregmae* (Gahan, 1931) (Figs. 7–9)**

Material examined: 3♀♀ 2♂♂, reared from *A. gossypii* Glover on *Capsicum annuum* L., April 2014, IIHR, Bengaluru, Leg. D. Lokeshwari.


Diagnosis: *Lipolexis oregmae* is closely related to *L. gracilis* from which can be

separated by a series of characters pertaining to the dorsal surface of petiole and female genitalia. In *L. oregmae*, the petiole is smooth dorsally, bearing crenulated longitudinal carinae laterally (Fig. 8); while petiole in *L. gracilis* has a prominent central carina bifurcated near the middle. The ovipositor sheaths in *L.*

oregmae are short and slightly curved downwards (Fig. 9), in contrast to *L. gracilis* where those structure are long, and strongly curved downwards. In the forewing of *L. oregmae* the ratio of R1/stigma is 1.0–1.1 (Fig. 7), but 1.25–1.3 in *L. gracilis*.

Distribution in India: Widely distributed in India, Jammu and Kashmir, Karnataka, Manipur, Meghalaya, Sikkim, Tripura, Uttar Pradesh and West Bengal (Akhtar *et al.* 2011).

General distribution: Oriental, introduced to Central America.



Figures 7-9. Diagnostic characters of *Lipolexis oregmae* (Gahan), 7. Forewing, 8. Lateral aspect of petiol, 9. ovipositor sheath

Discussion

The association of two common, but very efficient parasitoids on melon aphid in Bengaluru brings potentially powerful biological control agents. *Lipolexis oregmae* has already successfully been used as biological control agent against *Toxoptera citricia* Kirkaldy (Persad *et al.* 2007). The second parasitoid, *Aphidius platensis* was classified as a native species in South America (Starý 1970). Later, its origin was again considered revealing Indian source,

which was explained by a series of accidental introductions to other continents (Starý 1972). *Aphidius platensis* was synonymized by *Aphidius colemani* Viereck which also showed the same host range and wide distribution (Starý 1975), therefore it was subsequently studied under the name of *A. colemani*. True *Aphidius colemani* has been frequently evaluated as the most efficient parasitoid of *A. gossypii* in greenhouses (van Steenis 1995). This parasitoid was widely used in the biological control of *A. gossypii* (van

Schelt *et al.* 1990). Existence of morphological differences and the discrete pattern of distribution for *A. colemani* have later been criticized (Takada 1998; Barahoei *et al.* 2013) and finally clarified by Tomanović *et al.* (2014). Based on this study, *A. platensis* has been re-validated, sharing the same host range pattern and the same origin as *A. colemani*. Furthermore, all the material from the central Asia (Iran) (Rakhshani *et al.* 2008; Zamani *et al.* 2007) is classified as *A. platensis* with no trace of true *A. colemani*. Consequently, the early evidence of *A. colemani* in India (Akhtar *et al.* 2011) may contribute to *A. platensis*. The occurrence of both species in the Oriental and Eastern Palaearctic regions need to be investigated in future works.

Lipolexis oregmae has a wide distribution in the Eastern Palaearctic and the Oriental including Guam (Miller *et al.* 2002), India (Starý and Bhagat 1978; Starý and Ghosh 1983); the Philippines, Hong Kong and Taiwan (Starý and Schlinger 1967; Chou 1981); Korea (Starý and Choi 2000) and Vietnam (Starý and Zelený 1983). Miller *et al.* (2002) re-examined the paratypes of *L. oregmae* and compared the petiole with that of *L. scutellaris*. As a result, the two species appear to be identical, and *L. scutellaris* was classified as a junior synonym of *L. oregmae*. Two other species, *L. pseudoscutellaris* Pramanik & Raychaudhuri and *L. myzackaiae* Pramanik & Raychaudhuri were also described from India (Pramanik and Raychaudhuri 1984). The original description of the first species leads to *L. oregmae*, and it was possibly classified as a new species because the authors may have overlooked drawings in the re-description of *L. oregmae* by Starý (1960). Also, in the case of *L. myzackaiae* there are some discrepancies between the description and the drawing of the petiole. Although (Pramanik and Raychaudhuri 1984) based the description of this species noting that the petiole laterally possess longitudinal carinae, they

were not shown in the attached figures. While we did not revise the type material, both species can be considered as conspecific with *L. oregmae*. In China, both *L. gracilis* (syn., *Lipolexis chinensis* Chen) and *L. oregmae* are present (Chen and Shi 2001). The third species, *L. wuyiensis* Chen, parasitoid of *Oregmala nigera* (Zehnt.) described as a distinct species (Chen, 1981). However, in our opinion, it appears to be another synonym of *L. oregmae* revealing that it parasitizes the same hosts as true *L. oregmae* in its original area in the Philippines (Miller *et al.* 2002).

Further investigations including DNA analyses become necessary in order to reveal the entire parasitoid complex of *A. gossypii* in Bengaluru; especially the other host plants as refugia for this aphid. Collecting the additional material of *A. platensis* or the examination of previously recorded material from India is also necessary to confirm the coincidence of *A. colemani*.

Acknowledgments

The authors thank Director, Indian Institute of Horticultural Research, Bengaluru for encouragement. Thanks are also due to Indian Council of Agricultural Research (ICAR), New Delhi for financial support through the Out Reach Programme on Management of Sucking Pests of Horticultural Crops. The contribution by E. Rakhshani has supported by the grant No. 89-9198, University of Zabol.

References

- Akhtar, M.S., Dey, D. and Usmaniz, M.K. 2011. A catalogue of aphid parasitoids (Hymenoptera: Braconidae: Aphidiinae) from India. *Insecta Mundi*, 151: 1–31.
- Barahoei, H., Rakhshani, E., Madjdzadeh, S.M., Alipour, A., Taheri, S., Nader, E., Mitrovski-Bogdanović, A., Petrović-Obradović, O., Starý, P., Kavallieratos, N.G. and Tomanović, Ž. 2013. Aphid parasitoid species (Hymenoptera: Braconidae: Aphidiinae) of central

- submountains of Iran. *North-Western Journal of Zoology*, 9(1): 70–93.
- Blackman, R.L. and Eastop, V.F. 2000. *Aphids on the World's Crops: An Identification and Information Guide*, 2nd edn. Chichester, Wiley. 466pp.
- Burgio, G., Ferrari, R. and Nicoli, G. 1997. Biological and Integrated Control of *Aphis gossypii* Glover (Horn., Aphididae) in protected cucumber and melon. *Bollettino dell'Istituto di Entomologia "Guido Grandi" della Università degli Studi di Bologna*, 51: 171–178.
- Castle, S.J., Perring, T.M., Farrar, C.A. and Kishaba, A.N. 1992. Field and laboratory transmission of watermelon mosaic virus 2 and zucchini yellow mosaic virus by various aphid species. *Phytopathology*, 82(2): 235–240.
- Chen, J.H. 1981. [Description of a new species on aphid parasites of *Oregma lanigera* (Zehntner) (Hym.: Aphidiidae)]. *Journal of Fujian Agricultural University*, 10(1): 33–38. [in Chinese]
- Chen, J. and Shi, Q. 2001. *Systematic studies on Aphidiidae of China (Hymenoptera: Aphidiidae)*. Fujian Science and Technology Press, Fuzhou, 273 pp. [in Chinese]
- Chou, L.Y. 1981. The genera of Aphidiidae (Hymenoptera: Ichneumonidea) in Taiwan. *Journal of Agricultural Research, China*, 30: 308–323.
- Dharmadhikari, P.R. and Ramaseshiah, G. 1970. Recent records of aphidiids (Hymenoptera: Aphidiidae) in India. *Commonwealth Institute of Biological Control Technical Bulletin*, 13: 83–89.
- Frazer, B.D. 1988. Predators. pp: 217–230, In: Minks, A.K. and Harrewijn, P. (Eds.), *Aphids: Their Biology, Natural Enemies and Control*. Volume B. Elsevier, Amsterdam.
- Herron, G.A., Powisand, K. and Rophail, J. 2001. Insecticide resistance in *Aphis gossypii* Glover (Hemiptera: Aphididae), a serious threat to Australia cotton. *Australian Journal of Entomology*, 40: 85–91.
- Kottek, M., Grieser, J., Beck, C., Rudolf, B. and Rubel, F. 2006. World map of the Köppen-Geiger climate classification updated. *Meteorologische Zeitschrift*, 15(3): 259–263. DOI: <http://dx.doi.org/10.1127/09412948/2006/0130>
- Leclant, F., Deguine, J. P. 1994. Aphids (Hemiptera: Aphididae). pp: 285–323, In: Matthew, G. A., Tunstall, J. P. (Eds) *Insect Pests of Cotton*. Wallingford UK, CAB International.
- Miller, R.H., Pike, K.S. and Starý, P. 2002. Aphid parasitoids (Hymenoptera: Aphidiidae) on Guam. *Micronesica*, 34: 83–107.
- Persad, A.B., Hoy, M.A. and Nguyen, R. 2007. Establishment of *Lipolexis oregmae* (Hymenoptera: Aphidiidae) in a classical biological control program directed against the brown citrus aphid (Homoptera: Aphididae) in Florida. *Florida Entomologist*, 90: 204–213. DOI: <http://dx.doi.org/10.1653/0015-4040>
- Pimentel, D., Acquay, H., Biltonen, M., Rice, P., Silva, M., Nelson, J., Liper, V., Giordano, S., Harowitz, A. and D'amore, M. 1992. Environmental and economic costs of pesticide use. *BioScience*, 42(10): 750–760.
- Pramanik, D.R. and Raychaudhuri, D. 1984. New aphid-parasitoids (Hymenoptera: Aphidiidae) from Negaland, Northeast India. *Akitu*, 61: 1–4.
- Rabasse, J.M. and van Steenis, M.J. 1987. Biological control of aphids, pp: 235–243, In: Albajes, R. (Ed.), *Integrated Pest and Disease Management in Greenhouse Crops*. Kluwers Academic Publishers, Netherlands.
- Rakhshani, E., Barahoei, H., Ahmed, Z., Tomanović, Z., Janković, M., Petrović, A. and Starý, P. 2012. New species and additional evidence of aphid parasitoids (Hymenoptera, Braconidae, Aphidiinae) from India. *Zootaxa*, 3397: 45–54.
- Rakhshani, E., Talebi, A.A., Starý, P., Tomanović, Z., Kavallieratos, N.G. and Manzari, S. 2008. A review of *Aphidius* Nees (Hymenoptera: Braconidae: Aphidiinae) in Iran: host associations, distribution and taxonomic notes. *Zootaxa*, 1767: 37–54.
- Ramaseshiah, G. 1972. Supply of *Binodoxys indicus* (for the U.S.A.). Report of the Commonwealth Institute of Biological Control Technical Bulletin, 1971: 67.
- Ramaseshiah, G., Bhat, K. V. and Dharmadhikari, P. R. 1968. Influence of host aphid, host plant and temperature on the laboratory breeding of *Lysiphlebus testaceipes*. *Indian Journal of Entomology*, 30: 281–285

- Sethumadhavan, T.V. and Dharmadhikari, P. R. 1969. Notes on *Diaeretiella rapae* (M'Intosh) (Hymenoptera: Aphidiidae) parasitic on aphids on crucifers in India. *Commonwealth Institute of Biological Control Technical Bulletin*, 11: 173-177.
- Shannag, H.K., Thorvilson, H. and El-Shatnawi, M.D.K. 1998. Changes in photosynthetic and transpiration rates of cotton leaves infested with the cotton aphid, *Aphis gossypii*: unrestricted infestation. *Annals of Applied Biology*, 132(1): 13-18.
- Sharkey, M.J. and Wharton, R.A. 1997. Morphology and terminology, pp: 19-37, In: Wharton, R.A., Marsh, P.M. and Sharkey, M.J. (Eds.), *Manual of the New World Genera of the Family Braconidae (Hymenoptera)*. Special Publication 1. Washington, DC, International Society of Hymenopterists.
- Starý, P. 1960. The systematic position of *Diaeretus oregmae* Gahan (Hymenoptera: Aphidiidae). *Insecta Matsumurana*, 23: 109-111.
- Starý, P. 1970. *Biology of Aphid Parasites (Hymenoptera: Aphidiidae) with Respect to Integrated Control*. Dr. W. Junk, b.v., The Hague. 643pp.
- Starý, P. 1972. *Aphidius platensis* Brethes, its distribution and host range (Hym.: Aphidiidae). *Oriental Insects*, 6(3): 359-370.
- Starý, P. 1975. *Aphidius colemani* Viereck: its taxonomy, distribution and host range (Hymenoptera: Aphidiidae). *Acta Entomologica Bohemoslovaca*, 72: 156-163.
- Starý, P. 1988. Aphidiidae, pp: 171-184, In: Minks, A.K. and Harrewijn, P. (Eds.), *Aphids: Their Biology, Natural Enemies and Control*. Volume B. Elsevier, Amsterdam.
- Starý, P. and Bhagat, R.C. 1978. Records and descriptions of aphid parasitoids from Kashmir, India (Hymenoptera: Aphidiidae). *Acta Entomologica Bohemoslovaca*, 75: 387-393.
- Starý, P. and Choi, J.Y. 2000. An annotated review of tritrophic associations of aphid parasitoids (Hymenoptera, Braconidae, Aphidiinae) in Korea. *Insecta Koreana*, 17: 91-109.
- Starý, P. and Schlinger, E.I. 1967. *A Revision of the Far East Asian Aphidiidae (Hymenoptera)*. Series Entomologica. The Hague, 204 pp.
- Starý, P. and Zelený, J. 1983. Aphid parasitoids from Vietnam (Hymenoptera: Aphidiidae). *Acta Entomologica Bohemoslovaca*, 80: 190-195.
- Starý, P. and Ghosh, A.K. 1983. Aphid Parasitoids of India and Adjacent Countries (Hymenoptera: Aphidiidae). Zoological Survey of India. Technical Monograph No. 7: 1-96.
- Takada, H. 1998. A review of the *Aphidius colemani* (Hymenoptera: Braconidae; Aphidiinae) and closely related species indigenous to Japan. *Applied Entomology and Zoology*, 33: 59-66.
- Tomanović, Ž., Petrović, A., Mitrović, M., Kavallieratos, N.G., Starý, P., Rakhshani, E., Rakhshanipour, M., Popović, A., Shukshuk, A.H. and Ivanović, A. 2014. Molecular and morphological variability within the *Aphidius colemani* group with redescription of *Aphidius platensis* Brethes (Hymenoptera: Braconidae: Aphidiinae). *Bulletin of Entomological Research*, 104(5): 552-565. DOI: <http://dx.doi.org/10.1017/S0007485314000327>
- van Schelt, J., Douma, J.B. and Raveneberg, W.J. 1990. Recent developments in the control of aphids in sweet pepper and cucumber. *Bulletin IOBC/WPRS*, 13(5): 190-193.
- van Steenis, M. 1995. Evaluation of four aphidiine parasitoids for biological control of *Aphis gossypii*. *Entomologia Experimentalis et Applicata*, 75(2): 151-157. DOI: <http://dx.doi.org/10.1111/j.15707458.1995.tb01921.x>
- Vásquez, G.M., Orr, D.B. and Baker, J.R. 2006. Efficacy assessment of *Aphidius colemani* (Hymenoptera: Braconidae) for suppression of *Aphis gossypii* (Homoptera: Aphididae) in greenhouse-grown *Chrysanthemum*. *Journal of Economic Entomology*, 99(4): 1104-1111. DOI: <http://dx.doi.org/10.1093/jee/99.4.1104>
- Yu, D.S., van Achterberg, C. and Horstmann, K. 2012. World Ichneumonoidea. Taxonomy, Biology, Morphology and Distribution. Taxapad (Scientific names for information management) Interactive catalogue on DVD/CDROM. Vancouver, Available from: <http://www.taxapad.com>
- Zamani, A.A., Talebi, A.A., Fathipour, Y. and Baniameri, V. 2007. Effect of temperature on life history of *Aphidius colemani* and *Aphidius matricariae* (Hymenoptera: Braconidae), two parasitoids of *Aphis gossypii* and *Myzus persicae* (Homoptera: Aphididae). *Environmental Entomology*, 36(2): 263-271.

پارازیتوئیدهای شته *Aphis gossypii* Glover (Hymenoptera, Braconidae, Aphidiinae) در بنگلور، هندوستان

دایاناندان لوکشواری^{۱*}، نالور کوتاندارامان کریشنا کومار^۲ و احسان رخشانی^۳

۱ بخش حشره‌شناسی و نماتدشناسی، موسسه تحقیقات باغبانی هند، بنگلور، هندوستان

۲ بخش علوم باغبانی، مرکز تحقیقات کشاورزی هند، دهلی‌نو، هندوستان

۳ گروه گیاهپزشکی، دانشکده کشاورزی، دانشگاه زابل، ایران

* پست الکترونیکی نویسنده مسئول مکاتبه: lokibiotech@gmail.com

تاریخ دریافت: ۲۳ اسفند ۱۳۹۴، تاریخ پذیرش: ۱۳ خرداد ۱۳۹۵، تاریخ انتشار: ۱۷ خرداد ۱۳۹۵

چکیده: این تحقیق در بهار سال ۱۳۹۳ به منظور شناسایی پارازیتوئیدهای (Hym.:) *Aphis gossypii* Glover (Braconidae, Aphidiinae) (Hemi.: Aphididae) در مزارع خیار و هندوانه بنگلور هندوستان انجام گردید. سه گونه از زنبورهای زیرخانواده Aphidiinae شامل *Binodoxys aphidius platensis* Brethes, 1913 و *acalephae* (Marshall, 1896) و *Lipolexis oregmae* (Gahan, 1931) جمع‌آوری و شناسایی شدند که از بین آنها گونه *Aphidius platensis* برای اولین بار از کشور هند گزارش می‌شود. به نظر می‌رسد گزارشات قبلی گونه *Aphidius colemani* Viereck, 1912 در هندوستان مربوط به گونه *A. platensis* باشد. خصوصیات مرفولوژیک این سه گونه در مقایسه با گونه‌های نزدیک به طور مختصر مقایسه شد.

واژگان کلیدی: *Binodoxys acalephae*, *Lipolexis oregmae*, *Aphidius platensis*

گزارش جدید، مهار زیستی