



## The beetles (Coleoptera) associated with the Maamora cork oak forest in the north-west of Morocco

**Hamza Habbaz**

Laboratoire d'Ecologie et Environnement, Hassan II University of Casablanca, Faculty of Sciences Ben M'Sick, BP 7955 - Sidi Othman, Casablanca, Morocco [1]; Centre d'Innovation, de Recherche et de Formation, Agence Nationale des Eaux et Forêts, BP. 763, Rabat - Agdal, Morocco [2].

✉ [hamza.habbaz-etu@etu.univh2c.ma](mailto:hamza.habbaz-etu@etu.univh2c.ma)

<https://orcid.org/0009-0000-7117-1075>

**Noureddin Maatouf**

Centre d'Innovation, de Recherche et de Formation, Agence Nationale des Eaux et Forêts, BP. 763, Rabat - Agdal, Morocco.

✉ [noureddin\\_maatouf@yahoo.ca](mailto:noureddin_maatouf@yahoo.ca)

<https://orcid.org/0000-0002-6288-7378>

**Jean-Pierre Lumaret**

Université Paul Valéry Montpellier III, Route de Mende, 34199 Montpellier Cedex 5, France.

✉ [jean-pierre.lumaret@univ-montp3.fr](mailto:jean-pierre.lumaret@univ-montp3.fr)

<https://orcid.org/0000-0003-2926-3974>

**Latifa Rohi**

Laboratoire d'Ecologie et Environnement, Hassan II University of Casablanca, Faculty of Sciences Ben M'Sick, BP 7955 - Sidi Othman, Casablanca, Morocco.

✉ [rohilatifa@gmail.com](mailto:rohilatifa@gmail.com)

<https://orcid.org/0000-0002-4180-1117>

**ABSTRACT.** This study aims to explore and improve the understanding of the biodiversity of beetles in the Maamora forest, one of the largest cork oak forests in the world. A survey was carried out over two consecutive years (2021 and 2022), using both active and passive traps. A total of 8247 individuals were collected, representing 256 species, belonging to 42 families of Coleoptera order. Most of the beetle species belonged to the families Tenebrionidae (27 species), Carabidae (27 species) and Curculionidae (24 species). Out of the 256 species identified, 216 have been included in this document. We only selected species of high heritage value; 53 were saproxylic, including 6 rare species; 29 were endemic, and 26 species were identified as new records for the Moroccan entomofauna. These results provide the first checklist of beetle fauna in the Maamora forest, which will contribute new information on the distribution of beetles in the cork oak forests of Morocco.

**Keywords:** Biodiversity, inventory, distribution, endemic, saproxylic beetles

**Received:**  
December 20, 2023

**Accepted:**  
September 04, 2024

**Published:**  
January 01, 2025

**Subject Editor:**  
Sabu K Thomas

**Citation:** Habbaz, H., Maatouf, N., Lumaret, J.-P. & Rohi, L. (2025) The beetles (Coleoptera) associated with the Maamora cork oak forest in the north-west of Morocco. *Journal of Insect Biodiversity and Systematics*, 11 (1), 39–84.

## INTRODUCTION

Like the forests of the southern side of the Mediterranean basin, Moroccan forests contain a great variety of habitats rich in biodiversity (QCN, 2021). However, they are highly vulnerable, and face multiple crises characterized by significant biodiversity loss and disruption of forest habitats (Hoekstra et al., 2005). This vulnerability stems from both natural and anthropogenic pressures. The current changes, which are becoming increasingly stringent, are expected to intensify in the coming decades (Ali et al., 2022). To address these growing threats, it is necessary to adopt new management approaches and establish a coherent network of protected natural areas in Mediterranean countries (Galbraith & Stroud, 2022). Since 1996, the Maamora forest has been identified as a Site of Biological and Ecological Interest (SIBE) as part of the study on Protected Areas in Morocco (AEFCS, 1996). Located on the Atlantic coast of Morocco, this forest provides numerous ecological goods and services to society (recreational areas, pastures, timber, acorns, etc.). It is one of the most studied ecosystems in Morocco. Its flora is relatively well studied

**Corresponding author:** Hamza Habbaz, ✉ [hamza.habbaz-etu@etu.univh2c.ma](mailto:hamza.habbaz-etu@etu.univh2c.ma), [hamza96habbaz@gmail.com](mailto:hamza96habbaz@gmail.com)


**Copyright** © 2025, **Authors.** This is an open access article distributed under the terms of the Creative Commons NonCommercial Attribution License (CC BY NC 4.0), which permits Share - copy and redistribute the material in any medium or format, and Adapt - remix, transform, and build upon the material, under the Attribution-NonCommercial terms.

(Sauvage, 1961; Benabid, 2000; Aafi et al., 2005). The vascular flora of the Maamora forest represents 48% of the vascular flora of Moroccan cork oak forests and 9.3% of the overall vascular flora of Morocco (Sauvage, 1961; Benabid, 2000; Aafi, 2007). This forest is also home to diverse fauna, including over 700 arthropod species (Villemant & Fraval, 1993) and 150 bird species (Cherkaoui et al., 2007). However, studies on the entomological fauna are limited (Idrissi, 1982; Mahari, 1992; Villemant & Fraval, 1993; El Alami Idrissi, 2013), and information on the structure of beetle populations and their spatial distribution is relatively incomplete. This means that the forest managers have a significant responsibility to conserve this natural heritage. Over the past decades, human activities have exerted numerous pressures on this forest, leading to a reduction in its surface area. The continuous loss and alteration of the forest habitats are expected to have repercussions for the entomological community. The replacement and/or loss of native insect species often result in the destabilization of the original communities and a reduction in the ecological goods and services they provide (Murphy, 1988).

Knowledge about the conservation status of beetle populations in Morocco is limited. There is no list of rare and threatened species, highlighting the need to survey the Maamora forest entomofauna to help assign conservation priorities by identifying heritage species that require urgent protection measures. To update and complement existing information, a targeted inventory of beetles in the Maamora forest was conducted over two consecutive years (2021 and 2022) using various traps and visual collection techniques. The objectives of this study were to obtain a better understanding of the beetles in the forest and to assess the biological and heritage value of the forest habitats.

## MATERIAL AND METHODS

**Study area.** The Maamora forest is located between the cities of Salé and Kénitra, along the Atlantic coast in north-west Morocco (Fig. 1). It extends eastwards along an area 40 km wide and 70 km long, following a bioclimatic gradient ranging from sub-humid to semi-arid (Belghazi & Mounir, 2016). This forest covers approximately 135,000 hectares, including 71,000 hectares of commercial plantations and 64,461 hectares of cork oak woodlands (Belghazi & Mounir, 2016), representing 17% of the total area of cork oak forests in Morocco and 25% of the area of Atlantic cork oak forests (Aafi, 2007). The topography of the forest is mostly flat, interrupted by a southwest-northeast oriented hydrographic network that divides it into five distinct Cantons named A, B, C, D, and E, respectively, and distributed from the sea level up to 300 metres in elevation (Fig. 1). For the purposes of this study, three sampling sites were chosen within the forest, identified as S1, S2 and S3. Only three of the five Cantons were selected for trap installation: site S1 in Canton A, site S2 in Canton C and site S3 in Canton E. Cantons B and D were not included in the study, as Canton B has similar vegetation and climatic characteristics to Canton C, just as Canton D is similar to Canton E. The climate is of the Mediterranean type, tempered by the influence of the Atlantic Ocean. The average temperatures range between 1.8°C in January (year 2005) and 38°C in August (year 2012), with annual rainfall ranging from 403 to 557 mm, according to years. Rains is generally concentrated in November, December, and January, but their distribution can vary from year to year. The dry period is relatively long, lasting up to 6 months, but atmospheric humidity is high, especially in the western part, which partly compensates for the aridity of the climate. Climatic data classify the western sectors of the Maamora forest as "warm sub-humid bioclimate" and the eastern sectors as "temperate semi-arid bioclimate" (Belghazi & Mounir, 2016). The natural vegetation of the Maamora consists mainly of cork oak trees (*Quercus suber* L.) and a few specimens of the Maamora pear (*Pyrus mamorensis* Trabut), an endemic species to the region. Mastic trees (*Pistacia lentiscus* L.), wild olive (*Olea europaea* ssp. *oleaster*), and green olive trees or mock privet (*Phillyrea latifolia* L.) are found on soils with shallow sand and red soil. The dwarf palm, *Chamaerops humilis* L., grows in dense clumps in the more open areas. Phoenician juniper (*Juniperus phoenicea* L.) can be found along the Atlantic coast, particularly around Lake Boughaba. The shrub and herb communities are highly diverse, with 402 recorded species (Métro & Sauvage, 1955; Sauvage, 1961; Aafi, 2007). The parts of the forest which have been impacted by commercial plantations are covered with introduced species: pine (*Pinus pinaster* ssp. *atlantica* H. del Vill., *Pinus halepensis* Mill., and *Pinus pinea* L.), eucalyptus (mostly *Eucalyptus camaldulensis* Dehn), and black wattle (or tannin acacia) trees (*Acacia mearnsii* De Wild.).



**Figure 1.** Maamora forest in Morocco and location of the studied stations (S1, S2, S3).


Site S1 is close to the sea and has a low rate of reforestation with commercial plantations. Site S2 is in the core of the forest, with a medium level of reforestation with commercial plantations. Finally, site S3 is located at the edge of the forest, under more marked continental influences, and shows a high rate of reforestation with commercial plantations. These differences in geographical location and species composition between the three study sites reflect the variations in vegetation conditions and climate within the Maamora forest.

**Collection sites.** Three sampling stations (named S1, S2, and S3, respectively) were chosen according to their location in the forest, their distance from the sea, and whether the plots were fenced (Fig. 1). Fencing prevents pruning and wood cutting by the local population and prevents cattle grazing in the undergrowth, which preserves the herbaceous and low shrub cover.

*Station S1* (Canton A; Fig. 2A). Located in the western part of the forest (34°12'33.34"N, 06°35'54.78"W), at an elevation of 37 m, it belongs to the warm sub-humid bioclimate with maritime influences. This station was characterized by an open cork oak forest where cork oak was mixed with tannin acacia, within a semi-fenced plot.

*Station S2* (Canton C; Fig. 2C). Located in the middle part of the forest (34°04'57.73"N, 06°24'38.31"W), at an elevation of 134 m, it belongs to the warm sub-humid bioclimate with maritime influences. This plot of old cork oak trees with a very dense understory was completely fenced and thus, protected from grazing.

*Station S3* (Canton E; Fig. 2B). Located in the eastern part of the forest (34°09'36.1"N, 06°07'04.34"W), at an elevation of 164 m, it belongs to the semiarid bioclimate with temperate winters and continental influences. This unfenced station comprised a mixed cork oak forest with eucalyptus trees.



**Figure 2.** Sampling stations. **A.** Canton A (S1); **B.** Canton E (S3) and **C.** Canton C (S2).

**Sampling methods.** Several sampling methods were used: visual capture when checking the traps; and passive sampling systems with three types of traps:

- 1) Window trapping (interception traps): The traps consisted of a large transparent plexiglass panel (73 cm x 42 cm) that intercepted the beetles that struck it in flight. The insects were collected in a gutter fixed at the base of the trap, filled with water, and mixed with saline solution and a few drops of dishwashing detergent to reduce surface tension. This trapping method is effective for collecting xylophagous and saproxylic species (Ranius & Jansson, 2002; Brustel, 2004; Bouget et al., 2008). Their position was 1.20 m from the ground, in plots where dead wood was present (Bouget & Noblecourt, 2005).
- 2) Pitfall traps (Barber traps): The traps were made with plastic cups with a top diameter of 60 mm, placed with the opening at the ground level, and covered with wooden branches to prevent them from being trampled by animals. Pitfall traps allow for the capture of soil-dwelling invertebrates (Nageleisen & Bouget, 2009). The distance between the traps was 20 metres.
- 3) Coloured bowls: These traps consisted of coloured bowls (yellow (YCT), orange (OCT), white (WCT), and blue (BCT)) with a diameter of 15 cm and a height of 13 cm. They were filled halfway with a preservation mixture (soapy water + salt) and placed at a height of 1.5 m above the ground level, in cork oak clearings to attract species attracted to flowers.

The different traps were installed at three stations in Cantons A, C, and E. In each station, 13 traps were set up at a distance ranging from 15 to 20 metres apart (8 Barber traps, 4 coloured traps one of each of the following yellow, white, orange, and blue colours - and one window trap). Traps were checked every 3 weeks for a 7-month period (April to October) during two consecutive seasons (2021 and 2022). The specimens collected were first identified to the genus level using specific identification keys for each

family. For species confirmation, they were compared to verified specimens in the museum's collections of the CIRF (Centre for Innovation, Research, and Training) and the Museum of the Scientific Institute of Rabat (Morocco). Carabidae and Tenebrionidae identifications were confirmed by Dr. Piero Leo (Italian entomologist (Cagliari) specialist in Tenebrionidae). The specimens collected were counted and only individuals identified to the species level were studied. The species studied in this work are preserved in the museum's collections of the CIRF.

The distribution of the captured species and their level of endemism in Morocco were checked using the *Catalogue of Palaearctic Coleoptera* by Löbl & Smetana (2003, 2004, 2006, 2007, 2008, 2010, 2011, 2013). To assess the rarity and threat status of saproxylic species, we referred to the IUCN Red List of Threatened Species (2022), the European Red List of saproxylic beetles (Nieto & Alexander, 2010; Cáliz et al., 2018), the Atlas of endangered invertebrates of Spain (Critically Endangered and Endangered Species) (Verdu & Galante, 2009), the Red List of Invertebrates of Spain (Verdu & Galante, 2006), the Red List of Mediterranean saproxylic beetles (García et al., 2018), the Red List of saproxylic and phytophagous beetles of Limousin (Chambord et al., 2013), and the studies of several authors, including Chavanon (2018), Labrique & Chavanon (2001), Prudhomme (2016), Velle (2004, 2011), and Villemant & Fraival (1993).

## RESULTS

256 species belonging to 42 families were identified, but only 216 species were included in this study: 29 species were endemic to the region; 26 species were new to Morocco; 53 were considered saproxylic; 108 species were already known from Moroccan forest ecosystems. The families and species are presented in alphabetical order in Table 1. The species were classified according to the following criteria: [S] Saproxylic species; [R] Saproxylic species listed in the Red List of the Mediterranean region; [N] Species newly recorded for Morocco and [E] Endemic species. Table 2 compares the number of species according to the collection sites, with a distinction between saproxylic species (S), endemic species (E) and those that are new to Morocco (N).

### *Taxonomic hierarchy*

#### **Class Insecta Linnaeus, 1758**

#### **Order Coleoptera Linnaeus, 1758**

#### **Family Anobiidae Fleming, 1821**

#### ***Stegobium paniceum* (Linnaeus, 1758) [S]**

**Material examined.** 1 spec., Canton C (34°04'57.73"N, 06°24'38.31"W), 28.V.2021, BCT; 1 spec., 05.VIII.2021, WCT; 3 spec., 13.V.2022, window trap. 1 spec., Canton A (34°12'33.34"N, 06°35'54.78"W), 13.V.2022, window trap; 1 spec., 02.VI.2022, window trap.

**Distribution.** Palaearctic (Europe, Asia, and North Africa – Borowski, 2007).

#### ***Xyletinus bucephalus* (Illiger, 1807) [S]**

**Material examined.** 1 spec., Canton A (34°12'33.34"N, 06°35'54.78"W), 22.VII.2022, BCT.

**Distribution.** Palaearctic (Europe, Asia, and North Africa – Borowski, 2007).

#### **Family Anthicidae Latreille, 1819**

#### ***Notoxus monoceros* (Linnaeus, 1760) [N] [S]**

**Material examined.** 1 spec., Canton A (34°12'33.34"N, 06°35'54.78"W), 28.V.2021, Barber trap; 1 spec., 02.VI.2022, WCT. 117 specs., Canton C (34°04'57.73"N, 06°24'38.31"W), 02.VI.2022, Barber traps and YCT, OCT, WCT, and BCT; 2 specs., 23.VI.2022, OCT, and BCT.

**Distribution.** Palaearctic (Europe and Asia – Chandler et al., 2008).

***Notoxus trifasciatus* Rossi, 1792 [S]**

**Material examined.** 1 spec., Canton C (34°04'57.73"N, 06°24'38.31"W), 30.IV.2021, Barber trap; 5 specs., 13.V.2022, YCT, WCT, and BCT; 5 specs., 2.VI.2022, YCT, WCT, and BCT. 45 specs., Canton A (34°12'33.34"N, 06°35'54.78"W), 28.V.2021, Barber trap; 5 specs., 2.VII.2021, Barber trap; 1 spec., 13.V.2022, YCT; 7 specs., 2.VI.2022, OCT, WCT, and BCT; 6 specs., 23.VI.2022, WCT and window trap. 11 specs., Canton E (34°09'36.1"N, 06°07'04.34"W), 13.V.2022, Barber trap and WCT.

**Distribution.** Palaearctic (Europe, Asia, and North Africa – Chandler et al., 2008).

***Stricticollis transversalis* (A. Villa & J.B. Villa, 1833) [S]**

**Material examined.** 1 spec., Canton C (34°04'57.73"N, 06°24'38.31"W), 30.IV.2021, Barber trap; 5 specs., 13.V.2022, YCT, WCT, and BCT; 5 specs., 02.VI.2022, YCT, WCT, and BCT. 45 specs., Canton A (34°12'33.34"N, 06°35'54.78"W), 28.V.2021, Barber trap, 5 specs., 02.VII.2021, Barber trap; 1 spec., 13.V.2022, YCT; 7 specs., 02.VI.2022, OCT, WCT, and BCT; 6 specs., 23.VI.2022, WCT and window trap. 11 specs., Canton E (34°09'36.1"N, 06°07'04.34"W), 13.V.2022, Barber trap and WCT.

**Distribution.** Palaearctic (Europe, Asia, and North Africa – Chandler et al., 2008).

**Family Anthribidae Billberg, 1820*****Bruchela rufipes rufipes* (Olivier, 1790)**

**Material examined.** 2 specs., Canton A (34°12'33.34"N, 06°35'54.78"W), 20.IV.2022, Barber trap.

**Distribution.** Western Palaearctic (Europe and Morocco – Magnano & Alonso-Zaragoza, 2011).

**Family Brentidae Billberg, 1820*****Kalcapion semivittatum* (Gyllenhal, 1833)**

**Material examined.** 3 specs., Canton C (34°04'57.73"N, 06°24'38.31"W), 06.IV.2021, OCT.

**Distribution.** Palaearctic (Europe, Asia, and North Africa – Sforzi, 2011).

***Ceratapion robusticorne* (Desbrochers des Loges, 1866)**

**Material examined.** 1 spec., Canton C (34°04'57.73"N, 06°24'38.31"W), 20.IV.2022, BCT.

**Distribution.** Western Palaearctic (Italy, Malta, Portugal, Spain, Algeria, Morocco, and Tunisia – Sforzi, 2011).

**Family Buprestidae Leach, 1815*****Acmaeodera bipunctata kureimatica* Escalera, 1914**

**Material examined.** 1 spec., Canton E (34°09'36.1"N, 06°07'04.34"W), 02.VII.2021, WCT; 1 spec., 02.VI.2022, YCT; 1 spec., 23.VI.2022, Barber trap.

**Distribution.** North Africa (Algeria and Morocco – Volkovitsh, 2006).

***Acmaeoderella adspersula* (Illiger, 1803) [S]**

**Material examined.** 3 specs., Canton E (34°09'36.1"N, 06°07'04.34"W), 30.IV.2021, WCT and window trap; 10 specs., 28.V.2021, YCT, BCT, and OCT; 9 specs., 02.VII.2021, YCT, WCT, and OCT; 2 specs., 20.IV.2022, YCT; 10 specs., 13.V.2022, YCT, OCT, and WCT; 26 specs., 02.VI.2022, YCT, OCT, WCT, and BCT; 31 specs., 23.VI.2022, Barber traps and YCT, WCT, and BCT. 1 spec., Canton C (34°04'57.73"N, 06°24'38.31"W), 28.V.2021, Barber trap, 5 specs., 13.V.2022; YCT, WCT, and BCT; 326 specs., 02.VI.2022, Barber traps and YCT, OCT, WCT, and BCT; 42 specs., 23.VI.2022, Barber traps and YCT, OCT, WCT, and BCT; 3 specs., 22.VII.2022, Barber trap. 1 spec., Canton A (34°12'33.34"N, 06°35'54.78"W), 13.V.2022, Barber trap; 3 specs., 23.VI.2022, YCT, OCT, and WCT.

**Distribution.** Palaearctic (Europe, Asia, Morocco, Algeria, and Tunisia – Volkovitsh, 2006; Villemant & Fraival, 1993).

***Acmaeoderella discoidea* (Fabricius, 1787)**

**Material examined.** 1 spec., Canton E (34°09'36.1"N, 06°07'04.34"W), 30.IV.2021, WCT; 2 specs., 28.V.2021, YCT; 1 spec., 20.IV.2022, YCT; 2 specs., 13.V.2022, OCT and WCT. 1 spec., Canton C (34°04'57.73"N, 06°24'38.31"W), 30.IV.2021, Barber trap; 4 specs., 28.V.2021, Barber trap; 37 specs., 20.IV.2022, Barber trap; 80 specs., 13.V.2022, YCT and Barber trap; 2 specs., 02.VI.2022, Barber trap and OCT. 4 specs., Canton A (34°12'33.34"N, 06°35'54.78"W), 28.V.2021, Barber trap; 3 specs., 13.V.2022, WCT and Barber trap; 1 spec., 02.VI.2022, Barber trap.

**Distribution.** Palaearctic (Europe, North Africa, Syria, and Israel – Volkovitsh, 2006).

***Acmaeoderella lanuginosa* (Gyllenhal, 1817) (Fig. 3A)**

**Material examined.** 5 specs., Canton E (34°09'36.1"N, 06°07'04.34"W), 30.IV.2021, WCT; 5 specs., 28.V.2021, YCT and OCT; 3 specs., 20.IV.2022, YCT; 5 specs., 13.V.2022, YCT, WCT, and BCT; 1 spec., 02.VI.2022, BCT. 1 spec., Canton C (34°04'57.73"N, 06°24'38.31"W), 19.III.2021, Barber trap; 13 specs., 20.IV.2022, Barber trap; 28 specs., 02.VI.2022, Barber trap and YCT, OCT, and BCT; 3 specs., 23.VI.2022, Barber trap and BCT. 2 specs., Canton A (34°12'33.34"N, 06°35'54.78"W), 20.IV.2022, YCT, and Barber trap.

**Distribution.** Western Palaearctic (Europe, Morocco, Algeria, Tunisia, Sinai, and Israel – Volkovitsh, 2006).

***Agrilus biguttatus* (Fabricius, 1777) [S]**

**Material examined.** 1 spec., Canton A (34°12'33.34"N, 06°35'54.78"W), 13.V.2022, sight capturing.

**Distribution.** Palaearctic (Europe, Iran, Syria, Algeria, and Morocco – Jendek, 2006).

***Agrilus graminis mamorensis* Théry, 1930 [E] [S]**

**Material examined.** 5 specs., Canton C (34°04'57.73"N, 06°24'38.31"W), 30.IV.2021, YCT; 7 specs., 28.V.2021, YCT, OCT, WCT, and BCT; 1 spec., 02.VII.2021, OCT; 1 spec., 05.VIII.2021, WCT. 4 specs., Canton A (34°12'33.34"N, 06°35'54.78"W), 28.V.2021, YCT and WCT.

**Distribution.** Morocco (Jendek, 2006).

***Anthaxia anatolica ferulae* Gené, 1839**

**Material examined.** 1 spec., Canton C (34°04'57.73"N, 06°24'38.31"W), 20.IV.2022, BCT.

**Distribution.** Western Palaearctic (Spain, France, Italy, Portugal, Algeria, Tunisia, and Morocco – Bily, 2006).

***Anthaxia millefolii polychloros* Abeille de Perrin, 1894**

**Material examined.** 2 specs., Canton E (34°09'36.1"N, 06°07'04.34"W), 28.V.2021, YCT; 1 spec., 02.VI.2022, YCT. 1 spec., Canton A (34°12'33.34"N, 06°35'54.78"W), 13.V.2022, YCT; 1 spec., 02.VI.2022, YCT.

**Distribution.** Western Palaearctic (Europe and North Africa – Bily, 2006).

***Anthaxia nitidula* (Linnaeus, 1758) [S] [N]**


**Material examined.** 1 spec., Canton E (34°09'36.1"N, 06°07'04.34"W), 20.IV.2022, WCT; 2 specs., 13.V.2022, YCT and BCT.

**Distribution.** Western Palaearctic (Europe and Algeria – Bily, 2006; Prudhomme, 2016).

***Anthaxia umbellatarum umbellatarum* (Fabricius, 1787)**

**Material examined.** 1 spec., Canton E (34°09'36.1"N, 06°07'04.34"W), 28.V.2021, YCT; 1 spec., 02.VI.2022, YCT. 2 specs., Canton C (34°04'57.73"N, 06°24'38.31"W), 19.III.2021, Barber trap; 3 specs., 28.V.2021, YCT; 5 specs., 13.V.2022, YCT and OCT; 3 specs., 02.VI.2022, YCT. 1 spec., Canton A (34°12'33.34"N, 06°35'54.78"W), 20.IV.2022, YCT.

**Distribution.** Palaearctic (Europe, Asia, and North Africa – Bily, 2006).



**Figure 3.** The beetles associated with the Maamora cork oak forest in the north-west of Morocco. **A.** *Acmaeoderella lanuginosa* (Gyllenhall, 1817); **B.** *Perotis unicolor igniventris* Escalera; 1914, **C.** *Carabus faminii favieri* Fairmaire, 1859; **D.** *Pterostichus globosus pecoudi* (Antoine, 1922); **E.** *Sphodrus leucophthalmus* (Linnaeus, 1758); **F.** *Cicindela maroccana* Fabricius, 1801; **G.** *Trichoferus ilicis* Sama, 1987; **H.** *Chrysolina americana* (Linnaeus, 1758); **I.** *Cryptocephalus rugicollis* Olivier, 1792; **J.** *Adalia decempunctata* (Linnaeus, 1758); **K.** *Brachyderes pubescens* Boheman, 1833; **L.** *Attagenus trifasciatus* (Fabricius, 1787); **M.** *Dermestes undulatus* Brahm, 1790; **N.** *Lacon punctatus punctatus* (Herbst, 1779) (photos by H. Habbaz and N. Maatouf).

***Habroloma triangulare* (Lacordaire, 1835)**

**Material examined.** 1 spec., Canton C (34°04'57.73"N, 06°24'38.31"W), 02.VII.2021, WCT; 1 spec., 20.IV.2022, YCT; 2 specs., 02.VI.2022, YCT and OCT; 1 spec., 23.VI.2022, YCT; 2 specs., 22.VII.2022, YCT and BCT. 3 specs., Canton A (34°12'33.34"N, 06°35'54.78"W), 20.IV.2022, Barber trap.

**Distribution.** Western Palaearctic (Spain, France, Italy, Portugal, Algeria, Tunisia, and Morocco – Kuban, 2006).

***Perotis unicolor igniventris* Escalera, 1914 [E] [S] (Fig. 3B)**

**Material examined.** 1 spec., Canton C (34°04'57.73"N, 06°24'38.31"W), 20.IV.2022, WCT; 1 spec., 13.V.2022, BCT.

**Distribution.** Morocco (Kuban, 2006).

**Family: Cantharidae Imhoff, 1856*****Malthinus flaveolus* (Herbst, 1786) [N]**

**Material examined.** 11 specs., Canton C (34°04'57.73"N, 06°24'38.31"W), 19.III.2021, OCT, and WCT; 1 spec., 06.IV.2021, WCT.

**Distribution.** Western Palaearctic (Europe – Kazantsev & Brancuçi, 2007).

**Family Carabidae Latreille, 1802*****Acinopus picipes* (Olivier, 1795)**

**Material examined.** 1 spec., Canton E (34°09'36.1"N, 06°07'04.34"W), 28.V.2021, Barber trap; 1 spec., 16.IX.2021 Barber trap. 1 spec., Canton A (34°12'33.34"N, 06°35'54.78"W), 30.IV.2021, Barber trap; 5 specs., 28.V.2021, Barber trap; 1 spec., 16.IX.2021, Barber trap.

**Distribution.** Palaearctic (Europe, Asia and Morocco – Kataev et al., 2003; Benyahia et al., 2016).

***Agonum numidicum* (Lucas, 1846)**

**Material examined.** 1 spec., Canton A (34°12'33.34"N, 06°35'54.78"W), 20.IV.2022, Barber trap.

**Distribution.** Western Palaearctic (France, Italy, Portugal, Spain, Morocco – Bousquet, 2003a).

***Amara aenea* (De Geer, 1774)**

**Material examined.** 3 specs., Canton A (34°12'33.34"N, 06°35'54.78"W), 28.V.2021, Barber trap; 3 specs., 20.IV.2022, YCT and BCT; 5 specs., 13.V.2022, Barber trap.

**Distribution.** Palaearctic (Europe, Asia, and North Africa – Hieke, 2003).

***Anisodactylus binotatus* (Fabricius, 1787)**

**Material examined.** 3 specs., Canton A (34°12'33.34"N, 06°35'54.78"W), 13.V.2022, window trap.

**Distribution.** Palaearctic (Europe, Asia, and North Africa – Ito, 2003).

***Calathus circumseptus* Germar, 1824**

**Material examined.** 2 specs., Canton A (34°12'33.34"N, 06°35'54.78"W), 19.III.2021, Barber trap; 3 specs., 29.X.2021, Barber trap.

**Distribution.** Western Palaearctic (Europe, Algeria, and Morocco – Hovorka & Sciaky, 2003).

***Calathus fuscipes algericus* Gautier des Cottés, 1866**

**Material examined.** 2 specs., Canton C (34°04'57.73"N, 06°24'38.31"W), 19.III.2021, Barber trap; 1 spec., 30.IV.2021, Barber trap; 2 specs., 29.X.2021, Barber trap.

**Distribution.** North Africa (Algeria, Tunisia, and Morocco – Hovorka & Sciaky, 2003).

***Calathus mollis atticus* Gautier des Cottes, 1867**

**Material examined.** 1 spec., Canton A (34°12'33.34"N, 06°35'54.78"W), 22.VII.2022, YCT.

**Distribution.** North Africa (Algeria, Libya, Tunisia, and Morocco – Hovorka & Sciaky, 2003).

***Carabus faminii favieri* Fairmaire, 1859 [E] (Fig. 3C)**

**Material examined.** 1 spec., Canton E (34°09'36.1"N, 06°07'04.34"W), 29.X.2021, Barber trap. 7 specs., Canton C (34°04'57.73"N, 06°24'38.31"W), 19.III.2021, Barber trap; 2 specs., 06.IV.2021, Barber trap; 2 specs., 30.IV.2021, Barber trap; 1 spec., 28.V.2021, Barber trap, 2 specs., 29.X.2021, Barber trap. 5 specs., Canton A (34°12'33.34"N, 06°35'54.78"W), 19.III.2021, Barber trap; 2 specs., 29.X.2021, Barber trap.

**Distribution.** Morocco (Bousquet et al., 2003).

***Carabus rugosus rugosus* Fabricius, 1792 [E]**

**Material examined.** 1 spec., Canton C (34°04'57.73"N, 06°24'38.31"W), 06.IV.2021, Barber trap.

**Distribution.** Morocco (Bousquet et al., 2003).

***Cicindela maroccana* Fabricius, 1801 (Fig. 3F)**

**Material examined.** 1 spec., Canton E (34°09'36.1"N, 06°07'04.34"W), 29.X.2021, OCT; 1 spec., 02.VI.2022, Barber trap. 5 specs., Canton C (34°04'57.73"N, 06°24'38.31"W), 29.X.2021, Barber trap. 2 specs., Canton A (34°12'33.34"N, 06°35'54.78"W), 20.IV.2022, Barber trap; 1 spec., 13.V.2022, Barber trap.

**Distribution.** Western Palaearctic (Portugal, Spain, Tunisia, and Morocco – Puchkov & Matalin, 2003).

***Dromius agilis* (Fabricius, 1787) [N]**

**Material examined.** 1 spec., Canton E (34°09'36.1"N, 06°07'04.34"W), 29.X.2021, window trap.

**Distribution.** Palaearctic (Europe and Asia – Kabak, 2003).

***Graphipterus exclamationis* (Fabricius, 1792)**

**Material examined.** 2 specs., Canton A (34°12'33.34"N, 06°35'54.78"W), 05.VIII.2021, Barber trap.

**Distribution.** North Africa (Algeria, Tunisia, and Morocco – Hurka, 2003).

***Harpalus attenuatus* Stephens, 1828**

**Material examined.** 1 spec., Canton A (34°12'33.34"N, 06°35'54.78"W), 19.III.2021, Barber trap; 6 specs., 20.IV.2022, Barber trap; 2 specs., 13.V.2022, Barber trap; 5 specs., 02.VI.2022, Barber trap.

**Distribution.** Palaearctic (Europe, Asia, and North Africa – Kataev et al., 2003).

***Harpalus serripes serripes* (Quensel, 1806)**

**Material examined.** 1 spec., Canton A (34°12'33.34"N, 06°35'54.78"W), 20.IV.2022, Barber trap.

**Distribution.** Palaearctic (Europe, Asia, and North Africa – Kataev et al., 2003).

***Microlestes abeillei brisouti* Holdhaus, 1912**

**Material examined.** 1 spec., Canton A (34°12'33.34"N, 06°35'54.78"W), 06.IV.2021, window trap; 1 spec., 16.IX.2021, Barber trap; 3 specs., 29.X.2021, Barber trap. 3 specs., Canton C (34°04'57.73"N, 06°24'38.31"W), 20.IV.2022, YCT, BCT, and WCT; 1 spec., 13.V.2022, WCT.

**Distribution.** North Africa (Kabak, 2003).

***Ocys tachysoides* (Antoine, 1933)**

**Material examined.** 1 spec., Canton C (34°04'57.73"N, 06°24'38.31"W), 29.X.2021, window trap.

**Distribution.** Western Palaearctic (Portugal and Morocco – Marggi et al., 2003).

***Paradromius linearis linearis* (Olivier, 1795)**

**Material examined.** 1 spec., Canton C (34°04'57.73"N, 06°24'38.31"W), 13.V.2022, WCT.

**Distribution.** Palaearctic (Europe, Asia, and North Africa – Kabak, 2003).

***Philorhizus quadrisignatus* (Dejean, 1825) [N]**

**Material examined.** 1 spec., Canton C (34°04'57.73"N, 06°24'38.31"W), 05.VIII.2021, YCT.

**Distribution.** Palaearctic (Europe, Algeria, and Israel – Kabak, 2003).

***Poecilus baeticus gharbensis* (Alluaud, 1927) [E]**

**Material examined.** 4 specs., Canton A (34°12'33.34"N, 06°35'54.78"W), 30.IV.2021, Barber trap and YCT.

**Distribution.** Morocco (Bousquet, 2003b).

***Pterostichus elongatus* (Duftschmid, 1812)**

**Material examined.** 2 specs., Canton E (34°09'36.1"N, 06°07'04.34"W), 30.IV.2021, Barber trap; 9 specs., 29.X.2021, Barber trap; 1 spec., 20.IV.2022, Barber trap. 1 spec., Canton C (34°04'57.73"N, 06°24'38.31"W), 19.III.2021, Barber trap; 35 specs., 06.IV.2021, Barber trap; 13 specs., 30.IV.2021, Barber trap; 19 specs., 28.V.2021, Barber trap; 37 specs., 29.X.2021, Barber trap. 3 specs., Canton A (34°12'33.34"N, 06°35'54.78"W), 06.IV.2021, Barber trap; 2 specs., 30.IV.2021, Barber trap; 1 spec., 28.V.2021, Barber trap; 1 spec., 05.VIII.2021, Barber trap; 5 specs., 29.X.2021, Barber trap; 2 specs., 20.IV.2022, Barber trap.

**Distribution.** Palaearctic (Europe, Morocco, Afghanistan, Kazakhstan, and Turkey – Bousquet, 2003b).

***Pterostichus globosus pecoudi* (Antoine, 1922) [E] (Fig. 3D)**

**Material examined.** 1 spec., Canton C (34°04'57.73"N, 06°24'38.31"W), 19.III.2021, Barber trap.

**Distribution.** Morocco (Bousquet, 2003b).

***Scarites terricola terricola* Bonelli, 1813**

**Material examined.** 5 specs., Canton E (34°09'36.1"N, 06°07'04.34"W), 30.IV.2021, Barber trap; 3 specs., 28.V.2021, Barber trap; 1 spec., 29.X.2021, Barber trap; 9 specs., 20.IV.2022, Barber trap; 2 specs., 13.V.2022, Barber trap; 2 specs., 02.VI.2022, Barber trap; 1 spec., 23.VI.2022, Barber trap. 9 specs., Canton C (34°04'57.73"N, 06°24'38.31"W), 30.IV.2021, Barber trap; 6 specs., 28.V.2021, Barber trap; 3 specs., 29.X.2021, Barber trap; 6 specs., 20.IV.2022, Barber trap; 1 spec., 13.V.2022, Barber trap. 3 specs., Canton A (34°12'33.34"N, 06°35'54.78"W), 06.IV.2021, Barber trap; 6 specs., 20.IV.2022, Barber trap.

**Distribution.** Palaearctic (Europe, Asia, and North Africa – Balkenohl, 2003).

***Sphodrus leucophthalmus* (Linnaeus, 1758) (Fig. 3E)**

**Material examined.** 1 spec., Canton A (34°12'33.34"N, 06°35'54.78"W), 06.IV.2021, Barber trap; 1 spec., 29.X.2021, Barber trap.

**Distribution.** Palaearctic (Europe, Asia, and North Africa – Casale, 2003).

***Syntomus foveatus* (Geoffroy, 1785)**

**Material examined.** 2 specs., Canton A (34°12'33.34"N, 06°35'54.78"W), 30.IV.2021, Barber trap; 6 specs., 28.V.2021, Barber trap.

**Distribution.** Palaearctic (Europe, Asia, and North Africa – Kabak, 2003).

***Trymosternus colombati* Antoine, 1934 [E]**

**Material examined.** 1 spec., Canton C (34°04'57.73"N, 06°24'38.31"W), 30.IV.2021, Barber trap.

**Distribution.** Morocco (Kabak, 2003).

**Family Cerambycidae Latreille, 1802*****Nathrius brevipennis* (Mulsant, 1839) [S] [R]**

**Material examined.** 1 spec., Canton A (34°12'33.34"N, 06°35'54.78"W), 28.V.2021, window trap.

**Distribution.** Palaearctic (Europe, Asia, and North Africa – Sama & Löbl, 2010a).

***Stenopterus ater* (Linnaeus, 1767) [S] [R]**

**Material examined.** 1 spec., Canton C (34°04'57.73"N, 06°24'38.31"W), 13.V.2022, YCT. 1 spec., Canton A (34°12'33.34"N, 06°35'54.78"W), 13.V.2022, OCT.

**Distribution.** Western Palaearctic (Europe and North Africa – Sama & Löbl, 2010a).

***Stictoleptura tangeriana* (Tournier, 1875) [E] [S]**

**Material examined.** 2 specs., Canton E (34°09'36.1"N, 06°07'04.34"W), 20.IV.2022, WCT and BCT.

**Distribution.** Morocco (Villemant & Fraval, 1993; Sama & Löbl, 2010b).

***Trichoferus ilicis* Sama, 1987 [E] [S] [R] (Fig. 3G)**

**Material examined.** 1 spec., Canton C (34°04'57.73"N, 06°24'38.31"W), 19.IX.2021, window trap. 1 spec., Canton A (34°12'33.34"N, 06°35'54.78"W), 28.V.2021, WCT.

**Distribution.** Morocco (Sama & Löbl, 2010a).

**Family Chrysomelidae Latreille, 1802*****Bruchidius biguttatus* (Olivier, 1800)**

**Material examined.** 1 spec., Canton E (34°09'36.1"N, 06°07'04.34"W), 05.VIII.2021, BCT; 4 specs., 16.IX.2021, WCT, YCT, and OCT, and window trap; 1 spec., 20.IV.2022, window trap; 1 spec., 22.VII.2022, YCT. 1 spec., Canton A (34°12'33.34"N, 06°35'54.78"W), 29.X.2021, WCT; 5 specs., 23.VI.2022, YCT, WCT, and window trap.

**Distribution.** Palaearctic (Europe, Asia, and North Africa – Anton, 2010).

***Bruchidius cisti* (Fabricius, 1775) [N]**

**Material examined.** 1 spec., Canton A (34°12'33.34"N, 06°35'54.78"W), 23.VI.2022, OCT.

**Distribution.** Western Palaearctic (Europe, Cyprus, and Turkey – Anton, 2010).

***Bruchidius rubiginosus* (Desbrochers des Loges, 1869)**

**Material examined.** 1 spec., Canton E (34°09'36.1"N, 06°07'04.34"W), 20.IV.2022, Barber trap. 1 spec., Canton A (34°12'33.34"N, 06°35'54.78"W), 13.V.2022, WCT.

**Distribution.** Western Palaearctic (Europe and North Africa – Anton, 2010).

***Cassida flaveola* Thunberg, 1794 [N]**

**Material examined.** 1 spec., Canton A (34°12'33.34"N, 06°35'54.78"W), 13.V.2022, sight capturing.

**Distribution.** Palaearctic (Europe, Algeria, China, and Russia – Borowiec & Sekerka, 2010).

***Cassida vittata* Villers, 1789**

**Material examined.** 1 spec., Canton E (34°09'36.1"N, 06°07'04.34"W), 29.IX.2021, BCT.

**Distribution.** Palaearctic (Europe, Asia, and North Africa – Borowiec & Sekerka, 2010).

***Chrysolina americana* (Linnaeus, 1758) (Fig. 3H)**

**Material examined.** 2 specs., Canton C (34°04'57.73"N, 06°24'38.31"W), 30.IV.2021, WCT.

**Distribution.** Western Palaearctic (Europe, North Africa, and Turkey – Kippenberg, 2010).

***Chrysolina kocheri* (Codina Padilla, 1961) [E]**

**Material examined.** 1 spec., Canton A (34°12'33.34"N, 06°35'54.78"W), 20.IV.2022, Barber trap.

**Distribution.** Morocco (Kippenberg, 2010).

***Cryptocephalus rabatensis* Pic, 1953 [E]**

**Material examined.** 1 spec., Canton C (34°04'57.73"N, 06°24'38.31"W), 19.III.2021, YCT; 5 specs., 30.IV.2021, YCT, OCT, and WCT; 1 spec., 20.IV.2022, BCT. 1 spec., Canton A (34°12'33.34"N, 06°35'54.78"W), 19.III.2021, YCT; 2 specs., 30.IV.2021, YCT and WCT.

**Distribution.** Morocco (Lopatin et al., 2010).

***Cryptocephalus rugicollis* Olivier, 1792 (Fig. 3I)**

**Material examined.** 2 specs., Canton C (34°04'57.73"N, 06°24'38.31"W), 19.III.2021, YCT. 1 spec., Canton A (34°12'33.34"N, 06°35'54.78"W), 20.IV.2022, BCT.

**Distribution.** Western Palaearctic (Europe, North Africa, Jordan, and Turkey – Lopatin et al., 2010).

**Family Cleridae Latreille, 1802*****Opilo domesticus* (Sturm, 1837) [S] [R]**

**Material examined.** 5 specs., Canton C (34°04'57.73"N, 06°24'38.31"W), 05.VIII.2021, WCT and YCT; 2 specs., 19.IX.2021, WCT and window trap. 1 spec., Canton A (34°12'33.34"N, 06°35'54.78"W), 16.IX.2021, window trap.

**Distribution.** Western Palaearctic (Europe, Morocco, Madeira Island, Tunisia – Villemant & Fraival, 1993; Löbl et al., 2007).

**Family Coccinellidae Latreille, 1807*****Adalia decempunctata* (Linnaeus, 1758) (Fig. 3J)**

**Material examined.** 1 spec., Canton E (34°09'36.1"N, 06°07'04.34"W), 29.X.2021, window trap. 1 spec., Canton C (34°04'57.73"N, 06°24'38.31"W), 30.IV.2021, WCT; 2 specs., 05.VIII.2021, OCT. 2 specs., Canton A (34°12'33.34"N, 06°35'54.78"W), 19.III.2021, YCT; 1 spec., 06.IV.2021, YCT; 1 spec., 30.IV.2021, window trap; 1 spec., 29.X.2021, OCT.

**Distribution.** Palaearctic (Europe, Asia, and North Africa – Kovar, 2007).

***Coccinella algerica* Kovář, 1977**

**Material examined.** 4 specs., Canton E (34°09'36.1"N, 06°07'04.34"W), 20.IV.2022, YCT and WCT. 1 spec., Canton A (34°12'33.34"N, 06°35'54.78"W), 20.IV.2022, YCT; 1 spec., 13.V.2022, WCT.

**Distribution.** Western Palaearctic (Canarian Islands, Gibraltar, Italy, and North Africa – Bensusan et al., 2006; Kovar, 2007).

***Hyperaspis duvergeri* Fürsch, 1985**

**Material examined.** 1 spec., Canton A (34°12'33.34"N, 06°35'54.78"W), 02.VI.2022, YCT.

**Distribution.** Western Palaearctic (Europe, Morocco, and Algeria – Kovar, 2007).

***Oenopia lyncea lyncea* (Olivier, 1808)**

**Material examined.** 1 spec., Canton C (34°04'57.73"N, 06°24'38.31"W), 28.V.2021, window trap.

**Distribution.** Western Palaearctic (France, Italy, Portugal, Spain, Algeria, Morocco, and Tunisia – Kovar, 2007).

***Oenopia conglobata* (Linnaeus, 1758)**

**Material examined.** 2 specs., Canton A (34°12'33.34"N, 06°35'54.78"W), 05.VIII.2021, YCT.

**Distribution.** Palaearctic (Europe, Asia and Morocco – Kovar, 2007; Benyahia et al., 2016).

***Platynaspis luteorubra* (Goeze, 1777)**

**Material examined.** 1 spec., Canton C (34°04'57.73"N, 06°24'38.31"W), 02.VII.2021, YCT. 1 spec., Canton A (34°12'33.34"N, 06°35'54.78"W), 13.V.2022, BCT; 1 spec., 02.VI.2022, YCT.

**Distribution.** Palaearctic (Europe, Asia and North Africa – Kovar, 2007).

***Rhyzobius chrysomeloides* (Herbst, 1792)**

**Material examined.** 1 spec., Canton E (34°09'36.1"N, 06°07'04.34"W), 20.IV.2022, WCT.

**Distribution.** Western Palaearctic (Europe, North Africa, and Turkey – Kovar, 2007).

***Rhyzobius litura* (Fabricius, 1787)**

**Material examined.** 1 spec., Canton C (34°04'57.73"N, 06°24'38.31"W), 02.VII.2021, YCT.

**Distribution.** Western Palaearctic (Europe, North Africa, and Turkey – Kovar, 2007).

***Rodolia cardinalis* (Mulsant, 1850)**

**Material examined.** 1 spec., Canton C (34°04'57.73"N, 06°24'38.31"W), 06.IV.2021, OCT; 1 spec., 13.V.2022, window coloured trap. 1 spec., Canton A (34°12'33.34"N, 06°35'54.78"W), 20.IV.2022, YCT.

**Distribution.** Palaearctic (Europe, Asia, and North Africa – Kovar, 2007).

***Scymnus subvillosus* (Goeze, 1777)**

**Material examined.** 1 spec., Canton C (34°04'57.73"N, 06°24'38.31"W), 05.VIII.2021, OCT.

**Distribution.** Palaearctic (Europe, Asia, and North Africa – Kovar, 2007).

**Family Corylophidae LeConte, 1852*****Orthoperus anxius* Mulsant & Rey, 1861**

**Material examined.** 1 spec., Canton A (34°12'33.34"N, 06°35'54.78"W), 05.VIII.2021, WCT.

**Distribution.** Western Palaearctic (France, Italy, Spain, Algeria, and Morocco – Bowstead, 2007).

**Family Cryptophagidae Kirby, 1837*****Cryptophagus dentatus* (Herbst, 1793)**

**Material examined.** 1 spec., Canton C (34°04'57.73"N, 06°24'38.31"W), 06.IV.2021, window trap. 1 spec., Canton A (34°12'33.34"N, 06°35'54.78"W), 22.VII.2022, YCT.

**Distribution.** Palaearctic (Europe, Asia, and North Africa – Johnson et al., 2007).

**Family Curculionidae Latreille, 1802*****Brachycerus barbarus* (Linnaeus, 1758)**

**Material examined.** 1 spec., Canton C (34°04'57.73"N, 06°24'38.31"W), 29.X.2021, Barber trap.

**Distribution.** Western Palaearctic (Spain, France, Italy, Tunisia, and Morocco – Colonnelli, 2011).

***Brachycerus muricatus* Olivier, 1790**

**Material examined.** 1 spec., Canton C (34°04'57.73"N, 06°24'38.31"W), 29.X.2021, Barber trap.

**Distribution.** Western Palaearctic (Europe, Algeria, Morocco, Tunisia, and Turkey – Colonnelli, 2011).

***Brachyderes pubescens* Boheman, 1833 (Fig. 3K)**

**Material examined.** 2 specs., Canton C (34°04'57.73"N, 06°24'38.31"W), 28.V.2021, WCT and window trap; 4 specs., 02.VII.2021, YCT and OCT; 1 spec., 05.VIII.2021, window trap; 1 spec., 19.IX.2021, WCT; 3 specs., 29.X.2021, YCT, WCT, and BCT; 1 spec., 13.V.2022, BCT.

**Distribution.** Western Palaearctic (Spain, Portugal, France, Italy, Algeria, and Morocco – Villemant & Fraval, 1993; Pelletier, 2013).

***Brachypera zoilus* (Scopoli, 1763)**

**Material examined.** 2 specs., Canton A (34°12'33.34"N, 06°35'54.78"W), 29.X.2021, Barber trap.

**Distribution.** Palaearctic (Europe, Asia, and North Africa – Skuhrovec, 2013).

***Charagnus griseus* (Fabricius, 1775)**

**Material examined.** 1 spec., Canton E (34°09'36.1"N, 06°07'04.34"W), 02.VI.2022, Barber trap.

**Distribution.** Western Palaearctic (Europe, Algeria, Morocco, Libya, Syria, Cyprus, and Turkey – Velazquez de Castro, 2013).

***Ceutorynchus contractus* (Marsham., 1802)**

**Material examined.** 1 spec., Canton C (34°04'57.73"N, 06°24'38.31"W), 20.IV.2022, YCT.

**Distribution.** Palaearctic (Europe, Asia, Morocco, and Libya – Colonnelli, 2013).

***Coeliodes ilicis* (Bedel, 1885)**

**Material examined.** 1 spec., Canton C (34°04'57.73"N, 06°24'38.31"W), 30.IV.2021, YCT; 1 spec., 06.IV.2021, window trap. 1 spec., Canton A (34°12'33.34"N, 06°35'54.78"W), 06.IV.2021, window trap.

**Distribution.** Palaearctic (Europe, Asia, Algeria, and Morocco – Colonnelli, 2013).

***Coeliodes ruber* (Marsham, 1802)**

**Material examined.** 1 spec., Canton C (34°04'57.73"N, 06°24'38.31"W), 13.V.2022, WCT.

**Distribution.** Western Palaearctic (Europe, Algeria, Morocco, Syria, and Israel – Villemant & Fraval, 1993; Colonnelli, 2013).

***Gronops lunatus* (Fabricius, 1775)**

**Material examined.** 9 specs., Canton A (34°12'33.34"N, 06°35'54.78"W), 29.X.2021, Barber trap.

**Distribution.** Western Palaearctic (Europe, Canary Islands, Morocco, and Tunisia – Meregalli, 2013).

***Liophloeus tessulatus* (O. F. Müller, 1776) [N]**

**Material examined.** 1 spec., Canton A (34°12'33.34"N, 06°35'54.78"W), 05.VIII.2021, Barber trap.

**Distribution.** Western Palaearctic (Europe – Meregalli, 2013).

***Listroderes costirostris* Schoenherr, 1826**

**Material examined.** 8 specs., Canton A (34°12'33.34"N, 06°35'54.78"W), 29.X.2021, Barber trap.

**Distribution.** Morocco, France, Portugal, Canary Islands, South Korea (Meregalli, 2013), Japan, South Africa, Australia, Pacific Islands, Hawaii, New Caledonia, New Zealand, Norfolk Island, United States, Argentina, Bolivia, Brazil, Chile, Uruguay, Venezuela (Morrone, 2013).

***Maurobaris spoliata* (Boheman, 1836)**

**Material examined.** 2 specs., Canton A (34°12'33.34"N, 06°35'54.78"W), 20.IV.2022, Barber trap.

**Distribution.** Western Palaearctic (Greece, France, Italy, Spain, Tunisia, Algeria, and Morocco – Prena, 2011).

***Mecinus pascuorum* (Gyllenhal, 1813) [N]**

**Material examined.** 1 spec., Canton E (34°09'36.1"N, 06°07'04.34"W), 13.V.2022, Barber trap. 1 spec., Canton A (34°12'33.34"N, 06°35'54.78"W), 02.VI.2022, WCT.

**Distribution.** Palaearctic (Europe, Canary Islands, and Asia – Caldara, 2013).

***Mecinus pyrastrer* (Herbst, 1795)**

**Material examined.** 1 spec., Canton A (34°12'33.34"N, 06°35'54.78"W), 20.IV.2022, YCT.

**Distribution.** Palaearctic (Europe, Asia, and North Africa – Caldara, 2013).

***Micrelus ferrugatus* (Perris, 1847)**

**Material examined.** 2 specs., Canton A (34°12'33.34"N, 06°35'54.78"W), 19.III.2021, YCT; 1 spec., 06.IV.2021, YCT.

**Distribution.** Western Palaearctic (Europe, Canary Islands, Morocco, and Turkey – Colonnelli, 2013).

***Microplontus molitor* (Gyllenhal, 1837)**

**Material examined.** 1 spec., Canton A (34°12'33.34"N, 06°35'54.78"W), 29.X.2021, Barber trap.

**Distribution.** Western Palaearctic (Europe and North Africa – Caldara, 2013).

***Orchestes erythropus* (Germar, 1821)**

**Material examined.** 3 specs., Canton C (34°04'57.73"N, 06°24'38.31"W), 05.VIII.2021, OCT; 1 spec., 02.VI.2022, WCT.

**Distribution.** Western Palaearctic (Europe, Algeria, Morocco, and Tunisia – Villemant & Fraval, 1993; Caldara, 2013).

***Orchestes irroratus maroccanus* (Roudier, 1954)**

**Material examined.** 1 spec., Canton C (34°04'57.73"N, 06°24'38.31"W), 06.IV.2021, Barber trap.

**Distribution.** North Africa (Algeria, Morocco, and Tunisia – Villemant & Fraval, 1993; Caldara, 2013).

***Orchestes rusci* (Herbst, 1795) [N]**

**Material examined.** 1 spec., Canton A (34°12'33.34"N, 06°35'54.78"W), 22.VII.2022, YCT.

**Distribution.** Palaearctic (Europe and Asia – Caldara, 2013).

***Otiorhynchus cribricollis* Gyllenhal, 1834**

**Material examined.** 1 spec., Canton E (34°09'36.1"N, 06°07'04.34"W), 16.IX.2021, Barber trap.

**Distribution.** Palaearctic (Europe, Asia, and North Africa – Magnano & Alonso-Zarazaga, 2013).

**Family Dasytidae Laporte, 1840*****Dasytes nigroaeneus* Küster, 1850**

**Material examined.** 1 spec., Canton E (34°09'36.1"N, 06°07'04.34"W), 30.IV.2021, window trap; 7 specs., 20.IV.2022, YCT, WCT, and BCT. 307 specs., Canton C (34°04'57.73"N, 06°24'38.31"W), 19.III.2021, YCT, WCT, and OCT; 175 specs., 06.IV.2021, OCT and WCT, and window trap; 18 specs., 30.IV.2021, window trap, and YCT and OCT; 4 specs., 28.V.2021, window trap, and BCT; 13 specs., 20.IV.2022, OCT, YCT, and WCT, and window trap; 4 specs., 13.V.2022, OCT, YCT, and WCT. 2 specs., 2 specs., Canton A (34°12'33.34"N, 06°35'54.78"W), 19.III.2021, YCT; 06.IV.2021, window trap; 14 specs., 30.IV.2021, window trap, and WCT; 23 specs., 20.IV.2022, YCT, OCT, and BCT; 6 specs., 13.V.2022, YCT, OCT, and BCT; 2 specs., 02.VI.2022, WCT, and window trap.

**Distribution.** Western Palaearctic (Europe, Syria, Algeria, and Morocco – Mayor, 2007a).

***Dasytes terminalis* Jacquelin du Val, 1863**

**Material examined.** 3 specs., Canton E (34°09'36.1"N, 06°07'04.34"W), 30.IV.2021, YCT and WCT and a window trap; 2 specs., 20.IV.2022, OCT and WCT. 4 specs., Canton C (34°04'57.73"N, 06°24'38.31"W), 06.IV.2021, window trap; 2 specs., 30.IV.2021, YCT; 2 specs., 20.IV.2022, BCT, and Barber trap, 1 spec., 13.V.2022; BCT. 1 spec., Canton A (34°12'33.34"N, 06°35'54.78"W), 13.V.2022, OCT.

**Distribution.** Western Palaearctic (France, Portugal, Spain, Algeria, and Morocco – Villemant & Fraval, 1993; Mayor, 2007a).

## Family Dermestidae Latreille, 1804

### *Anthrenus angustefasciatus* Ganglbauer, 1904

**Material examined.** 1 spec., Canton C (34°04'57.73"N, 06°24'38.31"W), 30.IV.2021, window trap; 1 spec., 20.IV.2022, WCT; 2 specs., 13.V.2022, WCT, and window trap. 1 spec., Canton E (34°09'36.1"N, 06°07'04.34"W), 20.IV.2022, WCT. 1 spec., Canton A (34°12'33.34"N, 06°35'54.78"W), 20.IV.2022, WCT, 1 spec., 13.V.2022; WCT.

**Distribution.** Western Palaearctic (Spain, Portugal, Italy, and Morocco – Hava, 2007b).

### *Anthrenus museorum* (Linnaeus, 1761)

**Material examined.** 1 spec., Canton C (34°04'57.73"N, 06°24'38.31"W), 19.III.2021, WCT; 1 spec., 06.IV.2021, window trap; 3 specs., 13.V.2022, YCT; 1 spec., 02.VI.2022, YCT. 4 specs., Canton E (34°09'36.1"N, 06°07'04.34"W), 20.IV.2022, YCT, BCT, and WCT; 1 spec., 02.VI.2022, YCT. 1 spec., Canton A (34°12'33.34"N, 06°35'54.78"W), 13.V.2022, BCT.

**Distribution.** Palaearctic (Europe, Asia, and North Africa – Hava, 2007b).

### *Anthrenus pimpinellae* (Fabricius, 1775)

**Material examined.** 1 spec., Canton E (34°09'36.1"N, 06°07'04.34"W), 20.IV.2022, WCT. 1 spec., Canton C (34°04'57.73"N, 06°24'38.31"W), 13.V.2022, YCT. 1 spec., Canton A (34°12'33.34"N, 06°35'54.78"W), 13.V.2022, YCT.

**Distribution.** Palaearctic (Europe, Asia, Egypt, and Morocco – Hava, 2007b).

### *Attagenus smirnovi* Zhantiev, 1973

**Material examined.** 4 specs., Canton E (34°09'36.1"N, 06°07'04.34"W), 20.IV.2022, WCT and BCT; 1 spec., 13.V.2022, BCT. 4 specs., Canton C (34°04'57.73"N, 06°24'38.31"W), 20.IV.2022, Barber traps; 1 spec., 02.VI.2022, WCT. 47 specs., Canton A (34°12'33.34"N, 06°35'54.78"W), 20.IV.2022, YCT, WCT, BCT, and Barber trap; 8 specs., 13.V.2022, BCT, WCT, and Barber trap; 1 spec., 02.VI.2022, BCT.

**Distribution.** Palaearctic (Europe, Morocco, Oman, Russia, and Saudi Arabia – Hava, 2007b).

### *Attagenus trifasciatus* (Fabricius, 1787) (Fig. 3L)

**Material examined.** 1 spec., Canton E (34°09'36.1"N, 06°07'04.34"W), 30.IV.2021, WCT. 1 spec., Canton C (34°04'57.73"N, 06°24'38.31"W), 06.IV.2021, window trap; 1 spec., 30.IV.2021, WCT; 10 specs., 28.V.2021, WCT, OCT, YCT, and BCT, and window trap; 94 specs., 13.V.2022, WCT, OCT, YCT, and BCT, and window trap; 2 specs., 02.VI.2022, OCT; 1 spec., 23.VI.2022, Barber trap. 5 specs., Canton A (34°12'33.34"N, 06°35'54.78"W), 13.V.2022, YCT and OCT.

**Distribution.** Western Palaearctic (Europe, North Africa, and Israel – Hava, 2007b).

### *Attagenus unicolor unicolor* (Brahm, 1791)

**Material examined.** 1 spec., Canton C (34°04'57.73"N, 06°24'38.31"W), 13.V.2022, Barber trap. 14 specs., Canton A (34°12'33.34"N, 06°35'54.78"W), 20.IV.2022, OCT, WCT, BCT, and Barber trap, 2 specs., 13.V.2022; BCT and OCT.

**Distribution.** Palaearctic (Europe, Asia, and North Africa – Hava, 2007b).

### *Dermestes frichii* Kugelann, 1792

**Material examined.** 1 spec., Canton E (34°09'36.1"N, 06°07'04.34"W), 28.V.2021, YCT. 1 spec., Canton C (34°04'57.73"N, 06°24'38.31"W), 06.IV.2021, Barber trap; 3 specs., 28.V.2021, Barber trap. 1 spec., Canton A (34°12'33.34"N, 06°35'54.78"W), 30.IV.2021, Barber trap; 5 specs., 28.V.2021, Barber trap; 5 specs., 13.V.2022, Barber trap.

**Distribution.** Palaearctic (Europe, Asia, and North Africa – Hava, 2007a).

***Dermestes undulatus* Brahm, 1790 (Fig. 3M)**

**Material examined.** 1 spec., Canton A (34°12'33.34"N, 06°35'54.78"W), 23.VI.2022, sight capturing.

**Distribution.** Palaearctic (Europe, North Africa, and Asia – Hava, 2007a).

***Dermestes peruvianus* Laporte de Castelnau, 1840**

**Material examined.** 1 spec., Canton A (34°12'33.34"N, 06°35'54.78"W), 13.V.2022, BCT.

**Distribution.** Palaearctic and Neotropical (Europe, North Africa, Asia, and South America – Hava, 2007a).

***Thorictus castaneus castaneus* Germar, 1834**

**Material examined.** 1 spec., Canton A (34°12'33.34"N, 06°35'54.78"W), 20.IV.2022, Barber trap; 1 spec., 23.VI.2022, Barber trap.

**Distribution.** Western Palaearctic (Morocco, Algeria, Libya, Egypt, Lebanon, and Syria – Löbl, 2007).

**Family Elateridae Leach, 1815*****Ampedus balteatus* (Linnaeus, 1758) [N] [S] [R]**

**Material examined.** 3 specs., Canton C (34°04'57.73"N, 06°24'38.31"W), 13.V.2022, WCT.

**Distribution.** Palaearctic (Europe and Asia – Cate, 2007).

***Agriotes sordidus* (Illiger, 1807) [S]**

**Material examined.** 7 specs., Canton A (34°12'33.34"N, 06°35'54.78"W), 13.V.2022, window trap and BCT.

**Distribution.** Western Palaearctic (Europe, Algeria, Canary Islands, and Morocco – Cate, 2007).

***Cardiophorus rufipes* (Goeze, 1777) [S]**

**Material examined.** 2 specs., Canton C (34°04'57.73"N, 06°24'38.31"W), 30.IV.2021, window trap and WCT; 1 spec., 20.IV.2022, window trap; 1 spec., 02.VI.2022, window trap. 1 spec., Canton A (34°12'33.34"N, 06°35'54.78"W), 28.V.2021, window trap; 1 spec., 05.VIII.2021, window trap.

**Distribution.** Palaearctic (Europe, Asia, and North Africa – Cate, 2007).

***Cebrio maculicollis* Fairmaire, 1856 [E] [S]**

**Material examined.** 2 specs., Canton A (34°12'33.34"N, 06°35'54.78"W), 02.VI.2022, YCT, and BCT.

**Distribution.** Morocco (Sanchez-Ruiz & Löbl, 2007).

***Drasterius bimaculatus* (Rossi, 1790) [S]**

**Material examined.** 1 spec., Canton A (34°12'33.34"N, 06°35'54.78"W), 05.VIII.2021, Barber trap.

**Distribution.** Palaearctic (Europe, North Africa, and Asia – Cate, 2007).

***Lacon punctatus punctatus* (Herbst, 1779) [S] [R] (Fig. 3N)**


**Material examined.** 1 spec., Canton C (34°04'57.73"N, 06°24'38.31"W), 30.IV.2021, sight capturing.

**Distribution.** Palaearctic (Europe, Asia, and Morocco – Cate, 2007).

**Family Geotrupidae Latreille, 1802*****Thorectes distinctus* Marseul, 1878 (Fig. 4A)**

**Material examined.** 11 specs., Canton E (34°09'36.1"N, 06°07'04.34"W), 29.X.2021, Barber trap; 1 spec., 20.IV.2022, Barber trap. 138 specs., Canton C (34°04'57.73"N, 06°24'38.31"W), 19.III.2021, Barber trap; 43 specs., 06.IV.2021, Barber trap; 9 specs., 30.IV.2021, Barber trap; 1 spec., 05.VIII.2021, Barber trap; 54 specs., 29.X.2021, Barber trap; 19 specs., 20.IV.2022, Barber trap. 17 specs., Canton A (34°12'33.34"N, 06°35'54.78"W), 19.III.2021, Barber trap; 7 specs., 06.IV.2021, Barber trap; 2 specs., 16.IX.2021, Barber trap; 11 specs., 29.X.2021, Barber trap; 2 specs., 20.IV.2022, Barber trap.

**Distribution.** North Africa (Algeria and Morocco – Löbl et al., 2006c).



**Figure 4.** The beetles associated with the Maamora cork oak forest in the north-west of Morocco. **A.** *Thorectes distinctus* Marseul, 1878; **B.** *Eulasia goudouti* (Laporte, 1840); **C.** *Hycleus rufipalpis* (Escalera, 1909); **D.** *Euserica mamorensis* Baraud, 1965; **E.** *Sphodroxia maroccana* Ley, 1923; **F.** *Dignomus dilophus* (Illiger, 1807) (photos by A. Samih); **G.** *Aethiessa floralis* (Fabricius, 1787); **H.** *Oxythyrea funesta* (Poda, 1761); **I.** *Scarabaeus cicatricosus* Lucas, 1846; **J.** *Platypus cylindrus* (Fabricius, 1792); **K.** *Akis tingitana* Lucas, 1859; **L.** *Blaps ovipennis* Seidlitz, 1893; **M.** *Pachychila punctata mamorensis* Antoine, 1942; **N.** *Pimelia chrysomeloides subbris* Koch, 1941 (photos by H. Habbaz and N. Maatouf).

***Typhaeus typhoeus* (Linnaeus, 1758)**

**Material examined.** 1 spec., Canton E (34°09'36.1"N, 06°07'04.34"W), 29.X.2021, OCT. 2 specs., Canton A (34°12'33.34"N, 06°35'54.78"W), 29.X.2021, Barber trap; 1 spec., 20.IV.2022, Barber trap.

**Distribution.** Western Palaearctic (Europe and Morocco – Löbl et al., 2006c).

**Family Glaphyridae Macleay, 1819*****Anthypna meles* (Fabricius, 1792)**

**Material examined.** 1 spec., Canton E (34°09'36.1"N, 06°07'04.34"W), 13.V.2022, WCT. 2 specs., Canton C (34°04'57.73"N, 06°24'38.31"W), 13.V.2022, BCT, and Barber trap.

**Distribution.** North Africa (Algeria, Tunisia, and Morocco – Nikodym & Bezdek, 2006).

***Eulasia goudouti* (Laporte, 1840) [E] (Fig. 4B)**

**Material examined.** 11 specs., Canton E (34°09'36.1"N, 06°07'04.34"W), 20.IV.2022, YCT, WCT, OCT, and BCT; 1 spec., 13.V.2022, BCT. 12 specs., Canton C (34°04'57.73"N, 06°24'38.31"W), 20.IV.2022, OCT, YCT, WCT, and BCT; 5 specs., 13.V.2022, YCT and BCT. 536 specs., Canton A (34°12'33.34"N, 06°35'54.78"W), 20.IV.2022, YCT, OCT, WCT, and BCT; 243 specs., 13.V.2022, OCT, WCT, and BCT.

**Distribution.** Morocco (Nikodym & Bezdek, 2006).

**Family Histeridae Gyllenhal, 1808*****Hister moerens* Erichson, 1834**

**Material examined.** 1 spec., Canton C (34°04'57.73"N, 06°24'38.31"W), 06.IV.2021, window trap.

**Distribution.** Western Palaearctic (Europe, Algeria, Tunisia, Morocco, and Turkey – Mazur, 2004).

***Hister maroccanus* Schmidt, 1887 [E]**

**Material examined.** 1 spec., Canton E (34°09'36.1"N, 06°07'04.34"W), 20.IV.2022, OCT.

**Distribution.** Morocco (Mazur, 2004).

***Hypocaccus brasiliensis* (Paykull, 1811)**

**Material examined.** 1 spec., Canton C (34°04'57.73"N, 06°24'38.31"W), 20.IV.2022, Barber trap; 1 spec., 02.VI.2022, OCT. 1 spec., Canton A (34°12'33.34"N, 06°35'54.78"W), 20.IV.2022, Barber trap.

**Distribution.** Palaearctic (Europe, Asia, and North Africa – Mazur, 2004).

***Hypocaccus rugiceps* (Duftschmid, 1805) [N]**

**Material examined.** 2 specs., Canton E (34°09'36.1"N, 06°07'04.34"W), 28.V.2021, WCT and YCT; 1 spec., 29.X.2021, Barber trap; 2 specs., 20.IV.2022, BCT. 3 specs., Canton C (34°04'57.73"N, 06°24'38.31"W), 20.IV.2022, Barber trap.

**Distribution.** Palaearctic (Europe, and China – Mazur, 2004).

***Pactolinus major* (Linnaeus, 1767)**

**Material examined.** 1 spec., Canton C (34°04'57.73"N, 06°24'38.31"W), 28.V.2021, Barber trap.

**Distribution.** Palaearctic (Europe, Asia, and North Africa – Mazur, 2004).

***Saprinus acuminatus* (Fabricius, 1798)**

**Material examined.** 1 spec., Canton E (34°09'36.1"N, 06°07'04.34"W), 05.VIII.2021, Barber trap; 2 specs., 29.X.2021, YCT and OCT; 2 specs., 20.IV.2022, WCT; 3 specs., 02.VI.2022, Barber trap and YCT. 10 specs., Canton C (34°04'57.73"N, 06°24'38.31"W), 30.IV.2021, Barber trap; 3 specs., 29.X.2021, Barber trap; 1 spec., 13.V.2022, Barber trap; 1 spec., 02.VI.2022, Barber trap. 14 specs., Canton A (34°12'33.34"N, 06°35'54.78"W), 20.IV.2022, Barber trap, YCT, WCT, and BCT; 6 specs., 13.V.2022, YCT, WCT, and BCT.

**Distribution.** Palaearctic (Europe, Asia, and North Africa – Mazur, 2004).

***Saprinus chalcites* (Illiger, 1807)**

**Material examined.** 8 specs., Canton E (34°09'36.1"N, 06°07'04.34"W), 20.IV.2022, YCT and Barber trap. 1 spec., Canton C (34°04'57.73"N, 06°24'38.31"W), 02.VI.2022, Barber trap. 4 specs., Canton A (34°12'33.34"N, 06°35'54.78"W), 20.IV.2022, YCT and BCT; 2 specs., 23.VI.2022, OCT, and WCT; 1 spec., 22.VII.2022, YCT.

**Distribution.** Palaearctic (Europe, Asia, and North Africa – Mazur, 2004).

***Saprinus figuratus* Marseul, 1855**

**Material examined.** 1 spec., Canton C (34°04'57.73"N, 06°24'38.31"W), 20.IV.2022, BCT; 11 specs., 02.VI.2022, Barber trap, YCT, OCT, and BCT. 23 specs., Canton A (34°12'33.34"N, 06°35'54.78"W), 13.V.2022, YCT, WCT, and BCT.

**Distribution.** Palaearctic (Spain, Asia, and North Africa – Mazur, 2004).

***Saprinus proximus simillimus* Wollaston, 1865**

**Material examined.** 1 spec., Canton C (34°04'57.73"N, 06°24'38.31"W), 06.IV.2021, Barber trap.

**Distribution.** North Africa (Mazur, 2004).

**Family Laemophloeidae Ganglbauer, 1899*****Cryptolestes ferrugineus* (Stephens, 1831)**

**Material examined.** 1 spec., Canton E (34°09'36.1"N, 06°07'04.34"W), 28.V.2021, window trap. 1 spec., Canton C (34°04'57.73"N, 06°24'38.31"W), 19.III.2021, OCT; 1 spec., 30.IV.2021, window trap. 1 spec., Canton A (34°12'33.34"N, 06°35'54.78"W), 05.VIII.2021, OCT.

**Distribution.** Palaearctic (Europe, Asia, Morocco, Canary Islands, and Madeira Island – Wegrzynowicz, 2007; Chavanon, 2018).

***Laemophloeus monilis* (Fabricius, 1787) [S]**

**Material examined.** 8 specs., Canton C (34°04'57.73"N, 06°24'38.31"W), 30.IV.2021, window trap; 1 spec., 13.V.2022, window trap.

**Distribution.** Western Palaearctic (Europe, Uzbekistan, Tunisia, Algeria and Morocco – Wegrzynowicz, 2007; Benyahia, 2016).

**Family Latridiidae Erichson, 1842*****Corticarina cavicollis* (Mannerheim, 1844) [N]**

**Material examined.** 4 specs., Canton C (34°04'57.73"N, 06°24'38.31"W), 06.IV.2021, window trap.

**Distribution.** Western Palaearctic and Nearctic (United States, Canada Venezuela, and Italy – Majka et al., 2009; Johnson, 2007).

***Corticarina curta* (Wollaston, 1854)**

**Material examined.** 1 spec., Canton E (34°09'36.1"N, 06°07'04.34"W), 28.V.2021, WCT; 7 specs., 16.IX.2021, WCT, BCT, OCT, and window; 1 spec., 20.IV.2022, YCT and BCT. 1 spec., Canton C (34°04'57.73"N, 06°24'38.31"W), 19.III.2021, OCT; 1 spec., 05.VIII.2021, window trap; 1 spec., 19.IX.2021, YCT; 1 spec., 29.X.2021, WCT; 1 spec., 20.IV.2022, WCT; 1 spec., 13.V.2022, BCT. 1 spec., Canton A (34°12'33.34"N, 06°35'54.78"W), 05.VIII.2021, OCT.

**Distribution.** Palaearctic (Europe, Asia, and North Africa – Johnson, 2007).

**Family Leiodidae Fleming, 1821*****Agathidium marocanum* (Hlisenkovsky, 1968) [E] [S]**

**Material examined.** 14 specs., Canton E (34°09'36.1"N, 06°07'04.34"W), 29.X.2021, Barber trap.

**Distribution.** Morocco (Perreau, 2004).

***Catops coracinus* Kellner, 1846**

**Material examined.** 2 specs., Canton C (34°04'57.73"N, 06°24'38.31"W), 19.III.2021, window trap; 23 specs., 06.IV.2021, Barber trap. 1 spec., Canton E (34°09'36.1"N, 06°07'04.34"W), 20.IV.2022, YCT.

**Distribution.** Palaearctic (Europe, Asia, Algeria, Tunisia, and Morocco – Perreau, 2004).

***Ptomaphagus tenuicornis mauritanicus* Jeannel, 1934**

**Material examined.** 3 specs., Canton C (34°04'57.73"N, 06°24'38.31"W), 29.X.2021, Barber trap.

**Distribution.** Western Palaearctic (Italy, Egypt, Libya, Tunisia, and Morocco – Perreau, 2004; de Faria e Silva et al., 2013).

**Family Meloidae Gyllenhaal, 1810*****Actenodia distincta* (Chevrolat, 1840)**

**Material examined.** 1 spec., Canton E (34°09'36.1"N, 06°07'04.34"W), 13.V.2022, YCT.

**Distribution.** Western Palaearctic (Italy, Morocco, Tunisia, and Algeria – Bologna, 2008).

***Cerocoma schaefferi* (Linnaeus, 1758) [N]**

**Material examined.** 1 spec., Canton E (34°09'36.1"N, 06°07'04.34"W), 20.IV.2022, WCT.

**Distribution.** Western Palaearctic (Europe and Turkey – Bologna, 2008).

***Hycleus duodecimpunctatus* (Olivier, 1811) [N]**

**Material examined.** 1 spec., Canton E (34°09'36.1"N, 06°07'04.34"W), 28.V.2021, OCT; 1 spec., 23.VI.2022, YCT. 1 spec., Canton A (34°12'33.34"N, 06°35'54.78"W), 20.IV.2022, Barber trap.

**Distribution.** Western Palaearctic (France, Portugal, and Spain – Bologna, 2008).

***Hycleus rufipalpis* (Escalera, 1909) (Fig. 4C)**

**Material examined.** 6 specs., Canton E (34°09'36.1"N, 06°07'04.34"W), 30.IV.2021, WCT; 27 specs., 28.V.2021, YCT, OCT, WCT, and BCT; 85 specs., 02.VII.2021, YCT, OCT, WCT, and BCT; 48 specs., 05.VIII.2021, YCT, OCT, WCT, and BCT; 22 specs., 13.V.2022, YCT, OCT, and WCT; 25 specs., 02.VI.2022, YCT, OCT, WCT, and BCT; 32 specs., 23.VI.2022, Barber trap, YCT, WCT, and BCT; 31 specs., 22.VII.2022, Barber trap and YCT. 2 specs., Canton C (34°04'57.73"N, 06°24'38.31"W), 30.IV.2021, Barber trap; 28 specs., 28.V.2021, WCT, YCT, and Barber trap; 38 specs., 02.VII.2021, Barber trap, YCT, OCT, and BCT; 1 spec., 20.IV.2022, BCT; 166 specs., 13.V.2022, Barber trap, YCT, WCT, OCT, and BCT; 575 specs., 02.VI.2022, Barber trap, YCT, WCT, OCT, and BCT; 79 specs., 23.VI.2022, Barber trap, YCT, WCT, OCT, and BCT; 2 specs., 22.VII.2022, Barber trap. 4 specs., Canton A (34°12'33.34"N, 06°35'54.78"W), 28.V.2021, Barber trap; 1 spec., 02.VII.2021, BCT; 14 specs., 13.V.2022, Barber trap, YCT, WCT, and BCT; 20 specs., 02.VI.2022, Barber trap, YCT, OCT, WCT, and BCT; 26 specs., 23.VI.2022, Barber trap, YCT, WCT, and BCT; 1 spec., 22.VII.2022, BCT.

**Distribution.** Western Palaearctic (Canary Islands and Morocco – Bologna, 2008).

**Family Melolonthidae Leach, 1819*****Euserica mamorensis* Baraud, 1965 [E] (Fig. 4D)**

**Material examined.** 2 specs., Canton C (34°04'57.73"N, 06°24'38.31"W), 30.IV.2021, OCT and WCT; 1 spec., 28.V.2021, window trap; 1 spec., 05.VIII.2021, window trap; 3 specs., 02.VI.2022, window trap. 4 specs., Canton A (34°12'33.34"N, 06°35'54.78"W), 30.IV.2021, Barber trap; 37 specs., 28.V.2021, Barber trap, YCT, OCT, WCT, BCT, and window trap; 5 specs., 02.VII.2021, OCT, WCT, BCT, and window trap; 1 spec., 02.VI.2022, window trap; 1 spec., 23.VI.2022, window trap. 3 specs., Canton E (34°09'36.1"N, 06°07'04.34"W), 13.V.2022, window trap; 1 spec., 23.VI.2022, window trap.

**Distribution.** Morocco (Bezdek, 2006a).

***Hoplia argentea* (Poda, 1761) [N]**

**Material examined.** 1 spec., Canton A (34°12'33.34"N, 06°35'54.78"W), 20.IV.2022, BCT.

**Distribution.** Western Palaearctic (Europe – Smetana, 2006a).

***Hoplia philanthus gagates* (Bedel, 1911)**

**Material examined.** 1 spec., Canton C (34°04'57.73"N, 06°24'38.31"W), 20.IV.2022, OCT. 1 spec., Canton A (34°12'33.34"N, 06°35'54.78"W), 20.IV.2022, OCT.

**Distribution.** Western Palaearctic (Spain and Morocco – Micó et al., 2003; Smetana, 2006a).

***Sphodroxia maroccana* Ley, 1923 [E] (Fig. 4E)**

**Material examined.** 1 spec., Canton E (34°09'36.1"N, 06°07'04.34"W), 02.VII.2021, Barber trap; 5 specs., 05.VIII.2021, Barber trap. 5 specs., Canton C (34°04'57.73"N, 06°24'38.31"W), 05.VIII.2021, Barber trap. 1 spec., Canton A (34°12'33.34"N, 06°35'54.78"W), 02.VII.2021, Barber traps; 4 specs., 23.VI.2022, BCT and YCT; 1 spec., 22.VII.2022, BCT.

**Distribution.** Morocco (Bezdek, 2006b).

**Family Melyridae Leach, 1815*****Charopus rotundatus* Erichson, 1840**

**Material examined.** 1 spec., Canton C (34°04'57.73"N, 06°24'38.31"W), 06.IV.2021, WCT.

**Distribution.** Western Palaearctic (Europe, Tunisia, Algeria, and Morocco – Mayor, 2007b).

***Colotes javeti* Jacquelin du Val, 1853**

**Material examined.** 2 specs., Canton C (34°04'57.73"N, 06°24'38.31"W), 06.IV.2021, Barber trap.

**Distribution.** Western Palaearctic (Spain, France, Italy, Portugal, Egypt, Tunisia, Algeria, and Morocco – Mayor, 2007b).

***Colotes punctatus* (Erichson, 1840)**

**Material examined.** 1 spec., Canton E (34°09'36.1"N, 06°07'04.34"W), 28.V.2021, 1 spec., OCT; 05.VIII.2021, window trap; 1 spec., 16.IX.2021, Barber trap; 1 spec., 13.V.2022, Barber trap; 1 spec., 02.VI.2022, WCT. 1 spec., Canton A (34°12'33.34"N, 06°35'54.78"W), 16.IX.2021, Barber trap; 2 specs., 22.VII.2022, Barber trap and WCT.

**Distribution.** Western Palaearctic (Europe and North Africa – Mayor, 2007b).

**Family Mordellidae Latreille, 1802*****Medimorda bipunctata* (Germar, 1827)**

**Material examined.** 2 specs., Canton E (34°09'36.1"N, 06°07'04.34"W), 02.VI.2022, WCT and YCT. 5 specs., Canton C (34°04'57.73"N, 06°24'38.31"W), 02.VI.2022, YCT, OCT, WCT, and BCT; 4 specs., 23.VI.2022, WCT and BCT. 1 spec., Canton A (34°12'33.34"N, 06°35'54.78"W), 23.VI.2022, WCT.

**Distribution.** Palaearctic (Europe, Asia, and North Africa – Horak, 2008).

***Mordella aculeata* (Linnaeus, 1758) [N]**

**Material examined.** 1 spec., Canton C (34°04'57.73"N, 06°24'38.31"W), 30.IV.2021, WCT; 6 specs., 20.IV.2022, WCT and OCT; 20 specs., 13.V.2022, YCT, OCT, and WCT; 3 specs., 02.VI.2022, YCT, OCT, and WCT. 3 specs., Canton E (34°09'36.1"N, 06°07'04.34"W), 20.IV.2022, WCT and BCT; 3 specs., 13.V.2022, YCT; 1 spec., 02.VI.2022, WCT. 5 specs., Canton A (34°12'33.34"N, 06°35'54.78"W), 13.V.2022, OCT, YCT, and WCT; 2 specs., 02.VI.2022, WCT.

**Distribution.** Palaearctic (Europe and Asia (Horak, 2008); Albania, Austria, Belgium, Bosnia Herzegovina, Bulgaria, Belarus, China, Croatia, Czech Republic, Denmark, France, Georgia, Germany, Greece, Hungary, Italy, Japan, Kazakhstan, Macedonia, Poland, Portugal, Romania, Russia, Serbia and Montenegro, Slovakia, Slovenia, South Korea, Spain, Sweden, Switzerland, Syria, Taiwan, and Turkey, Ukraine (Ruzzier et al., 2017).

***Variimorda ragusai* (Emery, 1876) [S]**

**Material examined.** 2 specs., Canton E (34°09'36.1"N, 06°07'04.34"W), 28.V.2021, WCT.

**Distribution.** Western Palaearctic (France, Italy, and Morocco – Horak, 2008).

**Family Nitidulidae Latreille, 1802*****Carpophilus hemipterus* (Linnaeus, 1758)**

**Material examined.** 5 specs., Canton A (34°12'33.34"N, 06°35'54.78"W), 13.V.2022, window trap.

**Distribution.** Palaearctic (Europe, Asia, Algeria, Canary Islands, Egypt, Libya, Madeira Island, Tunisia and Morocco – Jelinek & Audisio, 2007; Chavanon, 2018).

***Epuraea aestiva* (Linnaeus, 1758) [N] [S]**

**Material examined.** 26 specs., Canton C (34°04'57.73"N, 06°24'38.31"W), 13.V.2022, WCT, OCT, YCT, and BCT.

**Distribution.** Palaearctic (Europe and Asia – Jelinek & Audisio, 2007).

***Epuraea unicolor* (Olivier, 1790)**

**Material examined.** 1 spec., Canton C (34°04'57.73"N, 06°24'38.31"W), 30.IV.2021, YCT.

**Distribution.** Palaearctic (Europe, Asia, and North Africa – Jelinek & Audisio, 2007).

***Brassicogethes aeneus* (Fabricius, 1775)**

**Material examined.** 3 specs., Canton C (34°04'57.73"N, 06°24'38.31"W), 19.III.2021, OCT; 8 specs., 06.IV.2021, OCT and WCT; 1 spec., 28.V.2021, WCT; 1 spec., 20.IV.2022, YCT; 1 spec., 02.VI.2022, WCT. 1 spec., Canton A (34°12'33.34"N, 06°35'54.78"W), 02.VI.2022, window trap. 1 spec., Canton E (34°09'36.1"N, 06°07'04.34"W), 20.IV.2022, BCT.

**Distribution.** Palaearctic (Europe, Asia, and North Africa – Jelinek & Audisio, 2007).

**Family Oedemeridae Latreille, 1810*****Oedemera barbara* (Fabricius, 1792) [S]**

**Material examined.** 1 spec., Canton C (34°04'57.73"N, 06°24'38.31"W), 28.V.2021, YCT. 1 spec., Canton E (34°09'36.1"N, 06°07'04.34"W), 20.IV.2022, WCT; 1 spec., 13.V.2022, YCT.

**Distribution.** Western Palaearctic (Europe, North Africa, and Turkey – Švihla, 2008). Larvae in rotten wood, saproxylophagous. Anthophagous flower-feeding adults (Prudhomme, 2016).

***Oedemera simplex* (Linnaeus, 1767) [S]**

**Material examined.** 4 specs., Canton C (34°04'57.73"N, 06°24'38.31"W), 13.V.2022, YCT, OCT, and BCT; 2 specs., 02.VI.2022, YCT and WCT.

**Distribution.** Western Palaearctic (Europe and North Africa – Švihla, 2008). Larvae in rotten wood, saproxylophagous. Anthophagous flower-feeding adults (Prudhomme, 2016).

**Family Ptinidae Latreille, 1802*****Dignomus dilophus* (Illiger, 1807) [S] (Fig. 4F)**

**Material examined.** 1 spec., Canton E (34°09'36.1"N, 06°07'04.34"W), 05.VIII.2021, window trap. 1 spec., Canton C (34°04'57.73"N, 06°24'38.31"W), 29.X.2021, window trap. 1 spec., Canton A (34°12'33.34"N, 06°35'54.78"W), 19.III.2021, Barber trap.

**Distribution.** Western Palaearctic (Spain, Portugal, Algeria, Tunisia, and Morocco – Borowski, 2007).

***Dignomus gibbicollis* (P. H. Lucas, 1846) [S]**

**Material examined.** 2 specs., Canton C (34°04'57.73"N, 06°24'38.31"W), 06.IV.2021, Barber trap; 2 specs., 13.V.2022, Barber trap. 1 spec., Canton A (34°12'33.34"N, 06°35'54.78"W), 19.III.2021, Barber trap.

**Distribution.** Western Palaearctic (Spain, Portugal, Algeria, and Morocco – Borowski, 2007).

***Mesocoelopus niger* (P. W. J. Müller, 1821) [S]**

**Material examined.** 1 spec., Canton A (34°12'33.34"N, 06°35'54.78"W), 13.V.2022, Barber trap.

**Distribution.** Western Palaearctic (Europe, Algeria, Morocco, and Turkey – Borowski, 2007).

***Mizodorcatoma dommeri* (Rosenhauer, 1856) [S]**

**Material examined.** 1 spec., Canton E (34°09'36.1"N, 06°07'04.34"W), 13.V.2022, BCT.

**Distribution.** Western Palaearctic (France, Italy, Spain, Algeria, Morocco, and Tunisia – Borowski, 2007).

***Ptinus bidens* Olivier, 1790 [N]**

**Material examined.** 1 spec., Canton C (34°04'57.73"N, 06°24'38.31"W), 13.V.2022, window trap.

**Distribution.** Western Palaearctic (Europe and the Canary Islands – Borowski, 2007).

***Ptinus spitzyi* A. Villa & G. B. Villa, 1838**

**Material examined.** 1 spec., Canton E (34°09'36.1"N, 06°07'04.34"W), 29.X.2021, Barber trap. 1 spec., Canton C (34°04'57.73"N, 06°24'38.31"W), 19.III.2021, OCT. 1 spec., Canton A (34°12'33.34"N, 06°35'54.78"W), 19.III.2021, Barber trap.

**Distribution.** Western Palaearctic (Europe and North Africa – Borowski, 2007).

**Family Phalacridae Leach, 1815*****Olibrus affinis* (Sturm, 1807)**

**Material examined.** 4 specs., Canton C (34°04'57.73"N, 06°24'38.31"W), 02.VII.2021, WCT, BCT, and OCT.

**Distribution.** Palaearctic (Europe, Asia, and North Africa – Švec, 2007).

***Stilbus testaceus* (Panzer, 1796)**

**Material examined.** 1 spec., Canton C (34°04'57.73"N, 06°24'38.31"W), 13.V.2022, OCT. 1 spec., Canton A (34°12'33.34"N, 06°35'54.78"W), 20.IV.2022, Barber trap.

**Distribution.** Palaearctic (Europe, Asia, and North Africa – Švec, 2007).

**Family Scarabaeidae Latreille, 1802*****Aethiessa floralis* (Fabricius, 1787) (Fig. 4G)**

**Material examined.** 3 specs., Canton C (34°04'57.73"N, 06°24'38.31"W), 30.IV.2021, Barber traps; 20 specs., 20.IV.2022, WCT and Barber traps; 10 specs., 13.V.2022, YCT, OCT, WCT, BCT, and Barber traps. 8 specs., Canton E (34°09'36.1"N, 06°07'04.34"W), 20.IV.2022, WCT, YCT, and BCT. 114 specs., Canton A (34°12'33.34"N, 06°35'54.78"W), 20.IV.2022, YCT, OCT, WCT, BCT, and Barber trap; 62 specs., 13.V.2022, OCT, YCT, WCT, BCT, and Barber trap; 1 spec., 23.VI.2022, YCT.

**Distribution.** Western Palaearctic (Spain, Malta, France, Israel, and North Africa – Smetana, 2006b).

***Otophorus haemorrhoidalis* (Linnaeus, 1758)**

**Material examined.** 2 specs., Canton A (34°12'33.34"N, 06°35'54.78"W), 23.VI.2022, YCT and WCT.

**Distribution.** Palaearctic (Europe, Asia, Algeria, and Morocco – Dellacasa & Dellacasa, 2006).

***Labarrus lividus* (Olivier, 1789)**

**Material examined.** 1 spec., Canton A (34°12'33.34"N, 06°35'54.78"W), 23.VI.2022, Barber trap.

**Distribution.** Palaearctic (Europe, Asia, and North Africa – Dellacasa & Dellacasa, 2006).

***Esymus merdarius* (Fabricius, 1775)**

**Material examined.** 1 spec., Canton A (34°12'33.34"N, 06°35'54.78"W), 02.VI.2022, window trap.

**Distribution.** Palaearctic (Europe, Asia, Algeria, and Morocco – Dellacasa & Dellacasa, 2006).

***Subrinus sturmi* (Harold, 1870)**

**Material examined.** 1 spec., Canton E (34°09'36.1"N, 06°07'04.34"W), 23.VI.2022, BCT.

**Distribution.** Palaearctic (Europe, Asia, and North Africa – Dellacasa & Dellacasa, 2006).

***Onitis alexis septentrionalis* Balthasar, 1942**

**Material examined.** 1 spec., Canton A (34°12'33.34"N, 06°35'54.78"W), 22.VII.2022, BCT.

**Distribution.** Palaearctic (Europe, Asia, and North Africa – Bezdek & Krell, 2006).

***Onthophagus maki* (Illiger, 1803)**

**Material examined.** 1 spec., Canton A (34°12'33.34"N, 06°35'54.78"W), 13.V.2022, OCT; 1 spec., 23.VI.2022, WCT.

**Distribution.** Western Palaearctic (Spain, France, Portugal, Italy, and North Africa – Löbl et al., 2006b).

***Onthophagus similis* (Scriba, 1790)**

**Material examined.** 1 spec., Canton A (34°12'33.34"N, 06°35'54.78"W), 02.VI.2022, BCT.

**Distribution.** Palaearctic (Europe, Asia, and North Africa – Löbl et al., 2006b).

***Onthophagus vacca* (Linnaeus, 1767)**

**Material examined.** 3 specs., Canton C (34°04'57.73"N, 06°24'38.31"W), 06.IV.2021, Barber trap; 14 specs., 28.V.2021, Barber trap; 5 specs., 29.X.2021, Barber trap; 5 specs., 20.IV.2022, Barber trap; 4 specs., 13.V.2022, Barber trap; 7 specs., 02.VI.2022, Barber trap. 2 specs., Canton E (34°09'36.1"N, 06°07'04.34"W), 20.IV.2022, Barber trap; 1 spec., 02.VI.2022, Barber trap. 2 specs., Canton A (34°12'33.34"N, 06°35'54.78"W), 20.IV.2022, OCT; 2 specs., 13.V.2022, YCT and Barber trap.

**Distribution.** Palaearctic (Europe, Asia, and Morocco – Löbl et al., 2006b).

***Oryctes nasicornis grypus* (Illiger 1803)**

**Material examined.** 1 spec., Canton E (34°09'36.1"N, 06°07'04.34"W), 06.IV.2021, sight capturing.

**Distribution.** Palaearctic (Europe, Algeria, Tunisia, and Morocco – Krell, 2006).

***Oxythyrea funesta* (Poda, 1761) [S] (Fig. 4H)**

**Material examined.** 3 specs., Canton E (34°09'36.1"N, 06°07'04.34"W), 30.IV.2021, WCT; 45 specs., 28.V.2021, YCT, OCT, WCT, and BCT; 31 specs., 20.IV.2022, YCT, OCT, WCT, and BCT; 57 specs., 13.V.2022, YCT, OCT, WCT, and BCT; 14 specs., 02.VI.2022, YCT, OCT, and WCT. 1 spec., Canton C (34°04'57.73"N, 06°24'38.31"W), 20.IV.2022, Barber trap; 30 specs., 13.V.2022, YCT, OCT, and BCT; 14 specs., 02.VI.2022, YCT, OCT, WCT, and BCT; 1 spec., 23.VI.2022, OCT. 1 spec., Canton A (34°12'33.34"N, 06°35'54.78"W), 28.V.2021, Barber trap; 2 specs., 02.VII.2021, Barber trap and OCT; 14 specs., 20.IV.2022, Barber trap, OCT, WCT, and BCT; 52 specs., 13.V.2022, YCT, WCT, OCT, and BCT; 18 specs., 02.VI.2022, YCT, OCT, and WCT; 42 specs., 23.VI.2022, YCT, WCT, and BCT.

**Distribution.** Palaearctic (Europe, Asia, and North Africa – Smetana, 2006b).

***Phyllopertha horticola* (Linnaeus, 1758) [N]**

**Material examined.** 3 specs., Canton A (34°12'33.34"N, 06°35'54.78"W), 20.IV.2022, OCT and YCT.

**Distribution.** Palaearctic (Europe and Asia – Zorn, 2006).

***Protaetia (Netocia) morio heyrovskyi* (Balthasar, 1935)**

**Material examined.** 2 specs., Canton C (34°04'57.73"N, 06°24'38.31"W), 13.V.2022, YCT and OCT. 2 specs., Canton A (34°12'33.34"N, 06°35'54.78"W), 20.IV.2022, OCT, and WCT.

**Distribution.** North Africa (Algeria, Tunisia, and Morocco – Smetana, 2006b).

***Scarabaeus cicatricosus* Lucas, 1846 (Fig. 4I)**

**Material examined.** 1 spec., Canton E (34°09'36.1"N, 06°07'04.34"W), 05.VIII.2021, Barber trap; 1 spec., 13.V.2022, Barber trap. 1 spec., Canton C (34°04'57.73"N, 06°24'38.31"W), 19.III.2021, Barber trap; 1 spec., 06.IV.2021, Barber trap; 2 specs., 28.V.2021, Barber trap; 1 spec., 29.X.2021, Barber trap; 1 spec., 20.IV.2022, Barber trap.

**Distribution.** Western Palaearctic (Spain, Portugal, Algeria, and Morocco – Löbl et al., 2006a).

***Scarabaeus sacer* Linnaeus, 1758**

**Material examined.** 4 specs., Canton E (34°09'36.1"N, 06°07'04.34"W), 20.IV.2022, Barber trap. 1 spec., Canton A (34°12'33.34"N, 06°35'54.78"W), 13.V.2022, Barber trap.

**Distribution.** Palaearctic (Europe, Asia, and North Africa – Löbl et al., 2006a).

***Tropinota squalida pilosa* Brullé, 1832**

**Material examined.** 10 specs., Canton E (34°09'36.1"N, 06°07'04.34"W), 20.IV.2022, OCT, WCT, and BCT. 1 spec., Canton C (34°04'57.73"N, 06°24'38.31"W), 20.IV.2022, Barber trap; 1 spec., 13.V.2022, WCT. 3 specs., Canton A (34°12'33.34"N, 06°35'54.78"W), 20.IV.2022, WCT, BCT, and Barber trap; 2 specs., 13.V.2022, YCT.

**Distribution.** Palaearctic (Asia and North Africa – Smetana, 2006b).

***Tropinota hirta* (Poda, 1761)**

**Material examined.** 1 spec., Canton C (34°04'57.73"N, 06°24'38.31"W), 13.V.2022, Barber trap.

**Distribution.** Western Palaearctic (Europe, Morocco, Kazakhstan, and Turkey – Smetana, 2006b).

**Family Scolytidae Latreille, 1804*****Dryocoetes autographus* (Ratzeburg, 1837) [N] [S]**

**Material examined.** 1 spec., Canton C (34°04'57.73"N, 06°24'38.31"W), 30.IV.2021, window trap.

**Distribution.** Palaearctic (Europe, Asia, Algeria, and Madeira Island – Knizek, 2011).

***Ernoporicus fagi* (Fabricius, 1798) (= *Eidophelus fagi*) [N]**

**Material examined.** 1 spec., Canton A (34°12'33.34"N, 06°35'54.78"W), 16.IX.2021, BCT; 1 spec., 29.X.2021, OCT.

**Distribution.** Western Palaearctic (Europe and Turkey – Knizek, 2011).

***Hypoborus ficus* Erichson, 1836**

**Material examined.** 1 spec., Canton E (34°09'36.1"N, 06°07'04.34"W), 30.IV.2021, WCT.

**Distribution.** Palaearctic (Europe, Asia, and North Africa – Knizek, 2011).

***Platypus cylindrus* (Fabricius, 1792) (Fig. 4)**

**Material examined.** 1 spec., Canton E (34°09'36.1"N, 06°07'04.34"W), 30.IV.2021, window trap; 1 spec., 02.VII.2021, window trap; 1 spec., 23.VI.2022, BCT. 34 specs., Canton C (34°04'57.73"N, 06°24'38.31"W), 06.IV.2021, window trap. 1 spec., Canton A (34°12'33.34"N, 06°35'54.78"W), 13.V.2022, window trap; 2 specs., 23.VI.2022, window trap and OCT.

**Distribution.** Palaearctic (Europe, Asia, and North Africa – Knizek, 2011).

***Xyleborus monographus* (Fabricius, 1792) [S]**

**Material examined.** 1 spec., Canton E (34°09'36.1"N, 06°07'04.34"W), 02.VII. 2021, window trap. 1 spec., Canton C (34°04'57.73"N, 06°24'38.31"W), 19.III.2021, OCT; 7 specs., 06.IV.2021, window trap; 2 specs., 28.V.2021, window trap; 1 spec., 02.VII.2021, WCT; 1 spec., 05.VIII.2021, window trap; 1 spec., 20.IV.2022, window trap; 1 spec., 02.VI.2022, window trap; 8 specs., 22.VII.2022, window trap. 2 specs., Canton A (34°12'33.34"N, 06°35'54.78"W), 28.V.2021, window trap; 1 spec., 02.VII.2021, window trap; 1 spec., 20.IV.2022, YCT; 2 specs., 23.VI.2022, window trap and BCT.

**Distribution.** Palaearctic (Europe, Asia, Algeria, and Morocco – Villemant & Fraval, 1993; Knizek, 2011).

**Family Scaptiidae Mulsant, 1856*****Scaptia ophthalmica* Mulsant, 1856 (*S. schotti* Leblanc, 2012) [S]**

**Material examined.** 3 specs., Canton E (34°09'36.1"N, 06°07'04.34"W), 05.VIII.2021, window traps. 1 spec., Canton C (34°04'57.73"N, 06°24'38.31"W), 28.V.2021, WCT; 1 spec., 02.VII.2021, YCT; 2 specs., 05.VIII.2021, YCT and OCT; 1 spec., 19.IX.2021, YCT and window trap. 1 spec., Canton A (34°12'33.34"N, 06°35'54.78"W), 05.VIII.2021, OCT.

**Distribution.** Western Palaearctic (France, Spain, Italy, and Morocco – Leblanc et al., 2008). The specimens from Corsica, Algeria, Morocco, and Tunisia were considered as a new species and described by Leblanc (2012) under the name of *Scaptia schotti* n. sp. Saproxylophagous larvae, flower-feeding adults.

***Anaspis flava* (Linnaeus, 1758) [N] [S]**

**Material examined.** 2 specs., Canton C (34°04'57.73"N, 06°24'38.31"W), 19.III.2021, OCT; 12 specs., 06.IV.2021, WCT and window trap.

**Distribution.** Western Palaearctic (Europe and Lebanon – Leblanc et al., 2008).

**Family Scydmaenidae Leach, 1815*****Scydmaenus tingitanus* (Franz, 1962) [E] [S]**

**Material examined.** 1 spec., Canton C (34°04'57.73"N, 06°24'38.31"W), 02.VII.2021, Barber trap.

**Distribution.** Morocco (Vit, 2004).

**Family Silphidae Latreille, 1806*****Silpha tristis* Illiger, 1798**

**Material examined.** 1 spec., Canton E (34°09'36.1"N, 06°07'04.34"W), 30.IV.2021, Barber trap. 1 spec., Canton A (34°12'33.34"N, 06°35'54.78"W), 20.IV.2022, Barber trap.

**Distribution.** Palaearctic (Europe, Asia, and Morocco – Rozicka & Schneider, 2004; El Harche et al., 2022).

***Silpha olivieri* Bedel, 1887**

**Material examined.** 1 spec., Canton A (34°12'33.34"N, 06°35'54.78"W), 20.IV.2022, Barber trap.

**Distribution.** Western Palaearctic (Europe, Algeria, Tunisia, and Morocco – Rozicka & Schneider, 2004).

***Thanatophilus sinuatus* (Fabricius, 1775)**

**Material examined.** 1 spec., Canton C (34°04'57.73"N, 06°24'38.31"W), 19.III.2021, Barber trap.

**Distribution.** Palaearctic (Europe, Asia to Japan, Algeria, Tunisia, and Morocco – Rozicka & Schneider, 2004; El Harche et al., 2022).

**Family Staphylinidae Latreille, 1802*****Bolitobius cingulatus* (Mannerheim, 1830) [N]**

**Material examined.** 1 spec., Canton A (34°12'33.34"N, 06°35'54.78"W), 28.V.2021, Barber trap.

**Distribution.** Palaearctic (Europe, Asia – Smetana, 2004).

***Eusphalerum torquatum torquatum* (Marsham, 1802)**

**Material examined.** 1 spec., Canton C (34°04'57.73"N, 06°24'38.31"W), 19.III.2021, Barber trap.

**Distribution.** Western Palaearctic (Europe, Algeria, Madeira Island, and Morocco – Smetana, 2004).

***Gabrius appendiculatus* Sharp, 1910 [N]**

**Material examined.** 1 spec., Canton C (34°04'57.73"N, 06°24'38.31"W), 29.X.2021, Barber trap. 9 specs., Canton E (34°09'36.1"N, 06°07'04.34"W), 20.IV.2022, YCT, OCT, WCT, and BCT. 1 spec., Canton A (34°12'33.34"N, 06°35'54.78"W), 20.IV.2022, Barber trap.

**Distribution.** Palaearctic (Europe and Asia – Smetana, 2004).

***Micropeplus staphylinoides* Marsham, 1802**

**Material examined.** 2 specs., Canton C (34°04'57.73"N, 06°24'38.31"W), 06.IV.2021, Barber trap.

**Distribution.** Western Palaearctic (Europe, Algeria, Tunisia and Morocco – Smetana, 2004; Benyahia et al., 2016).

***Ontholestes marginalis* (Gené, 1836)**

**Material examined.** 1 spec., Canton A (34°12'33.34"N, 06°35'54.78"W), 20.IV.2022, YCT.

**Distribution.** Western Palaearctic (Spain, Portugal, Italy, France, Tunisia, Algeria, and Morocco – Smetana, 2004; Tronquet et al., 2019).

***Oxytelus sculptus* Gravenhorst, 1806 [S]**

**Material examined.** 1 spec., Canton E (34°09'36.1"N, 06°07'04.34"W), 30.IV.2021, Barber trap; 1 spec., 29.X.2021, window trap. 36 specs., Canton C (34°04'57.73"N, 06°24'38.31"W), 19.III.2021, Barber trap; 7 specs., 06.IV.2021, Barber trap; 46 specs., 29.X.2021, Barber trap and YCT. 1 spec., Canton A (34°12'33.34"N, 06°35'54.78"W), 19.III.2021, Barber trap.

**Distribution.** Palaearctic (Europe, Asia, and Morocco – Smetana, 2004).

***Xantholinus linearis* (Olivier, 1795)**

**Material examined.** 1 spec., Canton C (34°04'57.73"N, 06°24'38.31"W), 28.V.2021, window trap. 1 spec., Canton A (34°12'33.34"N, 06°35'54.78"W), 05.VIII.2021, window trap.

**Distribution.** Palaearctic (Europe, Asia, and North Africa; introduced in the Nearctic region – Smetana, 2004; Tronquet et al., 2019).

**Family Tenebrionidae Latreille, 1802*****Adelostoma sulcatum* Duponchel, 1827**

**Material examined.** 1 spec., Canton E (34°09'36.1"N, 06°07'04.34"W), 28.V.2021, Barber trap. 1 spec., Canton C (34°04'57.73"N, 06°24'38.31"W), 22.VII.2022, Barber trap.

**Distribution.** Palaearctic (Spain, Greece, Asia, and North Africa – Löbl et al., 2008a; Labrique & Chavanon, 2001).

***Akis tingitana* Lucas, 1859 [E] (Fig. 4K)**

**Material examined.** 2 specs., Canton C (34°04'57.73"N, 06°24'38.31"W), 30.IV.2021, Barber trap; 2 specs., 28.V.2021, Barber trap. 1 spec., Canton A (34°12'33.34"N, 06°35'54.78"W), 06.IV.2021, Barber trap.

**Distribution.** Morocco (Löbl et al., 2008a).

***Alphasida (Glabrasida) iblanensis reymondi* Antoine, 1952 [E]**

**Material examined.** 1 spec., Canton E (34°09'36.1"N, 06°07'04.34"W), 30.IV.2021, Barber trap. 1 spec., Canton C (34°04'57.73"N, 06°24'38.31"W), 06.IV.2021, Barber trap; 1 spec., 30.IV.2021, Barber trap; 2 specs., 28.V.2021, Barber trap.

**Distribution.** Morocco (Soldati, 2008; Pérez-Vera & Ávila, 2012).

***Arthrodeis globulosus* Escalera, 1922 [E] [S]**

**Material examined.** 2 specs., Canton C (34°04'57.73"N, 06°24'38.31"W), 02.VI.2022, Barber trap. 2 specs., Canton A (34°12'33.34"N, 06°35'54.78"W), 13.V.2022, Barber trap; 1 spec., 23.VI.2022, Barber trap.

**Distribution.** Morocco (Löbl et al., 2008a; Viñolas et al. 2017).

***Blaps ovipennis* Seidlitz, 1893 [E] [S] (Fig. 4L)**

**Material examined.** 1 spec., Canton C (34°04'57.73"N, 06°24'38.31"W), 06.IV.2021, Barber trap; 1 spec., 05.VIII.2021, Barber trap; 3 specs., 19.IX.2021, Barber trap; 1 spec., 29.X.2021, Barber trap; 3 specs., 20.IV.2022, Barber trap; 1 spec., 02.VI.2022, Barber trap.

**Distribution.** Morocco (Löbl et al., 2008a).

***Cheirodes brevicollis* (Wollaston, 1864) [S]**

**Material examined.** 1 spec., Canton A (34°12'33.34"N, 06°35'54.78"W), 13.V.2022, OCT.

**Distribution.** Palearctic (Spain, Malta, Italy, Asia, and North Africa – Iwan & Löbl, 2008; Chavanon, 2018).

***Cheirodes sardous sardous* Gené, 1839 [S]**

**Material examined.** 1 spec., Canton C (34°04'57.73"N, 06°24'38.31"W), 19.III.2021, Barber trap.

**Distribution.** Palearctic (Europe, Asia, and North Africa – Iwan & Löbl, 2008; Chavanon, 2018).

***Erodium granipennis* Fairmaire, 1871 [E]**

**Material examined.** 5 specs., Canton E (34°09'36.1"N, 06°07'04.34"W), 30.IV.2021, Barber trap; 8 specs., 28.V.2021, Barber trap; 1 spec., 02.VII.2021, Barber trap; 22 specs., 20.IV.2022, Barber trap; 16 specs., 13.V.2022, Barber trap; 133 specs., 02.VI.2022, Barber trap; 33 specs., 23.VI.2022, Barber trap. 9 specs., Canton C (34°04'57.73"N, 06°24'38.31"W), 19.III.2021, Barber trap; 3 specs., 06.IV.2021, Barber trap; 47 specs., 30.IV.2021, Barber trap; 35 specs., 28.V.2021, Barber trap; 93 specs., 20.IV.2022, Barber trap; 38 specs., 13.V.2022, Barber trap; 15 specs., 02.VI.2022, Barber trap; 1 spec., 23.VI.2022, Barber trap. 16 specs., Canton A (34°12'33.34"N, 06°35'54.78"W), 28.V.2021, Barber trap; 1 spec., 29.IX.2021, Barber trap; 20 specs., 20.IV.2022, Barber trap; 15 specs., 13.V.2022, Barber trap; 2 specs., 02.VI.2022, Barber trap; 1 spec., 23.VI.2022, Barber trap.

**Distribution.** Morocco (Löbl et al., 2008a).

***Heliotaurus ruficollis tangerianus* Escalera, 1922 [E]**

**Material examined.** 1 spec., Canton A (34°12'33.34"N, 06°35'54.78"W), 20.IV.2022, WCT.

**Distribution.** Morocco (Novak & Pettersson, 2008).

***Isomira melanophtalma* (Lucas, 1846)**

**Material examined.** 1 spec., Canton C (34°04'57.73"N, 06°24'38.31"W), 19.III.2021, OCT; 3 specs., 02.IV.2021, window trap; 2 specs., 30.IV.2021, window trap and OCT; 4 specs., 13.V.2022, OCT and window trap. 6 specs., Canton E (34°09'36.1"N, 06°07'04.34"W), 20.IV.2022, window trap.

**Distribution.** Western Palaearctic (Portugal, France, Italy, Croatia, Bosnia, Algeria, and Morocco – Novak & Pettersson, 2008).

***Latheticus oryzae* Waterhouse, 1880**

**Material examined.** 1 spec., Canton C (34°04'57.73"N, 06°24'38.31"W), 02.VII.2021, YCT.

**Distribution.** Palaearctic (Europe, Asia, and North Africa – Löbl et al., 2008a).

***Misolampus goudotii* Guérin-Méneville, 1834 [S]**

**Material examined.** 1 spec., Canton E (34°09'36.1"N, 06°07'04.34"W), 05.VIII.2021, Barber trap.

**Distribution.** Western Palaearctic (Morocco, Algeria, Balearic Islands – Löbl et al., 2008a; Rosas-Ramos et al., 2020).

***Pachychila alluaudi* (Peyerimhoff, 1925) [E] [S]**

**Material examined.** 1 spec., Canton A (34°12'33.34"N, 06°35'54.78"W), 20.IV.2022, Barber trap.

**Distribution.** Morocco (Löbl et al., 2008a).

***Pachychila punctata mamorensis* Antoine, 1942 [E] [S] (Fig. 4M)**

**Material examined.** 2 specs., Canton E (34°09'36.1"N, 06°07'04.34"W), 28.V.2021, Barber trap; 8 specs., 05.VIII.2021, Barber trap. 2 specs., Canton C (34°04'57.73"N, 06°24'38.31"W), 06.IV.2021, Barber trap; 11 specs., 30.IV.2021, Barber trap; 17 specs., 28.V.2021, Barber trap; 4 specs., 02.VII.2021, Barber trap; 18 specs., 05.VIII.2021, Barber trap; 15 specs., 19.IX.2021, Barber trap; 1 spec., 29.X.2021, Barber trap; 11 specs., 20.IV.2022, Barber trap; 2 specs., 13.V.2022, Barber trap; 1 spec., 02.VI.2022, Barber trap; 4 specs., 23.VI.2022, Barber trap; 149 specs., 22.VII.2022, Barber trap. 5 specs., Canton A (34°12'33.34"N, 06°35'54.78"W), 28.V.2021, Barber trap; 3 specs., 02.VII.2021, Barber trap; 11 specs., 05.VIII.2021, Barber trap; 3 specs., 20.IV.2022, Barber trap; 1 spec., 13.V.2022, Barber trap.

**Distribution.** Morocco (Löbl et al., 2008a).

***Pimelia capillata* Solier, 1836 [S]**

**Material examined.** 1 spec., Canton E (34°09'36.1"N, 06°07'04.34"W), 13.V.2022, Barber trap; 1 spec., 02.VI.2022, Barber trap.

**Distribution.** North Africa (Morocco et Algeria Löbl et al., 2008a).

***Pimelia platynota* Fairmaire, 1875 [E] [S]**

**Material examined.** 1 spec., Canton E (34°09'36.1"N, 06°07'04.34"W), 20.IV.2022, Barber trap; 1 spec., 13.V.2022, Barber trap; 2 specs., 02.VI.2022, Barber trap. 31 specs., Canton C (34°04'57.73"N, 06°24'38.31"W), 20.IV.2022, Barber trap; 2 specs., 22.VII.2022, Barber trap. 1 spec., Canton A (34°12'33.34"N, 06°35'54.78"W), 20.IV.2022, Barber trap; 1 spec., 22.VII.2022, Barber trap.

**Distribution.** Morocco (Löbl et al., 2008a).

***Pimelia chrysomeloides subris* Koch, 1941 [E] [S] (Fig. 4N)**

**Material examined.** 60 specs., Canton C (34°04'57.73"N, 06°24'38.31"W), 19.III.2021, Barber trap; 39 specs., 06.IV.2021, Barber trap; 32 specs., 30.IV.2021, Barber trap; 17 specs., 28.V.2021, Barber trap; 3 specs., 29.X.2021, Barber trap; 4 specs., 20.IV.2022, Barber trap; 1 spec., 13.V.2022, Barber trap; 2 specs., 22.VII.2022, Barber trap. 2 specs., Canton A (34°12'33.34"N, 06°35'54.78"W), 06.IV.2021, Barber trap; 1

spec., 28.V.2021, Barber trap; 3 specs., 05.VIII.2022, Barber trap; 2 specs., 16.IX.2021, Barber trap. 1 spec., Canton E (34°09'36.1"N, 06°07'04.34"W), 13.V.2022, Barber trap; 1 spec., 02.VI.2022, Barber trap.

**Distribution.** Morocco (Löbl et al., 2008a; Mas-Peinado et al., 2021).

### *Scaurus gigas* Waltl, 1835 [S]

**Material examined.** 2 specs., Canton C (34°04'57.73"N, 06°24'38.31"W), 02.VII.2021, Barber trap; 1 spec., 29.X.2021, Barber trap; 6 specs., 20.IV.2022, Barber trap; 1 spec., 13.V.2022, Barber trap; 4 specs., 02.VI.2022, Barber trap; 2 specs., 22.VII.2022, Barber trap. 1 spec., Canton A (34°12'33.34"N, 06°35'54.78"W), 05.VIII.2021, Barber trap; 1 spec., 16.IX.2021, Barber trap; 1 spec., 22.VII.2022, Barber trap. 1 spec., Canton E (34°09'36.1"N, 06°07'04.34"W), 20.IV.2022, Barber trap; 1 spec., 13.V.2022, Barber trap.

**Distribution.** Western Palaearctic (Spain, Portugal, Algeria, and Morocco – Löbl et al., 2008b).

### *Sepidium aliferum* Erichson, 1841 [S]

**Material examined.** 10 specs., Canton C (34°04'57.73"N, 06°24'38.31"W), 19.III.2021, Barber trap; 12 specs., 06.IV.2021, Barber trap; 3 specs., 30.IV.2021, Barber trap; 5 specs., 20.IV.2022, Barber trap. 12 specs., Canton A (34°12'33.34"N, 06°35'54.78"W), 20.IV.2022, Barber trap.

**Distribution.** Western Palaearctic (Spain, Algeria, Tunisia, and Morocco – Löbl et al., 2008a).

### *Stenosis mamorensis* Théry, 1924 [E] [S]

**Material examined.** 1 spec., Canton C (34°04'57.73"N, 06°24'38.31"W), 30.IV.2021, Barber trap. 1 spec., Canton A (34°12'33.34"N, 06°35'54.78"W), 06.IV.2021, Barber trap; 2 specs., 28.V.2021, Barber trap; 3 specs., 02.VII.2021, Barber trap; 2 specs., 05.VIII.2021, Barber trap; 3 specs., 16.IX.2021, Barber trap; 3 specs., 20.IV.2022, Barber trap; 1 spec., 02.VI.2022, Barber trap; 1 spec., 23.VI.2022, Barber trap. 2 specs., Canton E (34°09'36.1"N, 06°07'04.34"W), 22.VII.2022, Barber trap.

**Distribution.** Morocco (Löbl et al., 2008a).

### *Zophosis minuta* (Fabricius, 1775) [S]

**Material examined.** 3 specs., Canton C (34°04'57.73"N, 06°24'38.31"W), 06.IV.2021, Barber trap; 17 specs., 30.IV.2021, Barber trap; 13 specs., 28.V.2021, Barber trap; 1 spec., 05.VIII.2021, Barber trap; 4 specs., 19.IX.2021, Barber trap; 57 specs., 20.IV.2022, Barber trap; 55 specs., 13.V.2022, Barber trap; 28 specs., 02.VI.2022, Barber trap; 4 specs., 23.VI.2022, Barber trap. 4 specs., Canton A (34°12'33.34"N, 06°35'54.78"W), 28.V.2021, Barber trap; 4 specs., 02.VII.2021, Barber trap; 18 specs., 05.VIII.2021, Barber trap; 33 specs., 20.IV.2022, Barber trap; 27 specs., 13.V.2022, Barber trap; 21 specs., 02.VI.2022, Barber trap; 4 specs., 23.VI.2022, Barber trap; 3 specs., 22.VII.2022, Barber trap. 1 spec., Canton E (34°09'36.1"N, 06°07'04.34"W), 20.IV.2022, Barber trap; 3 specs., 13.V.2022, Barber trap; 2 specs., 02.VI.2022, Barber trap; 2 specs., 23.VI.2022, Barber trap; 1 spec., 22.VII.2022, Barber trap.

**Distribution.** Western Palaearctic (Portugal, Spain, Morocco and Tunisia – Lefèvre, 1885; Löbl et al., 2008a; López-Pérez, 2016).

## Family Trogidae MacLeay, 1819

### *Trox fabricii* Reiche, 1853 [S]

**Material examined.** 2 specs., Canton C (34°04'57.73"N, 06°24'38.31"W), 19.III.2021, Barber trap; 3 specs., 06.IV.2021, Barber trap; 2 specs., 19.IX.2021, Barber trap; 12 specs., 29.X.2021, Barber trap.

**Distribution.** Western Palaearctic (Spain, Portugal, Italy, and North Africa – Pittino, 2006).

## Family Trogossitidae Latreille, 1802

### *Tenebroides mauritanicus* (Linnaeus, 1758) [S]

**Material examined.** 1 spec., Canton A (34°12'33.34"N, 06°35'54.78"W), 29.X.2021, OCT.

**Distribution.** Palaearctic (Europe, Asia, and North Africa – Kolibac, 2007).

**Table 1.** List of species collected in the Maamora forest at sites S1, S2 and S3, and their status: [S] Saproxylic species; [R] Saproxylic species listed in the Red List of the Mediterranean region; [N] Species newly recorded for Morocco and [E] Endemic species.

Families	Species	Status	S1	S2	S3	
<b>Anobiidae</b>	<i>Stegobium paniceum</i> (Linnaeus, 1758)	S	*	*		
	<i>Xyletinus bucephalus</i> (Illiger, 1807)	S	*			
<b>Anthicidae</b>	<i>Notoxus monoceros</i> (Linnaeus, 1761)	N, S	*	*		
	<i>Notoxus trifasciatus</i> Rossi, 1792	S	*	*	*	
	<i>Stricticollis transversalis</i> (A. Villa & J.B. Villa, 1833)	S	*	*		
<b>Anthribidae</b>	<i>Bruchela rufipes rufipes</i> (Olivier, 1790)		*			
<b>Brentidae</b>	<i>Kalcapion semivittatum</i> (Gyllenhal, 1833)			*		
	<i>Ceratapion robusticorne</i> (Desbrochers des Loges, 1866)			*		
<b>Buprestidae</b>	<i>Acmaeodera bipunctata kureimatica</i> Escalera, 1914				*	
	<i>Acmaeoderella adpersula</i> (Illiger, 1803)	S	*	*	*	
	<i>Acmaeoderella discoidea</i> (Fabricius, 1787)		*	*	*	
	<i>Acmaeoderella lanuginosa</i> (Gyllenhal, 1817)		*	*	*	
	<i>Agrilus biguttatus</i> (Fabricius, 1777)	S	*			
	<i>Agrilus graminis mamorensis</i> Théry, 1930	E, S	*	*		
	<i>Anthaxia anatolica ferulae</i> Gené, 1839			*		
	<i>Anthaxia millefolii polychloros</i> Abeille de Perrin, 1894		*		*	
	<i>Anthaxia nitidula</i> (Linnaeus, 1758)	N, S			*	
	<i>Anthaxia umbellatarum umbellatarum</i> (Fabricius, 1787)		*	*	*	
	<i>Habroloma triangulare</i> (Lacordaire, 1835)			*		
	<i>Perotis unicolor igniventris</i> Escalera, 1914	E, S		*	*	
<b>Cantharidae</b>	<i>Malthinus flaveolus</i> (Herbst, 1786)	N		*		
<b>Carabidae</b>	<i>Acinopus picipes</i> (Olivier, 1795)		*		*	
	<i>Agonum numidicum</i> (Lucas, 1846)		*			
	<i>Amara aenea</i> (De Geer, 1774)		*			
	<i>Anisodactylus binotatus</i> (Fabricius, 1787)		*			
	<i>Calathus circumseptus</i> Germar, 1824		*			
	<i>Calathus fuscipes algericus</i> Gautier des Cottés, 1866			*		
	<i>Calathus mollis atticus</i> Gautier des Cottés, 1867		*			
	<i>Carabus faminii favieri</i> Fairmaire, 1859	E	*	*	*	
	<i>Carabus rugosus rugosus</i> Fabricius, 1792	E		*		
	<i>Cicindela maroccana</i> Fabricius, 1801		*	*	*	
	<i>Dromius agilis</i> (Fabricius, 1787)	N			*	
	<i>Graphipterus exclamationis</i> (Fabricius, 1792)					
	<i>Harpalus attenuatus</i> Stephens, 1828		*			
	<i>Harpalus serripes serripes</i> (Quensel, 1806)		*			
	<i>Microlestes abeillei brisouti</i> Holdhaus, 1912		*	*		
	<i>Ocys tachysoides</i> (Antoine, 1933)			*		
	<i>Paradromius linearis linearis</i> (Olivier, 1795)			*		
	<i>Philorhizus quadrisignatus</i> (Dejean, 1825)	N		*		
	<i>Poecilus baeticus gharbensis</i> (Alluaud, 1927)	E	*			
	<i>Pterostichus elongatus</i> (Duftschmid, 1812)		*	*	*	
	<i>Pterostichus globosus pecoudi</i> (Antoine, 1922)	E		*		
	<i>Scarites terricola terricola</i> Bonelli, 1813		*	*	*	
	<i>Sphodrus leucophthalmus</i> (Linnaeus, 1758)		*			
	<i>Syntomus foveatus</i> (Geoffroy, 1785)		*			
	<i>Trymosternus colombati</i> Antoine, 1934	E		*		
	<b>Cerambycidae</b>	<i>Nathrius brevipennis</i> (Mulsant, 1839)	S, R	*		
		<i>Stenopterus ater</i> (Linnaeus, 1767)	S, R	*	*	
<i>Stictoleptura tangeriana</i> (Tournier, 1875)		E, S			*	
<i>Trichoferus ilicis</i> Sama, 1987		E, S, R	*	*		
<b>Chrysomelidae</b>	<i>Bruchidius biguttatus</i> (Olivier, 1800)		*		*	
	<i>Bruchidius cisti</i> (Fabricius, 1775)	N	*			
	<i>Bruchidius rubiginosus</i> (Desbrochers des Loges, 1869)		*		*	
	<i>Cassida flaveola</i> Thunberg, 1794	N	*			

Families	Species	Status	S1	S2	S3
	<i>Cassida vittata</i> Villers, 1789				*
	<i>Chrysolina americana</i> (Linnaeus, 1758)			*	
	<i>Chrysolina kocheri</i> (Codina Padilla, 1961)	E	*		
	<i>Cryptocephalus rabatensis</i> Pic, 1953	E	*	*	
	<i>Cryptocephalus rugicollis</i> Olivier, 1792		*	*	
<b>Cleridae</b>	<i>Opilo domesticus</i> (Sturm, 1837)	S, R	*	*	
<b>Coccinellidae</b>	<i>Adalia decempunctata</i> (Linnaeus, 1758)		*	*	*
	<i>Coccinella algerica</i> Kovář, 1977		*		*
	<i>Hyperaspis duvergeri</i> Fürsch, 1985		*		
	<i>Oenopia lyncea lyncea</i> (Olivier, 1808)			*	
	<i>Oenopia conglobata</i> (Linnaeus, 1758)		*		
	<i>Platynaspis luteorubra</i> (Goeze, 1777)		*	*	
	<i>Rhyzobius chrysomeloides</i> (Herbst, 1792)				*
	<i>Rhyzobius litura</i> (Fabricius, 1787)			*	
	<i>Rodolia cardinalis</i> (Mulsant, 1850)		*	*	
	<i>Scymnus subvillosus</i> (Goeze, 1777)			*	
<b>Corylophidae</b>	<i>Orthoperus anxius</i> Mulsant & Rey, 1861		*		
<b>Cryptophagidae</b>	<i>Cryptophagus dentatus</i> (Herbst, 1793)		*	*	
<b>Curculionidae</b>	<i>Brachycerus barbarus</i> (Linnaeus, 1758)			*	
	<i>Brachycerus muricatus</i> Olivier, 1790			*	
	<i>Brachyderes pubescens</i> Boheman, 1833			*	
	<i>Brachypera zoilus</i> (Scopoli, 1763)		*		
	<i>Charagmus griseus</i> (Fabricius, 1775)				*
	<i>Ceutorynchus contractus</i> (Marsham, 1802)			*	
	<i>Coeliodes ilicis</i> (Bedel, 1885)		*	*	
	<i>Coeliodes ruber</i> (Marsham, 1802)			*	
	<i>Gronops lunatus</i> (Fabricius, 1775)		*		
	<i>Liophloeus tessulatus</i> (O. F. Müller, 1776)	N	*		
	<i>Listroderes costirostris</i> Schoenherr, 1826		*		
	<i>Maurobaris spoliata</i> (Boheman, 1836)		*		
	<i>Mecinus pascuorum</i> (Gyllenhal, 1813)	N	*		*
	<i>Mecinus pyraster</i> (Herbst, 1795)		*		
	<i>Micrelus ferrugatus</i> (Perris, 1847)		*		
	<i>Microplontus molitor</i> (Gyllenhal, 1837)		*		
	<i>Orchestes erythropus</i> (Germar, 1821)			*	
	<i>Orchestes irroratus maroccanus</i> (Roudier, 1954)			*	
	<i>Orchestes rusci</i> (Herbst, 1795)	N	*		
	<i>Otiorhynchus cribricollis</i> Gyllenhal, 1834				*
<b>Dasytidae</b>	<i>Dasytes nigroaeneus</i> Küster, 1850		*	*	*
	<i>Dasytes terminalis</i> Jacquelin du Val, 1863		*	*	*
<b>Dermestidae</b>	<i>Anthrenus angustefasciatus</i> Ganglbauer, 1904		*	*	*
	<i>Anthrenus museorum</i> (Linnaeus, 1761)		*	*	*
	<i>Anthrenus pimpinellae</i> (Fabricius, 1775)		*	*	*
	<i>Attagenus smirnovi</i> Zhantiev, 1973		*	*	*
	<i>Attagenus trifasciatus</i> (Fabricius, 1787)		*	*	*
	<i>Attagenus unicolor unicolor</i> (Brahm, 1791)		*	*	
	<i>Dermestes frichii</i> Kugelann, 1792		*	*	*
	<i>Dermestes maculatus</i> De Geer, 1774		*		
	<i>Dermestes peruvianus</i> Laporte de Castelnau, 1840		*		
	<i>Thorictus castaneus castaneus</i> Germar, 1834		*		
<b>Elateridae</b>	<i>Ampedus balteatus</i> (Linnaeus, 1758)	N, S, R		*	
	<i>Agriotes sordidus</i> (Illiger, 1807)	S	*		
	<i>Cardiophorus rufipes</i> (Goeze, 1777)	S	*	*	
	<i>Cebrio maculicollis</i> Fairmaire, 1856	E, S	*		
	<i>Drasterius bimaculatus</i> (Rossi, 1790)	S	*		
	<i>Lacon punctatus punctatus</i> (Herbst, 1779)	S, R	*		
<b>Geotrupidae</b>	<i>Thorectes distinctus</i> Marseul, 1878		*	*	*
	<i>Typhaeus typhoeus</i> (Linnaeus, 1758)		*		*
<b>Glaphyridae</b>	<i>Anthypna meles</i> (Fabricius, 1792)			*	*

Families	Species	Status	S1	S2	S3
<b>Histeridae</b>	<i>Eulasia goudoti</i> (Laporte, 1840)	E	*	*	*
	<i>Hister moerens</i> Erichson, 1834			*	
	<i>Hister maroccanus</i> Schmidt, 1887	E			*
	<i>Hypocaccus brasiliensis</i> (Paykull, 1811)		*	*	
	<i>Hypocaccus rugiceps</i> (Duftschmid, 1805)	N		*	*
	<i>Pactolinus major</i> (Linnaeus, 1767)			*	
	<i>Saprinus acuminatus</i> (Fabricius, 1798)		*	*	*
	<i>Saprinus chalcites</i> (Illiger, 1807)		*	*	
	<i>Saprinus figuratus</i> Marseul, 1855		*	*	*
	<i>Saprinus proximus simillimus</i> Wollaston, 1865			*	
<b>Laemophloeidae</b>	<i>Cryptolestes ferrugineus</i> (Stephens, 1831)		*	*	*
	<i>Laemophloeus monilis</i> (Fabricius, 1787)	S		*	
<b>Latridiidae</b>	<i>Corticarina cavicollis</i> (Mannerheim, 1844)	N		*	
	<i>Corticarina curta</i> (Wollaston, 1854)		*	*	*
<b>Leiodidae</b>	<i>Agathidium marocanum</i> Hlisnikovsky, 1968	E, S			*
	<i>Catops coracinus</i> Kellner, 1846			*	*
	<i>Ptomaphagus tenuicornis mauritanicus</i> Jeannel, 1934			*	
<b>Meloidae</b>	<i>Actenodia distincta</i> (Chevrolat, 1840)				*
	<i>Cerocoma schaefferi</i> (Linnaeus, 1758)	N			*
	<i>Hycleus duodecimpunctatus</i> (Olivier, 1811)	N	*		*
	<i>Hycleus rufipalpis</i> (Escalera, 1909)		*	*	*
<b>Melolonthidae</b>	<i>Euserica mamorensis</i> Baraud, 1965	E	*	*	*
	<i>Hoplia argentea</i> (Poda, 1761)	N	*		
	<i>Hoplia philanthus gagates</i> (Bedel, 1911)		*	*	
	<i>Sphodroxia maroccana</i> Ley, 1923	E	*	*	*
<b>Melyridae</b>	<i>Charopus rotundatus</i> Erichson, 1840			*	
	<i>Colotes javeti</i> Jacquelin du Val, 1853			*	
	<i>Colotes punctatus</i> (Erichson, 1840)		*		*
<b>Mordellidae</b>	<i>Mediimorda bipunctata</i> (Germar, 1827)		*	*	*
	<i>Mordella aculeata</i> Linnaeus, 1758	N	*	*	*
	<i>Variimorda ragusai</i> (Emery, 1876)	S			*
<b>Nitidulidae</b>	<i>Carpophilus hemipterus</i> (Linnaeus, 1758)		*		
	<i>Epuraea aestiva</i> (Linnaeus, 1758)	N, S		*	
	<i>Epuraea unicolor</i> (Olivier, 1790)			*	
	<i>Brassicogethes aeneus</i> (Fabricius, 1775)		*	*	*
<b>Oedemeridae</b>	<i>Oedemera barbara</i> (Fabricius, 1792)	S		*	*
	<i>Oedemera simplex</i> (Linnaeus, 1767)	S		*	
<b>Ptinidae</b>	<i>Dignomus dilophus</i> (Illiger, 1807)	S	*	*	*
	<i>Dignomus gibbicollis</i> (P. H. Lucas, 1846)	S	*	*	
	<i>Mesocoelopus niger</i> (P. W. J. Müller, 1821)	S	*		
	<i>Mizodorcatoma dommeri</i> (Rosenhauer, 1856)	S			*
	<i>Ptinus bidens</i> Olivier, 1790	N		*	
	<i>Ptinus spitzyi</i> A. Villa & G. B. Villa, 1838		*	*	*
<b>Phalacridae</b>	<i>Olibrus affinis</i> (Sturm, 1807)			*	
	<i>Stilbus testaceus</i> (Panzer, 1796)		*	*	
<b>Scarabaeidae</b>	<i>Aethiessa floralis</i> (Fabricius, 1787)		*	*	*
	<i>Otophorus haemorrhoidalis</i> (Linnaeus, 1758)		*		
	<i>Labarrus lividus</i> (Olivier, 1789)		*		
	<i>Esymus merdarius</i> (Fabricius, 1775)		*		
	<i>Subrinus sturmi</i> (Harold, 1870)				*
	<i>Onitis alexis septentrionalis</i> Balthasar, 1942		*		
	<i>Onthophagus maki</i> (Illiger, 1803)		*		
	<i>Onthophagus similis</i> (Scriba, 1790)		*		
	<i>Onthophagus vacca</i> (Linnaeus, 1767)		*	*	*
	<i>Oryctes nasicornis grypus</i> (Illiger 1803)				*
	<i>Oxythyrea funesta</i> (Poda, 1761)	S	*	*	*
	<i>Phyllopertha horticola</i> (Linnaeus, 1758)	N	*		
	<i>Protaetia (Netocia) morio heyrovskyi</i> (Balthasar, 1935)		*	*	
	<i>Scarabaeus cicatricosus</i> Lucas, 1846			*	*

Families	Species	Status	S1	S2	S3
	<i>Scarabaeus sacer</i> Linnaeus, 1758		*	*	
	<i>Tropinota squalida pilosa</i> Brullé, 1832		*	*	*
	<i>Tropinota hirta</i> (Poda, 1761)			*	
Scolytidae	<i>Dryocoetes autographus</i> (Ratzeburg, 1837)	N, S	*	*	*
	<i>Ernoporicus fagi</i> (Fabricius, 1798) (= <i>Eidophelus fagi</i> )	N	*	*	*
	<i>Hypoborus ficus</i> Erichson, 1836			*	
	<i>Platypus cylindrus</i> (Fabricius, 1792)		*		
	<i>Xyleborus monographus</i> (Fabricius, 1792)	S			*
Scaptiidae	<i>Scaptia ophthalmica</i> Mulsant, 1856 ( <i>S. schotti</i> Leblanc, 2012)	S	*	*	*
	<i>Anaspis flava</i> (Linnaeus, 1758)	N, S		*	
Scydmaenidae	<i>Scydmaenus tingitanus</i> (Franz, 1962)	E, S		*	
Silphidae	<i>Silpha tristis</i> Illiger, 1798		*		*
	<i>Silpha olivieri</i> Bedel, 1887		*		
	<i>Thanatophilus sinuatus</i> (Fabricius, 1775)			*	
Staphylinidae	<i>Bolitobius cingulatus</i> (Mannerheim, 1830)	N	*		
	<i>Eusphalerum torquatum torquatum</i> (Marsham, 1802)			*	
	<i>Gabrius appendiculatus</i> Sharp, 1910	N	*	*	*
	<i>Micropeplus staphylinoides</i> Marsham, 1802			*	
	<i>Ontholestes marginalis</i> (Gené, 1836)		*		
	<i>Oxytelus sculptus</i> Gravenhorst, 1806	S	*	*	*
	<i>Xantholinus linearis</i> (Olivier, 1795)		*	*	
Tenebrionidae	<i>Adelostoma sulcatum</i> Duponchel, 1827			*	*
	<i>Akis tingitana</i> Lucas, 1859	E	*	*	
	<i>Alphasida</i> ( <i>Glabrasida</i> ) <i>iblanensis reymondi</i> Antoine, 1952	E		*	*
	<i>Arthrodeis globulosus</i> Escalera, 1922	E, S	*	*	
	<i>Blaps ovipennis</i> Seidlitz, 1893	E, S		*	
	<i>Cheirodes brevicollis</i> (Wollaston, 1864)	S	*		
	<i>Cheirodes sardous sardous</i> Gené, 1839	S		*	
	<i>Erodium granipennis</i> Fairmaire, 1871	S	*	*	*
	<i>Heliotaurus ruficollis tangerianus</i> Escalera, 1922	E	*		
	<i>Isomira melanophthalma</i> (Lucas, 1846)			*	*
	<i>Latheticus oryzae</i> Waterhouse, 1880			*	
	<i>Misolampus goudotii</i> Guérin-Méneville, 1834	S			*
	<i>Pachychila alluaudi</i> Peyerimhoff, 1925	E, S	*		
	<i>Pachychila punctata mamorensis</i> Antoine, 1942	E, S	*	*	*
	<i>Pimelia capillata</i> Solier, 1836	S			*
	<i>Pimelia platynota</i> Fairmaire, 1875	E, S	*	*	*
	<i>Pimelia chrysomeloides subbris</i> Koch, 1941	E, S	*	*	*
	<i>Scaurus gigas</i> Walzl, 1835	S	*	*	*
	<i>Sepidium aliferum</i> Erichson, 1841	S	*	*	
	<i>Stenosis mamorensis</i> Théry, 1924	E, S	*		*
<i>Zophosis minuta</i> (Fabricius, 1775)	S	*	*	*	
Trogidae	<i>Trox fabricii</i> Reiche, 1853	S		*	
Trogossitidae	<i>Tenebroides mauritanicus</i> (Linnaeus, 1758)	S		*	

**Table 2.** Number of species according to sampling sites: **S1** (warm sub-humid bioclimate with maritime influences; open cork oak forest); **S2** (warm sub-humid bioclimate with maritime influences; old cork oak trees with dense understory); and **S3** (semiarid bioclimate with temperate winters, a mixed cork oak forest with eucalyptus trees).

Species	Stations		
	S1	S2	S3
Saproxyllic	34	36	24
Endemic	18	18	13
new for Morocco	12	11	10
Total	139	131	86

## DISCUSSION

The beetle fauna of the cork oak forest of Maamora constitutes an essential link for the functioning and dynamics of this forest ecosystem, with a significant richness and diversity of the beetle population. 256 species belonging to 42 families were identified, most of them belonging to the Tenebrionidae and Carabidae families, each with 27 species, and Curculionidae with 24 species. Previous studies on the entomofauna of the Maamora forest have reported the presence of 104 beetle species (El Alami Idrissi, 2013; Villemant & Fraval, 1993; Idrissi, 1982). These results complete the studies of previous authors and provide useful information on the differentiated methodology for collecting these beetles (Table 3).

**Table 3.** Number of species according to sampling methods: pitfall traps, coloured bowls (yellow, orange, white, and blue), window traps and visual collection.

Families	Number of species			
	Pitfall traps	Coloured bowls	Window traps	Visual collection
Anobiidae		1	1	
Anthricidae	1	2		
Anthribidae	1			
Brentidae		2		
Buprestidae		11		1
Cantharidae		1		
Carabidae	19	2	3	
Cerambycidae		2	2	
Chrysomileidae		6	2	1
Cleridae			1	
Coccinilidae		9	1	
Corylophidae		1		
Cryptophagidae			1	
Curculionidae	12	8		
Dasytidae		2		
Dermetidae	2	7		1
Elateridae	1	3	1	1
Geotrupidae	2			
Glaphyridae		2		
Histeridae	4	4	1	
Laemophloeidae			2	
Latridiidae		1	1	
Leiodidae	2			
Meloidae		4		
Melolonthidae	1	2	1	
Melyridae	1	2		
Mordellidae		2	1	
Nitidulidae		3	1	
Oedemeridae		2		
Ptinidae	3	1	2	
Phalacridae		2		
Scarabaeidae	6	9	1	1
Scolytidae		2	3	
Scaptidae		2		
Scydmaenidae	1			
Silphidae	3			
Staphylinidae	4	2	1	
Tenebrionidae	17	3	1	
Trogidae	1			
Trogossitidae		1		
<b>Total species</b>	<b>81</b>	<b>101</b>	<b>27</b>	<b>5</b>

The pitfall traps collected 81 species. The colored bowls (yellow, orange, white and blue) contributed to the collection of 101 species. The window traps collected 27 species, and 5 species were collected visually from the field. Out of the sampled species, 53 were saproxylic, including 6 species considered rare according to the Red List of Saproxylic Beetles in the Mediterranean region: *Nathrius brevipennis*, *Stenopterus ater*, *Trichoferus ilicis*, *Opilo domesticus*, *Ampedus balteatus*, and *Lacon punctatus punctatus* (Nieto & Alexander, 2010; García et al., 2018). The results show a much lower number of species at site S3, compared with the two other sites (Table 2). The analysis of certain taxonomic groups, particularly saproxylic beetle assemblages, makes it possible to monitor trends in forest environments (Rossi & Vallauri, 2013). Saproxylic beetles are highly sensitive to changes and disturbances (such as fires, collection of dead wood, overgrazing, etc.), and their qualitative (rarity) and quantitative study can provide information on the degradation and resilience of the environment (Blandin, 1986; Speight, 1989).

#### AUTHOR'S CONTRIBUTION

The authors confirm their contribution to the paper as follows: All authors contributed to the study conception and design. H. Habbaz and N. Maatouf: compiled the species records and background data, taxonomic revisions of the species names, and drafted the original manuscript; J-P. Lumaret: revised the taxonomy, contributed to the analysis, and reviewed the literature; L. Rohi: drafted and revised the manuscript. The authors read and approved the final version of the manuscript.

#### FUNDING

This research received no specific grant from any funding agencies.

#### AVAILABILITY OF DATA AND MATERIAL

The specimens listed in this study are deposited in the insect collection of the Centre for Innovation, Research, and Training - National Agency for Water and Forests, BP. 763, Rabat - Agdal, Morocco, and are available from the curator, upon request.

#### ETHICS APPROVAL AND CONSENT TO PARTICIPATE

This study only included arthropod material, and all required ethical guidelines for the treatment and use of animals were strictly adhered to in accordance with international, national, and institutional regulations. No human participants were involved in any studies conducted by the authors for this article.

#### CONSENT FOR PUBLICATION

Not applicable.

#### CONFLICT OF INTERESTS

The authors declare that there is no conflict of interest regarding the publication of this paper.

#### ACKNOWLEDGMENTS

We would like to thank the Director of the Centre for Innovation, Research, and Training in Rabat (National Agency for Water and Forests) for his hospitality and support in carrying out this work. Thanks to Vikki Bengtsson (BSc Env. Biology, Pro Natura Foundation, Gothenburg, Sweden) for reviewing the manuscript. We are also grateful to Amine Samih (Doctoral student in the field of entomology - Hassan II University) and El Ayyachi El Mnouar (Technician in the field of forest management - National Agency for Water and Forests Rabat, Morocco) for their guidance, assistance, and contribution in collecting the field data. Special thanks to Dr. Piero Leo (Private entomologist, Cagliari, Italy) for his invaluable identification of some specimens. We thank also the reviewers for their constructive comments.

## REFERENCES

- Aafi, A. (2007) *Etude de la Diversité Floristique de l'Écosystème de Chêne-liège de la Forêt de la Mamora* (Unpublished Ph.D. Dissertation). Institut Agronomique et Vétérinaire Hassan II, Rabat. 190 p.
- Aafi, A., Achhal El Kadmiri, A., Benabide, A. & Rochdi, M. (2005) Richesse et biodiversité floristique de la subéraie de la Mamora. *Acta Botanica Malacitana*, 30, 127–138. <https://doi.org/10.24310/abm.v30i0.7187>
- AEFCS (Administration des Eaux et Forêts et de la Conservation des Sols) (1996) *Plan Directeur des Aires Protégées du Maroc*. Groupement BCEOMSECA, Montpellier France. Vol. 2: les Sites d'Intérêt Biologique et Écologique du Domaine Continental, Rabat. 410 p.
- Ali, E., Cramer, W., Carnicer, J., Georgopoulou, E., Hilmi, N.J.M., Le Cozannet, G. & Lionello, P. (2022) Cross-Chapter Paper 4: Mediterranean Region. In: Pörtner, H.-O., Roberts, D.C., Tignor, M., Poloczanska, E.S., Mintenbeck, K., Alegria, A., Craig, M., Langsdorf, S., Löschke, S., Möller, V., Okem, A. & Rama, B. (eds) *Climate Change 2022: Impacts, Adaptation and Vulnerability. Contribution of Working Group II to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge University Press, Cambridge, UK and New York, NY, USA, pp. 2233–2272. <https://doi.org/10.1017/9781009325844.021>
- Alonso Zaragaza, M.A. (2011) Anthribidae. In: Löbl, I. & Smetana, A. (eds) *Catalogue of Palaearctic Coleoptera*, Vol. 7. Apollo Books, Stenstrup, Denmark, pp. 90–109.
- Anton, K.W. (2010) Bruchinae. In: Löbl, I. & Smetana, A. (eds) *Catalogue of Palaearctic Coleoptera*, Vol. 6. Apollo Books, Stenstrup, Denmark, pp 339–353.
- Arahou, M. (2008) *Catalogue de l'entomofaune du Chêne vert du Moyen Atlas (Maroc)*. Documents de l'Institut Scientifique, Rabat, n° 22. Université Mohammed V - Agdal, Institut Scientifique, Rabat. 36 p.
- Balkenohl, M. (2003) Carabidae, Tribe: Scaritini. In: Löbl, I. & Smetana, A. (eds) *Catalogue of Palaearctic Coleoptera*, Vol. 1. Apollo Books, Stenstrup, pp. 231–234.
- Belghazi, B. & Mounir, F. (2016) *Analyse de la Vulnérabilité au Changement Climatique du Couvert Forestier. Forêt de la Maâmora (Maroc)*. FAO, Rapport Technique. Rabat. 124 p.
- Benabid, A. (2000) *Flore et Écosystèmes du Maroc: Evaluation et Préservation de la Biodiversité*. Ibis Press, Paris. 359 p.
- Bensusan, K.J., Muñoz Batet, J. & Perez, C.E. (2006) *Coccinella algerica* Kovář, 1977: a new species to the fauna of mainland Europe, and a key to the *Coccinella* Linnaeus, 1758 of Iberia, the Maghreb and the Canary Islands (Coleoptera, Coccinellidae). *Boletín Sociedad Entomológica Aragonesa*, 39, 323–327.
- Benyahia, Y. (2016) *Inventaire de la Biodiversité entomologique (Coléoptères et Hyménoptères) pour une gestion patrimoniale de sapinière du Parc National de Talassemtane (Rif, Maroc)*. (Unpublished Ph.D. Dissertation), Faculté des Sciences Ben M'sik-Casablanca, Morocco. 163 p.
- Benyahia, Y., Latifa, R., Valladarès, L., Maatouf, N., Courtin, O., El Antry, S. & Brustel, H. (2016) Preliminary list of Coleoptera heritage species of the Talassemtane National Park, Morocco. *Journal of Insect Biodiversity*, 4 (13), 1–30. <https://doi.org/10.12976/jib/2015.4.13>
- Bezdek, A. (2006a) Sericinae. In: Löbl, I. & Smetana, A. (eds) *Catalogue of Palaearctic Coleoptera*, Vol. 3. Apollo Books, Stenstrup, pp. 228–248.
- Bezdek, A. (2006b) Melolonthidae: Tribe: Melolonthini. In: Löbl, I. & Smetana, A. (eds) *Catalogue of Palaearctic Coleoptera*, Vol. 3. Apollo Books, Stenstrup, pp. 191–198.
- Bezdek, A. & Krell, F.T. (2006) Tribe: Onitini. In: Löbl, I. & Smetana, A. (eds) *Catalogue of Palaearctic Coleoptera*, Vol. 3. Apollo Books, Stenstrup, pp. 158–159.
- Billy, S. (2006) Buprestidae, Buprestinae: Trib. Anthaxiini. In: Löbl, I. & Smetana, A. (eds) *Catalogue of Palaearctic Coleoptera*, Vol. 3. Apollo Books, Stenstrup, pp. 369–381.
- Blandin, P. (1986) Bioindicateurs et diagnostic des systèmes écologiques. *Bulletin d'Ecologie*, 17 (4), 215–307.
- Bologna, M.A. (2008) Meloidae. In: Löbl, I. & Smetana, A. (eds) *Catalogue of Palaearctic Coleoptera*, Vol. 5. Apollo Books, Stenstrup, pp. 370–412.
- Borowski, J. (2007) Ptinidae. In: Löbl, I. & Smetana, A. (eds) *Catalogue of Palaearctic Coleoptera*, Vol. 4. Apollo Books, Stenstrup, pp. 328–362.
- Borowiec, L. & Sekerka, L. (2010) Cassidinae. In: Löbl, I. & Smetana, A. (eds) *Catalogue of Palaearctic Coleoptera*, Vol. 6. Apollo Books, Stenstrup, pp. 368–390.
- Bouget, C. & Noblecourt, T. (2005) Short-term development of ambrosia and bark beetle assemblages following a windstorm in French broadleaved temperate forests. *Journal of Applied Entomology*, 129 (6), 300–310. <https://doi.org/10.1111/j.1439-0418.2005.00970.x>

- Bouget, C., Brustel, H. & Nageleisen, L. M. (2005) Nomenclature des groupes écologiques d'insectes liés au bois: synthèse et mise au point sémantique. *Comptes Rendus Biologies*, 328 (10–11), 936–948. <https://doi.org/10.1016/j.crvi.2005.08.003>
- Bouget C., Brustel H., Brin A., Noblecourt T., (2008) Sampling saproxylic beetles with window flight traps: methodological insights. *Revue d'Ecologie (La Terre et la Vie)*, 10 (suppl.), 21–32. <https://doi.org/10.3406/revec.2008.1457>
- Bousquet, Y. (2003a) Carabidae: Tribe Platynini. In: Löbl, I. & Smetana, A. (eds) *Catalogue of Palaearctic Coleoptera*, Vol. 1. Apollo Books, Stenstrup, pp. 449–454.
- Bousquet, Y. (2003b) Carabidae: Tribe Pterostichini. In: Löbl, I. & Smetana, A. (eds) *Catalogue of Palaearctic Coleoptera*, Vol. 1. Apollo Books, Stenstrup, pp. 469–515.
- Bousquet, Y., Brezina, B., Davies, A., Farkac, J. & Smetana, A. (2003) Carabidae: Tribe Carabini. In: Löbl, I. & Smetana, A. (eds) *Catalogue of Palaearctic Coleoptera*, Vol. 1. Apollo Books, Stenstrup, pp. 118–200.
- Bowstead, S. (2007) Corylophidae. In: Löbl, I. & Smetana, A. (eds) *Catalogue of Palaearctic Coleoptera*, Vol. 4. Apollo Books, Stenstrup, pp. 631–635.
- Brin, A. & Brustel, H. (2006) Réponse des coléoptères saproxyliques à l'hétérogénéité des subéraies dans le massif des Maures (France). *Revue d'Ecologie (La Terre et la Vie)*, 61, 327–342. <https://doi.org/10.3406/revec.2006.1328>
- Brochier, S. & Micas, L. (2019) *Echantillonnage des Coléoptères Saproxyliques de la Réserve Biologique Intégrale d'Assan (France, Hautes-Alpes), Années 2013–2018*. Office National des Forêts, Réseau National d'Entomologie Forestière, France. 40 p.
- Brustel, H. (2001) *Coléoptères Saproxyliques et Valeur Biologique des Forêts Françaises: Perspectives pour la Conservation du Patrimoine Naturel* (Unpublished Ph.D. Dissertation), Institut National Polytechnique, Toulouse. 297 pp.
- Brustel, H. (2004) *Coléoptères Saproxyliques et Valeur Biologique des Forêts Françaises: perspectives pour la conservation du patrimoine naturel*. Les dossiers forestiers, n°13. Office National des Forêts, Paris. 320 p.
- Brustel, H. & Noblecourt, T. (2019) Coléoptères saproxyliques et valeur biologique des forêts françaises : perspectives pour le diagnostic et la conservation du patrimoine naturel. In: Brochier, S. & Micas, L. (eds) *Echantillonnage des Coléoptères saproxyliques de la Réserve Biologique Intégrale d'Assan (France, Hautes-Alpes), années 2013–2018*. Office National des Forêts, Réseau National d'Entomologie Forestière, pp. 2–4.
- Caldara, R. (2013) Curculioninae. In: Löbl, I. & Smetana, A. (eds) *Catalogue of Palaearctic Coleoptera*, Vol. 8. Brill, Leiden, pp. 117–172. [https://doi.org/10.1163/9789004259164\\_010](https://doi.org/10.1163/9789004259164_010)
- Cálix, M., Alexander, K.N.A., Nieto, A., Dodelin, B., Soldati, F., Telnov, D., VazquezAlbalate, X., Aleksandrowicz, O., Audisio, P., Istrate, P., Jansson, N., Legakis, A., Liberto, A., Makris, C., Merkl, O., Mugerwa Pettersson, R., Schlaghamersky, J., Bologna, M.A., Brustel, H., Buse, J., Novák, V. & Purchart, L. (2018) *European Red List of Saproxylic Beetle. Supplementary Material to the IUCN European Red List of Saproxylic Beetles*. IUCN, Brussels, Belgium. 15 p.
- Casale, A. (2003) Carabidae, Subtribe Sphodrina. In: Löbl, I. & Smetana, A. (eds) *Catalogue of Palaearctic Coleoptera*, Vol. 1. Apollo Books, Stenstrup, pp. 532–544.
- Cate, P.C. (2007) Elateridae. In: Löbl, I. & Smetana, A. (eds) *Catalogue of Palaearctic Coleoptera*, Vol. 4. Apollo Books, Stenstrup, pp. 94–209.
- Chambord, R., Chabrol, L., Brustel, H., Pantacchini, C., Plas, L., Rohr, G. & Vassel, L. (2013) *Première Liste Rouge des Coléoptères Saproxyliques et Phytophages du Limousin*. Société Entomologique du Limousin pour la DREAL Limousin. 22 p.
- Chandler, D.S., Uhmman, G., Nardi, G. & Telnov, D. (2008) Anthicidae. In: Löbl, I. & Smetana, A. (eds) *Catalogue of Palaearctic Coleoptera*, Vol. 5. Apollo Books, Stenstrup, pp. 421–455.
- Chavanon, G. (2018) *Catalogue des Coléoptères de la région Orientale du Maroc (Province de Guercif exceptée)*. *Travaux de l'Institut Scientifique, Série Zoologie*, 57, 1–193.
- Cherkaoui, S.I., Dakki, M., Selmi, S., Rguibi Idrissi, H. & Thévenot, M. (2007) Les oiseaux de la subéraie de la Maamora (Maroc): phénologie du peuplement, statut des espèces nicheuses et évolution depuis le début du vingtième siècle. *Alauda*, 75 (1), 15–32.
- Colonnelli, E. (2011) Brachyceridae. In: Löbl, I. & Smetana, A. (eds) *Catalogue of Palaearctic Coleoptera*, Vol. 7. Brill, Leiden, pp. 182–184.
- Colonnelli, E. (2013) Centorhynchinae. In: Löbl, I. & Smetana, A. (eds) *Catalogue of Palaearctic Coleoptera*, Vol. 8. Brill, Leiden, pp. 176–213.
- de Faria e Silva, I., Salgado Costas, J.M. & Serrano, A.R.M. (2013) New data on the Cholevidae (Coleoptera) from Portugal. *Graellsia*, 69 (1), 45–56. <https://doi.org/10.3989/graeellsia.2013.v69.078>

- Dellacasa, M. & Dellacasa, G. (2006) Aphodiinae: Tribe Aphodiini. In: Löbl, I. & Smetana, A. (eds) *Catalogue of Palaearctic Coleoptera*, Vol. 3. Apollo Books, Stenstrup, pp. 105–142.
- El Alami Idrissi, N. (2013) La faune de la litière de la subéraie de la Mamora. *Revue Marocaine des Sciences Agronomiques et Vétérinaires*, 1 (2), 50–57.
- El Harche, H., Chavanon, G., Dahmani, J., Fadli, M. (2022) Checklist of terrestrial beetles (Arthropoda, Insecta, Coleoptera) associated with agroecosystems in North–West Morocco. *Arxius de Miscel·lània Zoològica*, 20, 59–81. <https://doi.org/10.32800/amz.2022.20.0059>
- Galbraith, C.A. & Stroud, D.A. (2022) *Creating a Protected Area Network for Nature Recovery in England*. NECR441. Natural England. 100 p.
- García, N., Numa, C., Bartolozzi, L., Brustel, H., Buse, J., Norbiato, M., Recalde, J.I., Zapata, J.L., Dodelin, B., Alcázar, E., Barrios, V., Verdugo, A., Audisio, P., Micó, E., Otero, J. C., Bahillo, P., Viñolas, A., Valladares, L., Méndez, M., El Antry, S. & Galante, E. (2018) *The conservation status and distribution of Mediterranean saproxylic beetles*. Malaga, Spain: IUCN. xii + 58 p. <https://doi.org/10.2305/IUCN.CH.2018.RA.3.en>
- Hava, J. (2007a) Dermestidae. In: Löbl, I. & Smetana, A. (eds) *Catalogue of Palaearctic Coleoptera*, Vol. 4. Apollo Books, Stenstrup, pp. 299–302.
- Hava, J. (2007b) Dermestidae. In: Löbl, I. & Smetana, A. (eds) *Catalogue of Palaearctic Coleoptera*, Vol. 4. Apollo Books, Stenstrup, pp. 306–319.
- Hieke, F. (2003) Carabidae: subtribe, Amarina. In: Löbl, I. & Smetana, A. (eds) *Catalogue of Palaearctic Coleoptera*, Vol. 1. Apollo Books, Stenstrup, pp. 547–568.
- Horak, J. (2008) Mordellidae. In: Löbl, I. & Smetana, A. (eds) *Catalogue of Palaearctic Coleoptera*, Vol. 5. Apollo Books, Stenstrup, pp. 87–105.
- Hoekstra, J.M., Boucher, T.M., Ricketts, T.H. & Roberts, C. (2005) Confronting a biome crisis: global disparities of habitat loss and protection. *Ecology Letters*, 8, 23–29. <https://doi.org/10.1111/j.1461-0248.2004.00686.x>
- Hovorka, O. & Sciaky, R. (2003) Carabidae: Tribe Sphodrini; subtribe Calathina. In: Löbl, I. & Smetana, A. (eds) *Catalogue of Palaearctic Coleoptera*, Vol. 1. Apollo Books, Stenstrup, pp. 524–528.
- Hurka, K. (2003) Carabidae: Tribe Graphipterini. In: Löbl, I. & Smetana, A. (eds) *Catalogue of Palaearctic Coleoptera*, Vol. 1. Apollo Books, Stenstrup, pp. 359–360.
- Idrissi, L. (1982) Contribution à l'étude écologique des coléoptères sabulicoles de Sidi Boughaba (Maroc). *Bulletin de l'Institut Scientifique, Rabat*, 6, 157–178.
- Ito, N. (2003) Carabidae: subtribe Anisodactylina. In: Löbl, I. & Smetana, A. (eds), *Catalogue of Palaearctic Coleoptera*, Vol. 1. Apollo Books, Stenstrup, pp. 360–362.
- Iwan, D. & Löbl, I. (2008) Tenebrionidae: Tribe Melanimini. In: Löbl, I. & Smetana, A. (eds) *Catalogue of Palaearctic Coleoptera*, Vol. 5. Apollo Books, Stenstrup, pp. 257–258.
- Jelinek, J. & Audisio, P. (2007) Nitidulidae. In: Löbl, I. & Smetana, A. (eds) *Catalogue of Palaearctic Coleoptera*, Vol. 4. Apollo Books, Stenstrup, pp. 459–491.
- Jendek, E. (2006) Buprestidae. In: Löbl, I. & Smetana, A. (eds) *Catalogue of Palaearctic Coleoptera*, Vol. 3. Apollo Books, Stenstrup, pp. 388–421.
- Johnson, C. (2007) Latridiinae. In: Löbl, I. & Smetana, A. (eds) *Catalogue of Palaearctic Coleoptera*, Vol. 4. Apollo Books, Stenstrup, pp. 635–641.
- Johnson, C., Otero, J. C. & Leschen, R.A.B. (2007) Cryptophagidae. In: Löbl, I. & Smetana, A. (eds) *Catalogue of Palaearctic Coleoptera*, Vol. 4. Apollo Books, Stenstrup, pp. 513–531.
- Kabak, I. (2003) Carabidae: Tribe Lebiini. In: Löbl, I. & Smetana, A. (eds) *Catalogue of Palaearctic Coleoptera*, Vol. 1. Apollo Books, Stenstrup, pp. 408–438.
- Kataev, B.M., Wrase, D.W. & Ito, N. (2003) Carabidae: Subtribe Harpalina. In: Löbl, I. & Smetana, A. (eds) *Catalogue of Palaearctic Coleoptera*, Vol. 1. Apollo Books, Stenstrup, pp. 367–407.
- Kazantsev, S. & Brancuçi, M. (2007) Cantharidae. In: Löbl, I. & Smetana, A. (eds) *Catalogue of Palaearctic Coleoptera*, Vol. 4. Apollo Books, Stenstrup, pp. 234–298.
- Kippenberg, H. (2010) Chrysomelinae. In: Löbl, I. & Smetana, A. (eds) *Catalogue of Palaearctic Coleoptera*, Vol. 6. Apollo Books, Stenstrup, pp. 390–443.
- Knizek, M. (2011) Scolytinae. In: Löbl, I. & Smetana, A. (eds) *Catalogue of Palaearctic Coleoptera*, Vol. 7. Apollo Books, Stenstrup, pp. 204–250.
- Kolibac, J. (2007) Trogossitidae. In: Löbl, I. & Smetana, A. (eds) *Catalogue of Palaearctic Coleoptera*, Vol. 4. Apollo Books, Stenstrup, pp. 364–366.

- Kovar, I. (2007) Coccinellidae. In: Löbl, I. & Smetana, A. (eds) *Catalogue of Palaearctic Coleoptera*, Vol. 4. Apollo Books, Stenstrup, pp. 568–631.
- Krell, F.-T. (2006) Dynastinae. In: Löbl, I. & Smetana, A. (eds) *Catalogue of Palaearctic Coleoptera*, Vol. 3. Apollo Books, Stenstrup, pp. 277–283.
- Kubán, V. (2006) Buprestinae without Anthaxini. In: Löbl, I. & Smetana, A. (eds) *Catalogue of Palaearctic Coleoptera*, Vol. 3. Apollo Books, Stenstrup, pp. 369–387.
- Labrique, H. & Chavanon, G. (2001) Résultats d'une campagne de prospection entomologique dans le Maroc oriental (Coleoptera Tenebrionidae). *Cahiers Scientifiques du Muséum d'histoire naturelle de Lyon*, 3, 75–87. <https://doi.org/10.3406/mhnlly.2001.1323>
- Leblanc, P. (2012) Notes sur quelques Scaptia ouest-paléarctiques et description d'une espèce nouvelle (Coleoptera Scaptiidae). *Bulletin Mensuel de la Société linnéenne de Lyon*, 81 (3-4), 47–53. <https://doi.org/10.3406/linly.2012.13821>
- Leblanc, P., Levey, B. & Horak, J. (2008) Scaptiidae. In: Löbl, I. & Smetana, A. (eds) *Catalogue of Palaearctic Coleoptera*, Vol. 5. Apollo Books, Stenstrup, pp. 458–465.
- Lefèvre, M.E. (1885) *Exploration scientifique de la Tunisie publiée sous les auspices du ministère de l'Instruction publique. Sciences Naturelles, Zoologie - Coléoptères. Liste des Coléoptères recueillis en Tunisie en 1883 par M.A. Letourneux.* Imprimerie Nationale, Paris. 16 p. <https://doi.org/10.5962/bhl.title.53766>
- Löbl, I. (2007) Thorictinae. In: Löbl, I. & Smetana, A. (eds) *Catalogue of Palaearctic Coleoptera*, Vol. 4. Apollo Books, Stenstrup, pp. 302–306.
- Löbl, I., Krell, F.-T. & Kral, D. (2006a) Scarabaeidae: Tribe Scarabaeini. In: Löbl, I. & Smetana, A. (eds) *Catalogue of Palaearctic Coleoptera*, Vol. 3. Apollo Books, Stenstrup, pp. 176–177.
- Löbl, I., Krell, F.-T., Kral, D. & Ziani, S. (2006b) Scarabaeidae: Tribe Onthophagini. In: Löbl, I. & Smetana, A. (eds) *Catalogue of Palaearctic Coleoptera*, Vol. 3. Apollo Books, Stenstrup, pp. 159–175.
- Löbl, I., Nekolajev, G.V. & Kral, D. (2006c) Geotrupidae. In: Löbl, I. & Smetana, A. (eds) *Catalogue of Palaearctic Coleoptera*, Vol. 3. Apollo Books, Stenstrup, pp. 84–92.
- Löbl, I., Rolcik, J., Kolibac, J., & Gerstmeier, R. (2007) Cleridae. In: Löbl, I. & Smetana, A. (eds) *Catalogue of Palaearctic Coleoptera*, Vol. 4. Apollo Books, Stenstrup, pp. 367–384.
- Löbl, I., Merkl, O., Ando, K., Bouchard, P., Lillig, M., Masomuto, K., & Schawaller, W. (2008a) Tenebrionidae without Lagriini, Asidini, Blaptini, Melanimini, Opatrini, Pedinini, Platyscelidini & Helopini. In: Löbl, I. & Smetana, A. (eds) *Catalogue of Palaearctic Coleoptera*, Vol. 5. Apollo Books, Stenstrup, pp. 105–352.
- Löbl, I., Nabozhenko, M. & Merkl, O. (2008b) Tenebrionidae: Tribe Blaptini. In: Löbl, I. & Smetana, A. (eds) *Catalogue of Palaearctic Coleoptera*, Vol. 5. Apollo Books, Stenstrup, pp. 219–229.
- Löbl, I. & Smetana, A. (2003) *Archostemata - Myxophaga - Adepaga*. Vol. 1. *Catalogue of Palaearctic Coleoptera*. Apollo Books, Stenstrup, Denmark. 819 p.
- Löbl, I. & Smetana, A. (2004) *Hydrophiloidea - Histeroidea - Staphylinoidea*. Vol. 2. *Catalogue of Palaearctic Coleoptera*. Apollo Books, Stenstrup, Denmark. 942 p.
- Löbl, I. & Smetana, A. (2006) *Scarabaeoidea, Scirtoidea, Dascilloidea, Buprestoidea, Byrrhoidea*. Vol. 4. *Catalogue of Palaearctic Coleoptera*. Apollo Books, Stenstrup, Denmark. 690 p.
- Löbl, I. & Smetana, A. (2007) *Elateroidea, Derodontoidea, Bostrichoidea, Lymexyloidea, Cleroidea, Cucujoidea*. Vol. 4. *Catalogue of Palaearctic Coleoptera*. Apollo Books, Stenstrup, Denmark. 935 p.
- Löbl, I. & Smetana, A. (2008) *Tenebrionoidea*. Volume 5. *Catalogue of Palaearctic Coleoptera*. Apollo Books, Stenstrup, Denmark. 670 p. <https://doi.org/10.1163/9789004260900>
- Löbl, I. & Smetana, A. (2010) *Chrysomeloidea*. Vol. 6. *Catalogue of Palaearctic Coleoptera*. Apollo Books, Stenstrup, Denmark. 924 p. <https://doi.org/10.1163/9789004260917>
- Löbl, I. & Smetana, A. (2011) *Curculionoidea I*. Vol. 7. *Catalogue of Palaearctic Coleoptera*. Apollo Books, Stenstrup, Denmark. 373 p. <https://doi.org/10.1163/9789004260931>
- Löbl, I. & Smetana, A. (2013) *Curculionoidea II*. Vol. 8. *Catalogue of Palaearctic Coleoptera*. Apollo Books, Stenstrup, Denmark. 700 p. <https://doi.org/10.1163/9789004259164>
- Lopatin, I., Smetana, A. & Schöller, M. (2010) Cryptocephalinae. In: Löbl, I. & Smetana, A. (eds) *Catalogue of Palaearctic Coleoptera*, Vol. 6. Apollo Books, Stenstrup, pp. 564–619.
- López-Pérez, J.J. (2016) Aportación al conocimiento de *Zophosis minuta* Fabricius, 1775 (Coleoptera: Tenebrionidae, Pimeliinae) en la provincia de Huelva (S.O. de Andalucía, España) y otros Zophosini Leach, 1815 de Murcia y Marruecos. *Archivos Entomológicos*, 16, 275–278.

- Mahari, S. (1992) *Étude Synécologique des Niveaux de Peuplement Arthropodiens de l'Écosystème Chêne-liège dans les Subéraies de la Mamora et de Benslimane* (Unpublished Ph.D. Dissertation). University Saint-Jérôme, Aix-Marseille, France. 200 p.
- Magnano, L. & Alonso-Zarazaga, M.A. (2013) Tribe Otiiorhynchini. In: Löbl, I. & Smetana, A. (eds) *Catalogue of Palaearctic Coleoptera*, Vol. 8. Brill, Leiden, pp. 302–347.
- Majka, C.G., Langor, D. & Rucker, W.H. (2009) Latridiidae (Coleoptera) of Atlantic Canada: New records, keys to identification, new synonyms, distribution, and zoogeography. *The Canadian Entomologist*, 141 (4), 317–370. <https://doi.org/10.4039/n09-050>
- Marggi, W.A., Huber, C., Müller-Motzfeld, G. & Hartmann, M. (2003) Carabidae, Tribe Bembidiini; subtribe Bembidiina. In: Löbl, I. & Smetana, A. (eds) *Catalogue of Palaearctic Coleoptera*, Vol. 1. Apollo Books, Stenstrup, pp. 241–273.
- Mas-Peinado, P., Ruiz, J.L., Merkl, O., Buckley, D., & García-París, M. (2021) Taxonomy of the North Moroccan and Iberian species of the subgenus *Amblypteraca* (Coleoptera: Tenebrionidae: Pimeliinae: Pimelia). *Zootaxa*, 4963 (3), 457–482. <https://doi.org/10.11646/zootaxa.4963.3.4>
- Mayor, A. (2007a) Dasytidae. In: Löbl, I. & Smetana, A. (eds) *Catalogue of Palaearctic Coleoptera*, Vol. 4. Apollo Books, Stenstrup, pp. 388–415.
- Mayor, A. (2007b) Melyridae. In: Löbl, I. & Smetana, A. (eds) *Catalogue of Palaearctic Coleoptera*, Vol. 4. Apollo Books, Stenstrup, pp. 386–388.
- Mazur, S. (2004) Histeridae. In: Löbl, I. & Smetana, A. (eds) *Catalogue of Palaearctic Coleoptera*, Vol. 2. Apollo Books, Stenstrup, pp. 68–102.
- Meregalli, M. (2013) Cyclominae. In: Löbl, I. & Smetana, A. (eds) *Catalogue of Palaearctic Coleoptera*, Vol. 8. Brill, Leiden, pp. 245–251.
- Métro, A. & Sauvage, C. (1955) *Flore des Végétaux Ligneux de la Mamora. La Nature au Maroc*. Société des Sciences naturelles du Maroc, Rabat. 498 p.
- Micó, E., Piau, O., Galante, E. & Lumaret, J.-P. (2003) Taxonomy of Iberian *Hoplia* (Col., Scarabaeoidea, Hopliinae) based on mtDNA analysis. *Molecular Phylogenetics and Evolution*, 26, 348–353. [https://doi.org/10.1016/S1055-7903\(02\)00302-0](https://doi.org/10.1016/S1055-7903(02)00302-0)
- Morrone, J.J. (2013) The subtribes and genera of the tribe Listroderini (Coleoptera, Curculionidae, Cyclominae): Phylogenetic analysis with systematic and biogeographical accounts. *ZooKeys*, 273, 15–71. <https://doi.org/10.3897/zookeys.273.4116>
- Mouna, M. & Arahou, M. (1986) Comparaison des peuplements de coléoptères frondicoles de deux essences forestiers résineuses au Maroc: Cèdre d'Atlas (*Cedrus atlantica*) et Thuya (*Tetraclinis articulata*). *Bulletin de l'institut scientifique*, Rabat, 1, 133–142.
- Murphy, E.F. (1988) Some legal solutions for contemporary problems concerning groundwater and aquifers. *Journal of Natural Resources & Environmental Law*, 4 (1), 4.
- Nageleisen, L.M. & Bouget, C. (2009) *L'étude des Insectes en Forêt: Méthodes et Techniques, Éléments Essentiels pour une Standardisation. Synthèse des Réflexions Menées par le Groupe de Travail "Inventaires Entomologiques en Forêt. Les dossiers Forestiers n°19*, Office National des forêts, Paris. 144 p.
- Nieto, A. & Alexander, K.N.A. (2010) *European Red List of Saproxyllic Beetles*. Publications Office of the European Union, Luxembourg. 46 p.
- Nikodym, M. & Bezdek, A. (2006) Glaphyridae. In: Löbl, I. & Smetana, A. (eds) *Catalogue of Palaearctic Coleoptera*, Vol. 3. Apollo Books, Stenstrup, pp. 97–103.
- Novak, V. & Pettersson, R. (2008) Tenebrionidae: Alleculinae. In: Löbl, I. & Smetana, A. (eds) *Catalogue of Palaearctic Coleoptera*, Vol. 5. Apollo Books, Stenstrup, pp. 319–338.
- Pelletier, J. (2013) Tribe Brachyderini. In: Löbl, I. & Smetana, A. (eds) *Catalogue of Palaearctic Coleoptera*, Vol. 8. Leiden, Brill, pp. 255–263.
- Pérez-Vera, F., Ávila, J. M. (2012) Los Asidini marroquíes. Ensayo monográfico sobre la tribu Asidini (Coleoptera, Tenebrionidae) en el Reino de Marruecos. *Monografías Electrónicas S.E.A.*, 3, 1–344.
- Perreau, M. (2004) Leiodidae. In: Löbl, I. & Smetana, A. (eds) *Catalogue of Palaearctic Coleoptera*, Vol. 2. Apollo Books, Stenstrup, pp. 133–203.
- Pittino, R. (2006) Trogidae. In: Löbl, I. & Smetana, A. (eds) *Catalogue of Palaearctic Coleoptera*, Vol. 3. Apollo Books, Stenstrup, pp. 79–81.
- Prena, J. (2011) Tribe Baridini. In: Löbl, I. & Smetana, A. (eds) *Catalogue of Palaearctic Coleoptera*, Vol. 7. Apollo Books, Stenstrup, pp. 252–260.

- Prudhomme, J.C. (2016) Une étude locale de la biodiversité : inventaire des coléoptères du domaine de la fondation Pierre Vérots à Saint-Jean-de-Thurigneux (Ain, France). 3. Les coléoptères saproxyliques. *Bulletin Mensuel de la Société Linnéenne de Lyon*, 85 (1-2), 23–58. <https://doi.org/10.3406/linly.2016.17776>
- Puchkov, A.V. & Matalin, A.V. (2003) Carabidae: subfamily Cicindelinae; tribe Cicindelini. In: Löbl, I. & Smetana, A. (eds) *Catalogue of Palaearctic Coleoptera*, Vol. 1. Apollo Books, Stenstrup, pp. 99–115.
- QCN (2021) *Quatrième Communication Nationale du Maroc à la Convention Cadre des Nations Unies sur le Changement Climatique. Chapitre 6.4 : Vulnérabilités et Programmes nationaux d'adaptation aux effets néfastes du changement climatique. La Forêt*. Royaume du Maroc, Ministère de l'Énergie, des Mines et de l'Environnement, Département de l'Environnement, Rabat, pp. 183–192.
- Ranius, T., Jansson, N. (2002). A comparison of three methods to survey saproxylic beetles in hollow oaks. *Biodiversity and Conservation*, 11, 1759–1771. <https://doi.org/10.1023/A:1020343030085>
- Rankou, H., Culham, A., Jury, S.L. & Christenhusz, M.J.M. (2013) The endemic flora of Morocco. *Phytotaxa*, 78 (1), 1–69. <https://doi.org/10.11646/phytotaxa.78.1.1>
- Rosas-Ramos, N., Mas-Peinado, P., Gil-Tapetado, D., Recuero, E., Ruiz, J.L. & García-París, M. (2020) Catalogue, distribution, taxonomic notes, and conservation of the Western Palearctic endemic hunchback beetles (Tenebrionidae, *Misolampus*). *ZooKeys*, 963, 81–129. <https://doi.org/10.3897/zookeys.963.53500>
- Rossi, M. & Vallauri, D. (2013) *Evaluer la Naturalité. Guide Pratique*. version 1.2. WWF, Marseille. 154 p.
- Rozicka, J. & Schneider, I. (2004) Silphidae. In: Löbl, I. & Smetana, A. (eds) *Catalogue of Palaearctic Coleoptera*, Vol. 2. Apollo Books, Stenstrup, pp. 229–237.
- Ruzzier, E., Ghahari, H. & Horák, J. (2017) A checklist of the Iranian Mordellidae (Coleoptera: Tenebrionoidea). *Zootaxa*, 4320 (1), 146–158. <https://doi.org/10.11646/zootaxa.4320.1.8>
- Sama, G. & Löbl, I. (2010a) Cerambycinae. In: Löbl, I. & Smetana, A. (eds) *Catalogue of Palaearctic Coleoptera*, Vol. 6. Apollo Books, Stenstrup, pp. 143–206.
- Sama, G. & Löbl, I. (2010b) Lepturinae. In: Löbl, I. & Smetana, A. (eds) *Catalogue of Palaearctic Coleoptera*, Vol. 6. Apollo Books, Stenstrup, pp. 114–117.
- Sanchez-Ruiz, A. & Löbl, I. (2007) Cebrionina. In: Löbl, I. & Smetana, A. (eds) *Catalogue of Palaearctic Coleoptera*, Vol. 4. Apollo Books, Stenstrup, pp. 89–94.
- Sauvage, C. (1961) Flore des suberaies marocaines (Catalogues des Cryptogames vasculaires et des phanérogames). *Travaux de l'Institut Scientifique Chérifien, série Botanique*, 22, 1–252.
- Sforzi, A. (2011) Brentidae. In: Löbl, I. & Smetana, A. (eds) *Catalogue of Palaearctic Coleoptera*, Vol. 7. Apollo Books, Stenstrup, pp. 142–148.
- Skuhrovec, J. (2013) Hyperinae. In: Löbl, I. & Smetana, A. (eds) *Catalogue of Palaearctic Coleoptera*, Vol. 8. Brill, Leiden, pp. 423–437.
- Smetana, A. (2004) Staphylinidae. In: Löbl, I. & Smetana, A. (eds) *Catalogue of Palaearctic Coleoptera*, Vol. 2. Apollo Books, Stenstrup, pp. 237–699.
- Smetana, A. (2006a) Melolonthidae: Tribe Hopliini. In: Löbl, I. & Smetana, A. (eds) *Catalogue of Palaearctic Coleoptera*, Vol. 3. Apollo Books, Stenstrup, pp. 184–190.
- Smetana, A. (2006b) Cetoniinae. In: Löbl, I. & Smetana, A. (eds) *Catalogue of Palaearctic Coleoptera*, Vol. 3. Apollo Books, Stenstrup, pp. 283–313.
- Soldati, F. (2008) Pimeliinae: Tribe Asidini. In: Löbl, I. & Smetana, A. (eds) *Catalogue of Palaearctic Coleoptera*, Vol. 5. Apollo Books, Stenstrup, pp. 128–139.
- Speight, M.C.D. (1989) *Les Invertébrés Saproxyliques et leur Protection*. Collection Sauvegarde de la nature, Conseil de l'Europe, Strasbourg, 42. 77 p.
- Švec, Z. (2007) Phalacridae. In: I. Löbl & A. Smetana (eds) *Catalogue of Palaearctic Coleoptera*, Vol. 4. Appollo Books, Stenstrup, pp. 507–513.
- Švihla, V. (2008) Oedemeridae. In: Löbl, I. & Smetana, A. (eds) *Catalogue of Palaearctic Coleoptera*, Vol. 5. Apollo Books, Stenstrup, pp. 353–369.
- Tronquet, M., Benyahia, Y., Valladares, L., Brustel, H. (2019) Complément à la liste des espèces patrimoniales de Coléoptères du parc national de Talassemtane au Maroc: famille des Staphylinidae (sous-familles Pselaphinae et Scydmaeninae exclues) (Coleoptera, Staphylinidae). *Revue de l'Association Roussillonnaise d'Entomologie*, 28 (4), 251–264.
- IUCN (2022) *The IUCN Red List of Threatened Species*. Available from <https://www.iucnredlist.org/en> [Accessed December 2, 2023]

- Velazquez de Castro, A.J. (2013) Tribe: Sitonini. In: Löbl, I. & Smetana, A. (eds) *Catalogue of Palaearctic Coleoptera*, Vol. 8. Brill, Leiden, pp. 386–392.
- Velle, L. (2004) *Inventaire des coléoptères de la réserve naturelle du Val d'Allier*. Office National des Forêts, France. 30 p.
- Velle, L. (2011) *Inventaire des Coléoptères saproxyliques dans la Réserve Naturelle Nationale du Val d'Allier (F-03)*. Rapport d'étude pour le compte de la DREAL Auvergne. RNN Val d'Allier & Réseau entomologie de l'ONF. 69 p.
- Verdú, J.R. & Galante, E. (2006) *Libro Rojo de Los Invertebrados de España*. Dirección General de Conservación de la Naturaleza, Madrid. 411 p.
- Verdú, J.R. & Galante, E. (2009) *Atlas de los Invertebrados Amenazados de España (Especies en Peligro Crítico y en Peligro)*. Dirección General para la Biodiversidad, Ministerio de Medio Ambiente, Madrid. 340 p.
- Viñolas, A., Caballero-López, B. & Masó, G. (2017) The collection of type specimens belonging to the subfamily Pimeliinae (Coleoptera, Tenebrionidae) in the Natural Sciences Museum of Barcelona, Spain. *Arxius de Miscel·lània Zoològica*, 15, 30–92. <https://doi.org/10.32800/amz.2017.15.0030>
- Villemant, C. & Fraval, A. (1993) La faune entomologique du chêne-liège en forêt de la Mamora (Maroc). *Ecologia Mediterranea*, 19 (3–4), 89–98. <https://doi.org/10.3406/ecmed.1993.1725>
- Vit, S. (2004) Scydmaenidae: Tribe Scydmaenini. In: Löbl, I. & Smetana, A. (eds) *Catalogue of Palaearctic Coleoptera*, Vol. 2. Apollo Books, Stenstrup, pp. 224–228.
- Volkovitsh, M.G. (2006) Buprestidae. In: Löbl, I. & Smetana, A. (eds) *Catalogue of Palaearctic Coleoptera*, Vol. 3. Apollo Books, Stenstrup, pp. 325–342.
- Wegrzynowicz, P. (2007) Laemophloeidae. In: Löbl, I. & Smetana, A. (eds) *Catalogue of Palaearctic Coleoptera*, Vol. 4. Apollo Books, Stenstrup, pp. 503–507.
- Zorn, C. (2006) Rutelinae. In: Löbl, I. & Smetana, A. (eds) *Catalogue of Palaearctic Coleoptera*, Vol. 3. Apollo Books, Stenstrup, pp. 248–277.

## سخت‌بال پوشان (Coleoptera) مرتبط با جنگل بلوط مامورا در شمال غرب مراکش

حمزه هباز<sup>۱،\*</sup>، نورالدین معطوف<sup>۲</sup>، ژان-پییر لومار<sup>۳</sup>، لطیفه روحی<sup>۱</sup>

۱ آزمایشگاه اکولوژی و محیط زیست، دانشگاه حسن II کازابلانکا، دانشکده علوم بن مسیک، سیدی عثمان، کازابلانکا، مراکش.  
 ۲ مرکز نوآوری، تحقیق و آموزش، آژانس ملی آب و جنگل‌ها، رباط - آگدال، مراکش  
 ۳ دانشگاه پل والرئ مونیپلیه III، جاده ماند، مونیپلیه سیدکس ۵، فرانسه

\* پست الکترونیک نویسنده مسئول مکاتبه: [hamza.habbaz-etu@etu.univh2c.ma](mailto:hamza.habbaz-etu@etu.univh2c.ma)

تاریخ دریافت: ۲۹ آذر ۱۴۰۲ | تاریخ پذیرش: ۱۴ شهریور ۱۴۰۳ | تاریخ انتشار: ۱۲ دی ۱۴۰۳ |

**چکیده:** هدف این تحقیق، بررسی و بهبود دانش تنوع زیستی سخت‌بال‌پوشان در جنگل مامورا، یکی از بزرگترین جنگل‌های بلوط چوب‌پنبه‌ای در جهان، می‌باشد. مطالعات در طول دو سال متوالی (۲۰۲۱ و ۲۰۲۲) با استفاده از تله‌های فعال و غیرفعال انجام شد. در مجموع ۸۲۴۷ نمونه جمع‌آوری شد که نمایانگر ۲۵۶ گونه از ۴۲ خانواده از راسته Coleoptera بودند. بیشتر گونه‌های سوسک متعلق به خانواده‌های Tenebrionidae (۲۷ گونه)، Carabidae (۲۷ گونه) و Curculionidae (۲۴ گونه) بودند. از ۲۵۶ گونه شناسایی شده، ۲۱۶ مورد در این مقاله گنجانده شده‌اند. ما فقط گونه‌های با ارزش زیستی قابل توجه را انتخاب کردیم؛ ۵۳ مورد چوبخوار شامل ۶ گونه نادر؛ ۲۹ مورد اندمیک بودند و ۲۶ گونه به عنوان رکوردهای جدید برای فون مراکش شناسایی شدند. بر اساس نتایج این تحقیق، نخستین فهرست سخت‌بال‌پوشان در جنگل مامورا تهیه شده که اطلاعات جدیدی درباره توزیع سوسک‌ها در جنگل‌های بلوط چوب‌پنبه‌ای مراکش فراهم می‌کند.

**واژگان کلیدی:** تنوع زیستی، فهرست‌برداری، انتشار، بومی، سوسک‌های چوبخوار