


Original Article 

Arthropod diversity in Iran: Class Insecta–Polyneoptera, part I. Blattodea Brunner von Wattenwyl, 1882, Mantodea Burmeister, 1838, and Phasmatodea Jacobson & Bianchi, 1902

Zohreh Mirzaee¹  | Thies H. Büscher²  | Dominic Evangelista³  | Paul Eggleton⁴  | Leonid Anisyutkin⁵  | Frédéric Legendre⁶ 

1 Senckenberg German Entomological Institute, Eberswalder Str. 90, 15374 Müncheberg, Germany

2 Functional Morphology and Biomechanics, Kiel University, Am Botanischen Garten 1–9, 24118 Kiel, Germany; tbuescher@zoologie.uni-kiel.de

3 465 Morrill Hall, 505 S Goodwin Ave., University of Illinois, Urbana-Champaign: Urbana, Illinois, US 61801; dominice@illinois.edu

4 Soil Biodiversity Group, Department of Entomology, The Natural History Museum, London SW7 5BD, UK; p.eggleton@nhm.ac.uk

5 Laboratory of Insect Systematics, Zoological Institute of Russian Academy of Sciences, Universitetskaya nab., 1, 199034 Saint-Petersburg, Russia; leonid.dictyoptera@gmail.com

6 Institut de Systématique, Évolution, Biodiversité (UMR 7205), Muséum national d'Histoire naturelle, CNRS, SU, EPHE-PSL, UA, Paris F-75005, France; frederic.legendre@mnhn.fr

Corresponding author: Zohreh Mirzaee | [✉ zmirzaee1988@gmail.com](mailto:zmirzaee1988@gmail.com)

<https://zoobank.org/urn:lsid:zoobank.org:COCD9B32-5958-467E-8916-D966344BB875>

Academic Editor

Alireza Zamani

Received

October 11, 2026

Revised

February 14, 2026

Accepted

February 15, 2026

Published online

February 27, 2026

ABSTRACT. This work presents a comprehensive review of the insect orders Mantodea (praying mantises), Blattodea (cockroaches and termites), and Phasmatodea (stick insects) in Iran, based on literature, national checklists, and global taxonomic databases to develop an up-to-date faunistic inventory. Following the most recent accepted classifications, the review documents a total of three orders, 20 families, 44 genera, and 104 species. This includes 9 families and 19 genera, and 33 species of Mantodea, 10 families and 23 genera, and 65 species of Blattodea (including termites), and one family with two genera and 5 species of Phasmatodea. Among these groups, Mantodea is the most thoroughly documented, while Phasmatodea remains poorly studied, with only five species recorded to date. Integrating up-to-date taxonomic data is essential for resolving inconsistencies in older or unverified records and establishing a reliable baseline for biogeographical analysis, conservation planning, and the management of both native and invasive species. Especially for Phasmatodea, it is clear that the taxonomy of the Iranian fauna needs further study, both to reveal still undocumented diversity and to clarify the higher-level placement of the lineages occurring in Iran. Continued field surveys and integrative taxonomic research are crucial for fully documenting the diversity of these insect groups across Iran's varied habitats.

KEYWORDS: Biodiversity, Checklist, Cockroaches, Dictyoptera, Stick insects, Taxonomy, Termites

Citation: Mirzaee, Z., Büscher, T.H., Evangelista, D., Eggleton, P., Anisyutkin, L. & Legendre, F. (2026) Arthropod diversity in Iran: Class Insecta–Polyneoptera, part I. Blattodea Brunner von Wattenwyl, 1882, Mantodea Burmeister, 1838, and Phasmatodea Jacobson & Bianchi, 1902. *Journal of Insect Biodiversity and Systematics*, 12 (02), 357–394.

INTRODUCTION

The study of the insect orders Mantodea, Blattodea, and Phasmatodea in Iran has developed through a fragmented yet advancing research history. For much of the 20th century, taxonomic and faunistic accounts of these groups were embedded within broader Middle Eastern surveys (e.g., Uvarov 1922, 1938; Werner 1930; Kaltenbach 1963, 1982), with limited attention to Iranian-specific diversity. These early efforts nonetheless provided the groundwork for later targeted studies, especially in the case of Mantodea, which has become the most thoroughly documented of these groups. This work is part of an

Copyright © 2026 The Authors. This is an open access article distributed under the terms of the Creative Commons NonCommercial Attribution License (CC BY NC 4.0), which permits Share - copy and redistribute the material in any medium or format, and Adapt - remix, transform, and build upon the material, under the Attribution-NonCommercial terms.

ongoing series of reviews on the diversity of arthropods in Iran. The main objective of this initiative is to establish a comprehensive foundation for future investigations, promoting further taxonomic and biogeographical studies on Iranian arthropods. By documenting and synthesising available data, this series seeks to facilitate a deeper understanding of the country's arthropod fauna and encourage more extensive surveys across different regions and habitats.

Mantodea (praying mantises) in Iran were first recorded through pioneering works by Uvarov (1938, 1939), Beier (1956, 1968), and Kaltenbach (1982) in *Fauna of Saudi Arabia*, which provided the first records and descriptions of several new and poorly known species. These early accounts, based on restricted field collections and museum specimens, lacked a broad ecological or biogeographic scope. For decades thereafter, the Mantodea fauna of Iran continued to be treated sporadically, often as part of Arabian Peninsula or Middle Eastern surveys. A significant shift occurred over the past decade, driven by Iranian entomologists and reinforced by international collaborations, resulting in taxonomic revisions, species descriptions, and comprehensive biogeographical assessments (Mirzaee et al. 2023; Mirzaee 2024; Mirzaee et al. 2024a, 2024b; Kolnegari et al. 2025). These works increasingly employ molecular techniques to clarify species boundaries and integrate ecological data, addressing questions of behaviour, life history, and habitat use that had long been neglected (Schwarz & Roy 2019). Despite these advances, sampling gaps persist, especially in remote or ecologically varied regions.

Blattodea (cockroaches and termites) represents another important component of Iran's insect fauna, but remains relatively poorly studied at the national scale compared to many other insect groups. Despite their ecological significance (Bignell & Eggleton 2000; Sugimoto et al. 2000; Basset 2001; Tarli et al. 2014; Djernæs 2018; Pringle et al. 2019; Pérez-Gómez et al. 2023), national-level surveys have remained limited. The foundational checklist by Chopard (1921) and Mirzayans (1986) marked the first systematic catalogue of Iranian cockroaches, documenting 24 species across four families and eleven genera. Later works, particularly those by Hashemi-Aghdam and Oshaghi (2015), expanded this number to 26 species within 14 genera. Subsequent studies have emphasised both taxonomic and ecological aspects of Iranian Blattodea, identifying important synanthropic and pest species across urban and agricultural environments (Bey-Bienko 1950; Ghayourfar 1998, 1999, 2005; Ravan & Akhtar 1999; Hanafi-Bojd & Sadaghiani 2009). Nevertheless, large areas of the country, particularly in arid and mountainous zones, remain poorly sampled, and many older records lack voucher confirmation. This underlines the need for continued faunistic and molecular investigations to clarify species identities, update distributions, and align the national fauna with current higher-level taxonomy.

Comprehensive, verified species lists are crucial, not only for understanding biodiversity patterns, but also for applied fields such as ecological monitoring (Mirzayans 1986; Hashemi-Aghdam & Oshaghi 2015). Termite research in Iran has developed gradually from scattered early records to more systematic faunistic and ecological studies in recent decades (Ravan & Akhtar 1993; Ravan & Akhtar 1999; Ravan et al. 2015). The first mentions of termites in the country go back to mid- to late-20th-century work on wood-destroying insects in the Middle East, but the fauna remained poorly known, and many taxa were undescribed until more focused surveys began in the 1990s and 2000s (Ghayourfar 2005). Subsequent targeted collecting across arid and semi-arid regions, especially in southern and south-eastern Iran, revealed a diverse assemblage of genera such as *Anacanthotermes* Jacobson, 1905, *Microcerotermes* Silvestri, 1901, *Amitermes* Silvestri, 1901, and *Heterotermes* Froggatt, 1897, and led to the description of several new species (Ghayourfar 2005).

Phasmatodea (stick insects) remains one of the least studied insect groups in Iran. Only five species within two genera are currently known from the country, based primarily on historical collections and scattered reports. Early references include works by Brunner von Wattenwyl (1907), Mistshenko (1937), and Bey-Bienko (1946), with later summaries provided by Otte & Brock (2005), Brock (1991, 2007), and Brock & Büscher (2022). Most known specimens originate from the provinces of Kerman, Khorasan, and Qazvin, housed in the Zoological Museum of the Russian Academy of Sciences, St. Petersburg. The extremely limited data on Iranian Phasmatodea likely reflect both their cryptic habits and insufficient fieldwork, rather than genuinely low diversity. Given Iran's wide range of climatic and ecological zones, from humid Caspian forests to central deserts and southeastern subtropical habitats, further exploration will clearly reveal a richer phasmid fauna than currently recognised. Integrative taxonomic studies

combining morphology, biogeography, and molecular evidence are urgently needed to clarify the country's phasmid diversity and distribution (Brock & Büscher 2022).

Together, these three insect orders represent components of Iran's polyneopteran diversity. Accurate and up-to-date species lists are foundational for future research and biodiversity assessments. They enable informed ecological studies, support biogeographic analyses, and contribute to taxonomic stability by providing a reliable reference framework (Marques et al. 2024). Furthermore, they are vital for conservation efforts, particularly in identifying endemic or threatened taxa. Given the presence of several dubious species records in Iranian literature, the need for careful validation and critical taxonomic review is paramount (Kolnegari 2023). The integration of morphological, ecological, and molecular data will be essential in addressing these gaps and ensuring the long-term integrity of the country's biodiversity database (e.g., Mirzaee et al. 2024b for Mantodea).

MATERIAL AND METHODS

The data presented in this paper are compiled from published literature, national checklists, and global taxonomic databases, including GBIF and online open data platforms such as iNaturalist. These sources were identified through structured searches in Web of Science, Google Scholar, and Google search engine as well as through specialized online resources: Mantodea Species File Online (<http://mantodea.speciesfile.org>), Cockroach Species File Online (<https://cockroach.speciesfile.org/>), Isoptera Species File Online (<https://isoptera.speciesfile.org/>), the Termites Database (<https://termitologia.net/catalog/index.php>), and Phasmida Species File Online (<https://phasmida.speciesfile.org/>). The extracted data were then checked and refined based on the most recent taxonomic updates as well as the examination of museum collections. The classification and nomenclature employed in this study follow widely accepted recent frameworks: Schwarz and Roy (2019) and Otte et al. (2023) for Mantodea; Deng et al. (2026) for Blattodea, including termites; and Brock (1991, 2007), Otte and Brock (2005), Brock and Büscher (2022), and Brock et al. (2025) for Phasmatodea. However, the taxonomic placement of the tribe Gratidiini Cliquennois, 2005, which includes all phasmid taxa known from Iran, and the phylogenetic relationships of these taxa require taxonomic reassessment (Bank & Bradler 2020). The taxonomic placement herein, consequently, requires revision based on molecular evidence. Bacillidae Brunner von Wattenwyl, 1893, and Gratidiini are para- or polyphyletic, and numerical representations at certain taxonomic levels in the checklist lack real phylogenetic meaning. These counts are provided only for comparative purposes and reflect the current but likely inaccurate taxonomic framework.

The literature consulted for Mantodea spans more than a century of taxonomic and faunistic research, drawing on foundational monographs and regional works. Key contributions include the early comprehensive works of Werner (1905, 1930), Bolívar (1911, 1913), and Uvarov (1922, 1938), followed by significant revisions and faunal accounts by Ramme (1951), Beier (1956), Deeleman-Reinhold (1957), Ebner (1963), La Greca and Lombardo (1982, 1987), Kaltenbach (1963, 1982), and Ehrmann (2002). More recent studies, such as those by Rabieh et al. (2016), Mirzaee and Pashaei Rad (2017), Kolnegari and Vafaei-Shoushtari (2018), Mirzaee and Sadeghi (2019, 2021), Moradzadeh et al. (2021), Kolnegari (2023), Kolnegari et al. (2025), and Mirzaee et al. (2023, 2024a, 2024b) provide updated regional records and taxonomic refinements. Research on Blattodea was founded on a combination of classical taxonomic references and modern regional surveys. Foundational sources include Desneux (1906), Bey-Bienko (1950), Harris (1957), Akhtar (1974), and Al-Alawy et al. (1990), complemented by later faunistic and applied studies such as Mirzayans (1986), Ghayourfar (1998, 1999, 2005), Hanafi-Bojd and Sadaghiani (2009), Hashemi-Aghdam and Oshaghi (2015), and Ravan & Akhtar (1999). Together, these works provide the primary framework for species identification and distributional interpretation. For Phasmatodea, the reference base integrates both historical and contemporary treatments, including Brunner von Wattenwyl (1907), Mistshenko (1937), Bey-Bienko (1946), Brock (1991, 2007), Otte & Brock (2005), Brock & Büscher (2022), and Brock et al. (2025). It should be noted that the numerical hierarchy applied to phasmatodean groups in the present work is purely organisational; current molecular evidence demonstrates that the superfamily-, family-, and tribe-level classifications used here are not monophyletic and remain in need of comprehensive revision.

Detailed species information has been extracted primarily from the original descriptions and distributional reports provided in these sources. The number of species, genera, and families is indicated alongside each main taxonomic group. For each taxon, the number presented on the left side of the slash (/) indicates the number for Iran, and the corresponding global total is mentioned at the right.

Abbreviations used for the zoological institutes and museums mentioned in this study are as follow: **AMNH** American Museum of Natural History, New York, USA; **BMNH** British Museum, London, UK; **ETHZ** Public University, Zürich, Switzerland; **HMIM** Hayk Mirzayans Insect Museum, Insect Taxonomy Research Department, Plant Pests and Diseases Research Institute, Tehran, Iran; **IAUA** Islamic Azad University Arak, Arak, Iran; **IEA** Istituto di Entomologia Agraria, Portici, Italy; **INER** National Institute of Entomology, Rom, Italy; **INHM** Iraq Natural History Museum, Baghdad, Iraq; **LSUK** Linnean Society United Kingdom, UK; **MNHN** Museum National d'Histoire Naturelle, Paris, France; **MNMS** National Museum of Natural Sciences Madrid, Spain; **MSNG** Museo Civico di Storia Naturale "giacomo Doria", Genova, Italy; **NCIP** National Collection of Insects, Pretoria, South Africa; **NHMB** Naturhistorisches Museum, Basel, Switzerland; **NHMM** Natuurhistorisch Museum Museum of Maastricht, Maastricht, the Netherlands; **NHMW** Naturhistorisches Museum, Wien, Austria; **NHRS** Naturhistoriska Riksmuseet, Stockholm, Sweden; **OUMNH** Oxford University Museum of Natural History, UK; **RIB** Institut Royal des Sciences Naturelles de Belgique, Bruxelles, Belgium; **SDEI** Senckenberg German Entomological Institute, Müncheberg, Germany; **SMNS** Natural History Museum Stuttgart, Germany; **USIU** United States International University, Kenya, Africa; **USNM** National Museum of Natural History Washington, USA; **UuzM** Uppsala University, Zoological Museum, Uppsala, Sweden; **ZIHS** Zoötomiska Institute, Hogskolas, Stockholm, Sweden; **ZIN RAS** Zoological Institute of the Russian Academy of Sciences, St. Petersburg, Russia; **ZMB** Museum für Naturkunde an der Universität Humbolt zu Berlin, Berlin, Germany; **ZMH** Zoologisches Museum für Hamburg, Hamburg, Germany; **ZSI** Zoological Survey of India, Calcutta, West Bengal, India; **ZUPL** Department of Zoology, University of Punjab, Lahore, Pakistan.

RESULTS

Class Insecta Linnaeus, 1758

Superorder Dictyoptera Latreille, 1829

Order Blattodea Wattenwyl Brunner Von, 1882

Epifamily Termitoidae Latreille, 1802 [5/13 families]

Family Heterotermitidae Froggatt, 1897 [2/3 genera]

Genus *Heterotermes* Froggatt, 1897 [1/32 species]

Heterotermes indicola (Wasmann, 1902)

Leucotermes indicola Wasmann, 1902:118–119. Lectotype soldier (NHMM).– India.

Distribution. Afghanistan, Bangladesh, India, Iran, Nepal, Pakistan.

Distribution in Iran. Fars, Isfahan, Kerman, Khorasan-e Razavi, Khuzestan, Mazandaran, Sistan-o Baluchistan, Tehran (Ravan & Akhtar 1998, 1999).

Remark. *Heterotermes indicola* is mainly in association with *Populus* plantations and other woody hosts in arid to semi-humid lowlands.

Genus *Reticulitermes* Holmgren, 1913 [2/25 species]

Reticulitermes clypeatus Lash, 1952

Reticulitermes clypeatus Lash, 1952:1–5. Holotype: imago (AMNH).– Israel.

Distribution. Iran, Iraq, Palestine/Israel, Romania.

Distribution in Iran. Western and southwestern provinces of Iran (Weidner 1960; Ghayourfar 2006).

Remarks. This species has been reported from the western and southwestern parts of Iran, where populations assigned to "*Reticulitermes* sp. near *lucifugus*" were later interpreted as belonging to the *R. clypeatus*–*R. lucifugus* complex (Weidner 1960; Ghayourfar 2006). Records from Iran are based on zoogeographical and faunistic studies that document *Reticulitermes* occurrences in the Zagros and adjacent regions, although most Iranian material has historically been referred only to "*Reticulitermes* sp. near *lucifugus*," indicating that the precise distribution and taxonomic status of *R. clypeatus* in Iran still require confirmation through integrative morphological and molecular analyses (Ghayourfar 2006; Luchetti et al. 2007).

***Reticulitermes lucifugus* Rossi, 1792**

Termes lucifugum Rossi, 1792:107. Syntypes: imago (ZMB).—"Shores of Mediterranean".

Syn. *Hemeroobius raphidioides* Villers, 1789:56. *Termes radicum* Latreille, 1794:84–85. *Termes parvulum* Illiger, 1807:16. *Termes pusillus* Walker, 1853:528–529.

Distribution. Algeria, Austria, Azerbaijan, Bosnia-Herzegovina, Bosnia and Herzegovina, Croatia, Cyprus, Egypt, France, Germany, Greece, Iran, Iraq, Italy, Madeira Island, Montenegro, Morocco, North Macedonia, Portugal, Romania, Russia, Serbia, Slovenia, Spain, Switzerland, Turkey, Ukraine.

Distribution in Iran. Northern Provinces (Austin 2006; Ghayourfar 2006; Krishna et al. 2013; Dedeine et al. 2016).

Remarks. *Reticulitermes lucifugus* has been reported from northern Iran, with records based on material identified as *Reticulitermes* sp. near *lucifugus* and treated within the *R. lucifugus* complex in zoogeographical and genetic studies (Austin 2006; Ghayourfar 2006; Krishna et al. 2013; Dedeine et al. 2016). These works indicate its presence in forested and montane habitats in northern Iran, but also stress that eastern Mediterranean populations (including Iran) may represent distinct, closely related taxa pending further taxonomic revision.

Family Hodotermitidae Desneux, 1904 [1/3 genera]

Genus *Anacanthotermes* Jacobson, 1905 [8/16 species]

***Anacanthotermes ahngerianus* (Jacobson, 1905)**

Hodotermes (*Anacanthotermes*) *ahngerianus* Jacobson, 1905:67. Syntypes: imago (AMNH).—Turkmenistan.

Syn. *Anacanthotermes ahngerianus opacus* Luppova, 1958:102, 104.

Distribution. Iran, Kazakhstan, Turkmenistan, Uzbekistan.

Distribution in Iran. Khorasan-e Razavi (Weidner 1960; Ghayourfar 1996).

Remarks. In Iran, *Anacanthotermes ahngerianus* is known mainly from rangelands and steppe habitats, with confirmed records from Khorasan Province in the northeast, where it occurs in pastures and causes damage to rural and sometimes urban structures (Weidner 1960; Ghayourfar 1996). The species is considered one of the dominant harvester termites of Iranian rangelands and is often treated together with other *Anacanthotermes* species in faunistic and ecological studies of the country's termite fauna (Harris 1967; Ghayourfar 1994, 1996).

***Anacanthotermes esmailii* Ghayourfar, 1998**

Anacanthotermes esmailii Ghayourfar, 1998:11, 12–15. Holotype: imago, soldier (HMIM).—Iran.

Distribution. Iran.

Distribution in Iran. Khorasan-e Razavi (Ghayourfar 1998).

***Anacanthotermes gurganiensis* Ravan & Akhtar, 1999**

Anacanthotermes gurganiensis Ravan & Akhtar, 1999:73–78. Holotype: major soldier (ZUPL).—Iran.

Distribution. Iran.

Distribution in Iran. Golestan (Ravan & Akhtar 1999).

***Anacanthotermes iranicus* Ravan & Akhtar, 1993**

Anacanthotermes iranicus Ravan & Akhtar, 1993:555–559. Holotype: soldier (ZUPL).– Iran.

Distribution. Iran.

Distribution in Iran. Khorasan-e Razavi (Ravan & Akhtar 1993).

***Anacanthotermes macrocephalus* (Desneux, 1906)**

Hodotermes macrocephalus Desneux, 1906:344–348. Syntypes: imago, soldier, worker (AMNH).– Pakistan.

Syn. *Anacanthotermes peshawarensis* Akhtar, 1974:40–44.

Distribution. India, Pakistan, Afghanistan, Iran.

Distribution in Iran. Unspecified.

***Anacanthotermes septentrionalis* (Jacobson, 1905)**

Hodotermes (Anacanthotermes) vagans septentrionalis Jacobson, 1905:65–67. Syntypes: imago (AMNH).– Iran.

Distribution. Afghanistan, Iran, Turkmenistan.

Distribution in Iran. Mazandaran (Jacobson 1905).

***Anacanthotermes turkestanicus* (Jacobson, 1905)**

Hodotermes (Anacanthotermes) turkestanicus Jacobson, 1905:61–65. Syntypes: imago, soldier, worker (AMNH).– Kazakhstan.

Distribution. Iran, Kazakhstan, Tajikistan, Turkmenistan, Uzbekistan.

Distribution in Iran. Unspecified.

***Anacanthotermes vagans* (Hagen, 1858)**

Hodotermes vagans Hagen, 1858:88–90, 240. Syntype imago (NHMW).– Iran.

Fig. 1A–B

Distribution. Afghanistan, Iran, Iraq, Kuwait, Pakistan, Saudi Arabia.

Distribution in Iran. Bushehr, "Karak Island" Persian Gulf (See Remark) (Hagen 1858).

Remarks. The Termite Database (<https://termitologia.net>) and the original species description (Hagen 1858) both list the locality as Shiraz, "Karak Island," Persian Gulf. We consider the name "Karak Island" to be erroneous and interpret it as Kharg (Khark) Island, which is situated in Bushehr Province in the northern Persian Gulf, rather than in Shiraz (Fars Province).

Family Kalotermitidae Froggatt, 1897 [1/23 genera]**Genus *Postelectrotermes* Krishna, 1961 [3/15 species]*****Postelectrotermes bidentatus* Ravan & Akhtar, 1999**

Postelectrotermes bidentatus Ravan & Akhtar, 1999:67–70. Holotype: major soldier (ZUPL).– Iran.

Distribution. Iran.

Distribution in Iran. Sistan-o Baluchistan (Ravan & Akhtar 1999).

***Postelectrotermes pasniensis* Akhtar, 1974**

Postelectrotermes pasniensis Akhtar, 1974:24–29. Holotype: soldier (ZUPL).– Pakistan.

Distribution. Iran, Pakistan.

Distribution in Iran. Khuzestan (Babmorad et al. 2014), and Sistan-o Baluchistan (Ravan 2010).

***Postelectrotermes zabuliensis* Ravan & Akhtar, 1999**

Postelectrotermes zabuliensis Ravan & Akhtar, 1999:70–73. Holotype: major soldier (ZUPL).– Iran.

Distribution. Iran.

Distribution in Iran. Sistan-o Baluchistan (Ravan & Akhtar 1999).

Family Psammotermitidae Holmgren, 1911 [1/2 subfamilies]

Subfamily Psammotermitinae Holmgren, 1911 [1/1 genus]

Genus Psammotermes Desneux, 1902 [2/6 species]

***Psammotermes prohybostoma* Ravan & Akhtar, 1999**

Psammotermes prohybostoma Ravan & Akhtar, 1999:78–81. Holotype: major soldier (ZUPL).– Iran.

Distribution. Iran.

Distribution in Iran. Sistan-o Baluchistan (Ravan & Akhtar 1999).

***Psammotermes rajasthanicus* Roonwal & Bose, 1960**

Psammotermes rajasthanicus Roonwal & Bose, 1960:38–39. Holotype: soldier (ZSI).– India.

Distribution. India, Pakistan, Iran.

Distribution in Iran. South and southeast of Iran (Ghayourfar 1996).

Family Termitidae Latreille 1802 [3/18 subfamilies]

Subfamily Amitermitinae Kemner, 1934 [1/14 genera]

Genus Amitermes Silvestri, 1901 [10/113 species]

***Amitermes baluchistanicus* Akhtar, 1974**

Amitermes baluchistanicus Akhtar, 1974:58–59. Holotype: soldier (ZUPL).– Pakistan.

Distribution. Iran, Pakistan.

Distribution in Iran. Fars, Hormozgan, Isfahan, Kerman, Khorasan-e Razavi, Khuzestan, Mazandaran, Sistan-o Baluchistan, Tehran (Ravan & Akhtar 1999).

***Amitermes belli* (Desneux, 1906)**

Termes belli Desneux, 1906:352–354. Syntypes: soldier, worker (AMNH).– Pakistan.

Distribution. India, Iran, Pakistan.

Distribution in Iran. Sistan-o Baluchistan (Ghayourfar 2001).

Remark. Iran lies at the junction of the Palearctic and Oriental realms, with most of its territory biogeographically Palearctic (Holt et al. 2013). Although the type locality of *A. belli* is Karachi in Sind (Oriental - Desneux 1906), the records attributed to Iran are treated here as Palaeartic, whereas India and Pakistan are listed under the Oriental region.

***Amitermes corpulentus* Al-Alawy, Abdul-Rassoul & Al-Azawi, 1990**

Amitermes corpulentus Al-Alawy, Abdul-Rassoul & Al-Azawi, 1990:25–27. Holotype: soldier (INHM).– Iraq.

Distribution. Iran, Iraq.

Distribution in Iran. Unspecified.

***Amitermes iranicus* Ghayourfar, 1999**

Amitermes iranicus Ghayourfar, 1999:20–22. Holotype: soldier (HMIM).– Iran.

Distribution. Iran.

Distribution in Iran. Kermanshah (Ghayourfar 1999).

***Amitermes kharrazii* Ghayourfar, 1999**

Amitermes kharrazii Ghayourfar, 1999:22–25. Holotype: imago, soldier, worker (HMIM).– Iran.

Distribution. Iran.

Distribution in Iran. Alborz and Tehran (Ghayourfar 1999).

***Amitermes messinae* (Fuller, 1922)**

Hamitermes messinae Fuller, 1922:125. Syntypes: imago (NCIP).– South Africa.

Syn. *Amitermes harleyi* Harris, 1957:425–427

Distribution. Egypt, Ethiopia, Iran, Kenya, Malawi, Saudi Arabia, South Africa, Sudan, Tanzania, Yemen, Zambia.

Distribution in Iran. Unspecified.

***Amitermes paravilis* Ravan & Akhtar, 1999**

Amitermes paravilis Ravan & Akhtar, 1999:81–84. Holotype: soldier (ZUPL).– Iran.

Distribution. Iran.

Distribution in Iran. Khuzestan (Ravan & Akhtar 1999).

***Amitermes stephensoni* Harris, 1957**

Amitermes stephensoni Harris, 1957:424–425. Holotype: soldier (BMNH).– Yemen.

Distribution. Guinea, Iran, Kenya, Nigeria, Oman, Saudi Arabia, Sudan, Yemen.

Distribution in Iran. Unspecified.

***Amitermes vilis* (Hagen, 1858)**

Termes (Termes) vilis Hagen, 1858:185–186, 240. Syntypes: imago (NHMW).– Iran.

Syn. *Amitermes wahrmani* Spaeth, 1964:27–29. *Amitermes rhizophagus* Belyaeva, 1974:63–73.

Distribution. Afghanistan, Iran, Iraq, Jordan, Oman, Palestine/Israel, Saudi Arabia, Turkmenistan, Yemen.

Distribution in Iran. Fars (Hagen 1858).

***Amitermes yasujensis* Ghayourfar, 2005**

Amitermes yasujensis Ghayourfar, 2005:64, 65–66. Holotype: soldier (HMIM).– Iran.

Distribution. Iran.

Distribution in Iran. Kohgiluyeh va Boyer Ahmad (Ghayourfar 2005).

Subfamily Microcerotermitinae Holmgren, 1910 [1/2 genera]**Genus *Microcerotermes* Silvestri, 1901 [11/152 species]*****Microcerotermes buettikeri* Chhotani & Bose, 1979**

Microcerotermes buettikeri Chhotani & Bose, 1979:77–80. Holotype: imago (NHMB).– Saudi Arabia.

Distribution. Iran, Saudi Arabia, Yemen.

Distribution in Iran. Khuzestan (Latifian et al. 2018).

***Microcerotermes chaudhryi* Akhtar, 1974**

Microcerotermes chaudhryi Akhtar, 1974:46–49. Holotype: soldier (ZUPL).– Pakistan.

Distribution. Iran, Pakistan.

Distribution in Iran. Unspecified.

***Microcerotermes chhotanii* Ghayourfar, 1998**

Microcerotermes chhotanii Ghayourfar, 1998:12, 15–17. Holotype: soldier (HMIM).– Iran.

Distribution. Iran.

Distribution in Iran. Kermanshah (Ghayourfar 1998).

***Microcerotermes dashlibronensis* Ghayourfar, 2005**

Microcerotermes dashlibronensis Ghayourfar, 2005:62, 64. Holotype: soldier (HMIM).– Iran.

Distribution. Iran.

Distribution in Iran. Golestan (Ghayourfar 2005).

***Microcerotermes diversus* Silvestri, 1920**

Microcerotermes diversus Silvestri, 1920:477–479. Syntypes: imago, soldier, worker (IEA).– Iraq.

Distribution. Iran, Iraq, Kuwait, Oman, Saudi Arabia, Yemen.

Distribution in Iran. Khuzestan (Habibpour et al. 2010).

***Microcerotermes gabrielis* Weidner, 1955**

Microcerotermes gabrielis Weidner, 1955:64–67. Syntypes: imago (ZMH).– Iraq.

Distribution. Afghanistan, Iran, Iraq, Saudi Arabia.

Distribution in Iran. Unspecified.

***Microcerotermes pakistanicus* Akhtar, 1974**

Microcerotermes pakistanicus Akhtar, 1974:54–56. Holotype: soldier (ZUPL).– Pakistan.

Distribution. India, Iran, Malaysia, Pakistan.

Distribution in Iran. Unspecified.

***Microcerotermes saravanensis* Ghayourfar, 2005**

Microcerotermes saravanensis Ghayourfar, 2005:62, 64. Holotype: soldier (HMIM).– Iran.

Distribution. Iran.

Distribution in Iran. Sistan-o Baluchistan (Ghayourfar 2005).

***Microcerotermes shahroudiensis* Ghayourfar, 2005**

Microcerotermes shahroudiensis Ghayourfar, 2005:39–44. Holotype: soldier (HMIM).– Iran.

Distribution. Iran.

Distribution in Iran. Semnan (Ghayourfar 2005).

***Microcerotermes sistaniensis* Ghayourfar & Akhtar, 2005**

Microcerotermes sistaniensis Ghayourfar & Akhtar, 2005:155–163. Holotype: imago, soldier (HMIM).– Iran.

Distribution. Iran.

Distribution in Iran. Sistan-o Baluchistan (Ghayourfar & Akhtar 2005).

***Microcerotermes varaminicus* Ghayourfar, 1999**

Microcerotermes varaminicus Ghayourfar, 1999:16, 18–20. Holotype: soldier (HMIM).– Iran.

Distribution. Iran.

Distribution in Iran. Tehran (Ghayourfar 1999).




Figure 1. Representative termite in Iran: **A.** *Anacanthotermes vagans* from Khuzestan: Dezful (photo credit by: Adel Ramezani); **B.** *Anacanthotermes vagans* from Khuzestan: Masjed Solyman (photo credit by: Farhad Eshraghi Mofrad); **C.** Unknown sp., Kermanshah province: Paveh (photo credit by: Fahmi Solimany); **D.** Unknown sp., Khuzestan: Dez (photo credit by: Morteza Johari); **E.** Unknown sp., Sistan-o Baluchistan: Saravan (photo credit by: Zahed Aram); **F.** Unknown sp., Isfahan: Najaf Abad; **G.** *Amitermes* sp. from Hormozgan: Siyahu (photo credit by: Mobin Kargarfard); **H.** *Amitermes* sp. from Tehran, photo credit by: Kian Saidabadi; **I.** Unknown sp. from Sabzevar (photo credit by: Kian Saidabadi).

Subfamily Macrotermitinae Kemner, 1934 [1/12 genera]

Genus *Microtermes* Wasmann, 1902 [2/70 species]

Microtermes mycophagus (Desneux, 1906)

Termes mycophagus Desneux, 1906:348–352. Syntypes: imago, soldier, worker (RIB).– Pakistan.

Distribution. India, Iran, Pakistan.

Distribution in Iran. Tehran and Mazandaran (Ravan et al. 2015).

Microtermes obesi Holmgren, 1912

Microtermes obesi Holmgren, 1912:787–788. Syntypes: imago (ZIHS).– India.

Syn. *Microtermes anandi* Holmgren, 1913:114; *Microtermes anandi curvignathus* Holmgren, 1913:114.

Distribution. Bangladesh, Bhutan, Cambodia, India, Iran, Malaysia, Myanmar, Pakistan, Sri Lanka, Thailand, and Vietnam.

Distribution in Iran. Unspecified.

Superfamily Blattoidea Latreille, 1810 [1/6 families]

Family Blattidae Latreille, 1810 [1/4 subfamilies]

Subfamily Blattinae Latreille, 1810 [2/25 genera]

Genus *Blatta* Linnaeus, 1758 [1/8 species]

Blatta orientalis Linnaeus, 1758

Blatta orientalis Linnaeus, 1758:1. Lectotype ♂ (LSUK), Paralectotypes ♂♂ ♀♀ (LSUK).—Southern Russia.

Syn. *Blatta badia* Saussure, 1863:150. *Blatta castanea* Blanchard, 1851:18. *Blatta culinaris* De Geer, 1773:530. *Blatta ferruginea* Thunberg, 1810. *Stylopyga orientalis gracilis* Adelung, 1903. *Blatta hemialata* Gistel, 1856:427. *Pulex imperator* Westwood, 1858. *Blatta lucifuga* Poda, 1761:49. *Kakerlac pallipes* Philippi, 1863. *Kakerlac platystetho* Philippi, 1863. *Blatta secunda* Schaeffer, 1769:222. *Stylopyga orientalis spontanea* Semenov-Tian-Shansky, 1909. *Blatta tertia* Schaeffer, 1769.

Distribution. Cosmopolitan (Marshall 1983).

Distribution in Iran. Almost all parts of Iran (Mirzayans 1986), Southwest Iran (Shahraki et al. 2010), Hamedan (Salehzadeh 1995; Mahjoob et al. 2010; Nejati et al. 2011), Tehran (Zarchi & Vatani 2009), Zanzan (Mohammadi 1998).

Genus *Periplaneta* Burmeister, 1838 [4/44 species]

Periplaneta americana (Linnaeus, 1758)

Fig. 2A

Blatta americana Linnaeus, 1758:424. Lectotype ♂ (LSUK). Unknown.

Syn. *Blatta aurelianensis* Fourcroy, 1785. *Periplaneta americana colorata* Rehn, 1902. *Blatta domicola* Risso, 1826. *Blatta ferrugineofusca* Gronovius, 1764. *Blatta heros* Eschscholtz, 1822. *Blatta kakkerlac* De Geer, 1773. *Blatta americana* Linnaeus, 1758. *Blatta orientalis* Sulzer, 1776. *Blatta siccifolia* Stoll, 1813. *Periplaneta stolidia* Walker, 1868.

Distribution. Cosmopolitan (Marshall 1983).

Distribution in Iran. Almost all parts of Iran (Mirzayans 1986), Southwest Iran (Shahraki et al. 2010), Hamedan (Salehzadeh 1995; Mahjoob et al. 2010; Nejati et al. 2011), Isfahan (Fathpour et al. 2003; Doroodgar et al. 2006), Khuzestan (Kassiri & Kazemi 2012), Kohkiluyeh va Boyer-Ahmad (Shahraki et al. 2011), Kurdistan (Vahabi et al. 2007), Mazandaran (Motavali Haghi et al. 1997), Tehran (Zarchi & Vatani 2009), Zanzan (Mohammadi 1998), Qom (Fahiminia et al. 2010).

Remarks. The original description of *Blatta americana* by Linnaeus did not clearly designate a precise type locality, and modern catalogues list the lectotype in LSUK without a specific locality label rather than a formal “Africa” as the type locality. Current distribution and phylogeographic work support an African (and Middle Eastern) origin of the species (Ma et al. 2019), which is why many modern sources describe its origin as African.

Periplaneta arabica (Bey-Bienko, 1938)

Shelfordella arabica Bey-Bienko, 1938:235. Holotype ♂ (ZIN RAS).— Saudi Arabia.

Fig. 2H

Distribution. Iran, Oman, Saudi Arabia, United Arab Emirates, Yemen.

Distribution in Iran. Hamedan-Hamedan (Mahjoob et al. 2010, Nejati et al. 2011), Dehloran, Ilam (Luo et al. 2023), Khorassan-e-Jonubi (Birjand), most southern states (Persian Gulf coast) (Mirzayans 1986), Yazd (Hanafi-Bojd & Sadaghiani 2009).

Periplaneta lateralis Walker, 1868

Periplaneta lateralis Walker, 1868:136. Holotype: Larva (BMNH).— Egypt.

Syn. *Shelfordella ahngeri* Adelung, 1910. *Paraloboptera sillemi* Hanitsch, 1935. *Periplaneta tartara* Saussure, 1874. *Shelfordella zarudnyi* Adelung, 1910.

Material examined. 2♂♂ adults, 7–9.VI.1957, R. Pasquier, Kermanshah Province (Qasr-e Shirin—**New record**, MNHN); 2♀♀ adults, 18.V.1955, G. Remaudière, Sistan-o Balouchestan (Kash—**New record**, MNHN; Jalgh, Pishin, Sarbaz, Saravan, Zabol).

Distribution. Northern Africa to Central Asia (Afghanistan, Egypt, Iran, Iraq, Jordan, Libya, Pakistan, Palestine/Israel, southern Russia, Sudan, Uzbekistan), with introduced populations in southern USA, Mexico, southern and central Europe, Japan, and other warm urban areas.

Distribution in Iran. Ardebil, Fars, Golestan, Guilan (Hashemi-Aghdam & Oshaghi 2015), Hamedan (Salehzadeh 1995), Hormozgan (Mirzayans 1986), Isfahan (Mirzayans 1986; Fathpour et al. 2003), Kerman (Mirzayans 1986), Kermanshah, Khuzestan (Mirzayans 1986), Kohkiluyeh va Boyer-Ahmad (Shahraki et al. 2011), Sistan-o Balouchestan, Southwest Iran (Shahraki et al. 2010), Tehran, Yazd (Hanafi-Bojd & Sadaghiani 2009).

Periplaneta monochroma Walker, 1871

Periplaneta monochroma Walker, 1871:37. Holotype ♂ (BMNH).– India.

Syn. *Shelfordella gussakovskii* Bei-Bienko, 1950. *Shelfordella tertia* Bei-Bienko, 1938.

Distribution. India, Tajikistan, Afghanistan, Iran.

Distribution in Iran. Fars (Hopkins & Beccaloni 2023), Hormozgan (Mirzayans 1986).

Superfamily Corydioidea Saussure, 1864 [1/2 families]

Family Corydiidae Saussure, 1864 [2/2 subfamilies]

Subfamily Corydiinae Saussure, 1864 [4/22 genera]

Genus *Polyphaga* Brullé, 1835 [4/5 species]

Polyphaga aegyptiaca (Linnaeus, 1758)

Blatta aegyptiaca Linnaeus, 1758:424. Syntype ♂ (UZM).– Somalia.

Syn. *Polyphaga brasiliiana* Saussure, 1864. *Blatta lata* Gistel, 1856. *Blatta omega* Costa, 1839. *Heterogamia punctata* Fischer von Waldheim, 1846. *Blatta sphaerica* Gistel, 1856. *Blatta spinifer* [sic] (Fischer von Waldheim, 1846). *Heterogamia spinipes* Fischer von Waldheim, 1846. *Steleopyga trichoprocta* Fischer von Waldheim, 1833.

Material examined. 1♂ adult, 14.VI.1957, R. Pasquier, Ilam Province (Tchavar–New record, MNHN); 3♂♂ adults, 10–12.VI.1957, R. Pasquier, Ilam Province (Mehran–New record, MNHN); ♂ adult, 11.VI.1957, R. Pasquier, Ilam Province (Mohassanab–New record, MNHN); 1♂ adult, 30.VI.1957, R. Pasquier, Khorassan Province (Guerayel–New record, MNHN); 1♀ adult, 5–10.V.1965, L. Matile, East Azerbaidjan Province (Tabriz–New record, MNHN).

Distribution. The range covers the whole Mediterranean region, southern parts of Europe (e.g., Italy, Balkans), and south-western Asia, reaching Iran in the east, and extends to parts of South America.

Distribution in Iran. Fars, Golestan, Ilam, (Mirzayans 1986), Isfahan (Fathpour et al. 2003, Kermanshah, Khuzestan (Vazirianzadeh et al. 2009), Lorestan, Mazandaran, Tehran.

Polyphaga indica Walker, 1868

Polyphaga indica Walker, 1868:14. Holotype ♀ (BMNH).– India.

Material examined. 1♀ adult, 19.IV.1966, Sistan-o Baluchistan province (Zabol–New record, MNHN).

Distribution. Central Asia, Iran, India, and Russia.

Distribution in Iran. Azarbaijan-e Sharghi, Fars, Isfahan, Kerman, Khorasan-e Razavi, Khuzestan, Mazandaran, Semnan, Sistan-o Balouchestan, Tehran (Mirzayans 1986).

Polyphaga saussurei (Dohrn, 1888)

Heterogamia saussurei Dohrn, 1888:131. Syntype ♀ (BMNH).– India.

Syn. *Polyphaga camelorum* Kirby, 1903. *Heterogamia pellucida* Redtenbacher, 1889.

Distribution. Afghanistan, Azerbaijan, India, Iran, Kazakhstan, Russia (southern), Tajikistan, Uzbekistan.

Distribution in Iran. Fars, Hormozgan, Khorasan-e Razavi, Sistan-o Balouchestan (Mirzayans 1986), and Yazd (Hanafi-Bojd & Sadaghiani 2009).

Remarks. *Polyphaga* sp. was recorded by Hashemi-Aghdam & Oshaghi (2015) from northern Tehran; it has not yet been formally described, and its material is deposited at the Entomology Museum of the School of Public Health, Tehran University of Medical Sciences.

Genus *Psammoblatta* Bey-Bienko, 1950 [4/5 species]

Psammoblatta livida (Brunner von Wattenwyl, 1865)

Heterogamia livida Brunner von Wattenwyl, 1865:359. Non-type: ♀ (BMNH).– Cyprus.

Syn. *Heterogamodes livida*: Chopard, 1929:308. *Arenivaga (Psammoblatta) livida*: Bey-Bienko, 1950:308

Distribution. Asia (central), Cyprus, Greece, Iran, Iraq, Syria, Turkey.

Distribution in Iran. Azarbaijan-e Sharghi (Jolfa, Aras) (Bey-Bienko 1950), Azarbaijan-e Sharghi (Tabriz-Khalat pushan) (Mirzayans 1986).

Psammoblatta longipes (Chopard, 1929)

Heterogamodes longipes Chopard, 1929:321. Holotype ♂, (MSNG?).– Iran.

Syn. *Arenivaga (Psammoblatta) longipes*: Bey-Bienko, 1950:314

Distribution. Iran.

Distribution in Iran. Fars, North of Iran (Chopard 1929; Bey-Bienko 1950), Tehran (Mirzayans 1986).

Remark. The question mark indicates uncertainty about the actual deposition of the holotype in this museum, because this information is based only on Chopard (1929:321), where he stated that the type was deposited there.

Psammoblatta roseni (Brancsik, 1898)

Heterogamia roseni Brancsik, 1898:19–20. No type data available.

Syn. *Heterogamodes roseni*: Chopard, 1929:314. *Arenivaga (Psammoblatta) roseni*: Bey-Bienko, 1950:311.

Distribution. Russia, Asia (central), Afghanistan, Iran.

Distribution in Iran. Isfahan (Fathpour et al. 2003), Khuzestan (Chopard 1929; Bey-Bienko 1950), Semnan, Sistan-o Balouchestan, Tehran (Mirzayans 1986).

Psammoblatta subhyalina (Chopard, 1921)

Polyphaga subhyalina Chopard, 1921:27. Holotype ♂ (BMNH), paratype ♂ (BMNH).– Iran.

Syn. *Heterogamodes subhyalina*: Chopard, 1929:318. *Arenivaga (Psammoblatta) subhyalina*: Bey-Bienko, 1950:310.

Hemelytroblatta subhyalina: Grandcolas, 1994:157.

Distribution. Iran.

Distribution in Iran. Qazvin (Chopard 1929; Bey-Bienko 1950), North of Iran (Bey-Bienko 1950), Fars (Mirzayans 1986).

Genus *Hemelytroblatta* Chopard, 1929 [3/25 species]

Hemelytroblatta (Mollidentoblatta) latifrons (Chopard, 1929)

Heterogamodes latifrons Chopard, 1929:319. Syntype ♂ (MNHN-EP735).– Iran.

Distribution. Iran, Iraq, Pakistan.

Distribution in Iran. Bushehr (Chopard 1929), Chaharmahal va Bakhtiari (Bey-Bienko 1950), Khuzestan (Mirzayans 1986).

Hemelytroblatta (Mollidentoblatta) marismortui (Janson, 1891)

Heterogamia (Polyphaga) maris-mortui Janson, 1891:184. Holotype ♂ (BMNH).– Egypt.

Syn. *Heterogamia maris-mortui* Janson, 1891

Distribution. Egypt (Sinai Peninsula), Iran, Iraq, Jordan, Palestine/Israel, Syria.

Distribution in Iran. Azarbaijan-e Gharbhi, Fars, Kermanshah, Khuzestan, Lorestan (Mirzayans 1986).

***Hemelytrobatta (Mollidentoblatta) persica* (Chopard, 1921)**

Polyphaga persica Chopard 1921:27:51. Holotype ♂ (BMNH).– Iran.

Distribution. Iran, Afghanistan.

Distribution in Iran. Fars, Isfahan, Kerman, Khorassan-e Jonubi (Chopard 1929), Khuzestan, Markazi, Qazvin, Sistan-o Balouchestan, and Yazd (Mirzayans 1986).

Genus *Leiopteroblatta* Chopard, 1969 [1/1 species]***Leiopteroblatta monodi* Chopard, 1969**

Leiopteroblatta monodi Chopard, 1969:228–230. Type: ♂ (MNHN-EP747).– Iran.

Distribution. Iran.

Distribution in Iran. Lut Desert (Mirzayans 1986).

Subfamily Euthyrrhaphinae Handlirsch, 1925 [1/3 genera]**Genus *Tivia* Walker, 1869** [1/15 species]***Tivia inconspicua* Bei-Bienko, 1950**

Tivia inconspicua Bey-Bienko, 1950:280. Holotype ♂ (ZIN RAS).– Iran.

Distribution. Iran.

Distribution in Iran. Hormozgan (Mirzayans 1986), Sistan-o Balouchestan (Bey-Bienko 1950).

Superfamily Blaberoidea Saussure, 1864 [3/5 families]**Family Blattellidae Karny, 1921** [3/125 genera]**Genus *Blattella* Caudell, 1903** [2/60 species]***Blattella biligata* (Walker, 1868)**

Ischnoptera biligata Walker, 1868:123. Lectotype ♂ (BMNH).– Ceylon.

Syn. *Blatta cercalis* Walker, 1868. *Ischnoptera evansi* Chopard, 1921. *Blatta mellea* Krauss, 1902. *Ischnoptera modestiformis* Karny, 1915. *Symploce mossambica* Princis, 1965. *Symploce persica* Bey-Bienko, 1935. *Phyllodromia unicolor* Brunner von Wattenwyl, 1893.

Material examined. 2 ♀♀ adults, 12.IV.1966, Sistan-o Baluchistan Province (Nikshar–**New record**, MNHN).

Distribution. Afghanistan, Burma (Myanmar), China, Congo (Republic of the Congo), Democratic Republic of the Congo, India, Iran, Iraq, Kenya, Mozambique, Pakistan, Palestine/Israel, Saudi Arabia, South Africa (Transvaal province), Sri Lanka, Taiwan, United Arab Emirates, Yemen (southern/coastal Afrotropical part), Zimbabwe.

Distribution in Iran. Hormozgan, Sistan-o Balouchestan (Bey-Bienko 1950, Mirzayans 1986).

***Blattella germanica* (Linnaeus, 1767)**

Blatta germanica Linnaeus. 1767:688. Syntypes (sex and exact number not stated in the original description); type depository not clearly specified in modern catalogues.

Syn. *Blatta asiatica* Pallas, 1773. *Blatta bivittata* Serville, 1838. *Phyllodromia cuneivittata* Hanitsch, 1925. *Blatta daurica* Laxmann, 1769. *Phyllodromia magna* Tepper, 1895. *Phyllodromia nitakana* Shiraki, 1931. *Blatta obliquata* Daldorf, 1793. *Ischnoptera paralella* Tepper, 1893. *Blatta paralella* Harris, 1835. *Ischnoptera paralella* [sic] (Princis, 1969). *Blattella germanica shuguroffi* Karny, 1908. *Blatta transfuga* Brünnich, 1763. *Ischnoptera bivittata* Thomas, 1876.

Fig. 21

Distribution. Cosmopolitan

Distribution in Iran. Almost all parts of Iran (Mirzayans 1986), Fars (Fallahzadeh et al. 2008), Hamedan

(Salehzadeh 1995; Mahjoob et al. 2010; Nejati et al. 2011), Hormozgan (Shahi et al. 2008), Isfahan (Fathpour et al. 2003; Doroodgar et al. 2006), Kerman (Kamyabi et al. 2006), Kermanshah (Limoe et al. 2011), Khorassan-e-Razavi (Naderinasab & Moravvej 2012), Kohkiluyeh va Boyer-Ahmad (Shahraki et al. 2011), Kordestan (Vahabi et al. 2007), Mazandaran (Motavali Haghi et al. 1997; Abedi & Dayer 2006; Enayati & Haghai 2007), Qom (Fahiminia et al. 2010), Southwest Iran (Shahraki et al. 2010), Tehran (Ladonni 1997; Mousavi et al. 2004; Baniardalani et al. 2006; Limoe et al. 2006; Nasirian 2007; Zarchi & Vatani 2009; Fakoorziba et al. 2010); Zanjan (Mohammadi 1998; Ghavami et al. 2005).

Genus *Parcoblatta* Hebard, 1917 [1/12 species]

***Parcoblatta* sp.**

Remark. This species was listed for Iran by Hashemi-Aghdam and Oshaghi (2015), who cited an earlier record from Isfahan (Fathpour et al. 2003) based on a study of cockroaches as reservoirs and vectors of drug-resistant *Salmonella* spp. In their paper, they noted that this represents a new record for the Iranian fauna and highlighted that its occurrence in the country requires confirmation through additional investigations. In the present work, this species is retained as reported by Hashemi-Aghdam and Oshaghi (2015). However, it is almost certainly a misidentification, given that *Parcoblatta* Hebard, 1917 is otherwise entirely endemic to North America (Hopkins & Beccaloni 2023).

Genus *Symploce* Hebard, 1916 [1/61 species]

***Symploce zarudniana* Bey-Bienko, 1950**

Symploce zarudniana Bey-Bienko, 1950:150. Holotype ♂ (probably in ZIN RAS).– Iran.

Distribution. Iran, Palestine/Israel.

Distribution in Iran. Fars, Hormozgan, Kerman, Makran, Sistan-o Balouchestan (Bey-Bienko 1950; Mirzayans 1986).

Family Ectobiidae Brunner von Wattenwyl, 1865 [1/14 genera]

Genus *Ectobius* Stephens, 1835 [1/68 genera]

***Ectobius (Ectobius) lapponicus* (Linnaeus, 1758)**

Blatta lapponica Linnaeus, 1758:425.– Middle Europe.

Remarks. This species is listed in the Cockroach Species File (Hopkins & Beccaloni 2023) with a distribution that includes Iran. However, we did not find any publication that currently confirms its presence in Iran, so this record must be regarded as tentative. In this study, the species is mentioned as potentially occurring in Iran, but its validity and occurrence require confirmation by future field and taxonomic work. Moreover, according to Bey-Bienko (1950), the southern border of the distribution of this species runs along the border of the forest-steppe, without entering Central Asia. Thus, the record of *E. lapponicus* from Iran is questionable and requires confirmation.

Genus *Phyllodromica* Fieber, 1853 [2/83 species]

***Phyllodromica (Phyllodromica) polita* (Krauss, 1888)**

Aphlebia polita Krauss, 1888:569.– Iran.

Distribution. Caucasus Mountains, Iran, Turkey.

Distribution in Iran. Guilan (Bey-Bienko 1950; Mirzayans 1986).

***Phyllodromica (Lobolampra) persa* Bey-Bienko, 1935**

Phyllodromica (Lobolampra) persa Bey-Bienko, 1935:122. Holotype ♂ (ZIN RAS).– Iran.

Distribution. Afghanistan, Iran, Tajikistan.

Distribution in Iran. Golestan (Bey-Bienko 1950).




Figure 2. Representative Cockroaches of Iran: **A.** *Periplaneta americana* nymph from Kerman: Baft, Torang Cave (photo credit by: Yaser Bakhshi); **B.** *Periplaneta* sp. nymph from Kohgiluyeh va Boyer-Ahmad: Gachsaran, Gakal Cave (photo credit by: Yaser Bakhshi); **C.** *Polyphaga* sp. from Eshtehard (photo credit by: Houman Doroudi); **D.** Blattidae gen. sp. from Pebdeh Cave, southern Zagros Mountains (photo credit by: Yaser Bakhshi); **E.** Blattidae gen. sp. from Tange Zendan Cave, Ghaemieh (photo credit by: Yaser Bakhshi); **F.** *Periplaneta* sp. nymph from Gorgan (photo credit by: Daryoosh Mansouri); **G.** *Supella longipalpa*, nymph from Karaj (photo credit by: Amin Ghaffari); **H.** *Periplaneta arabica* from Markazi: Neydeh Cave (photo credit by: Yaser Bakhshi); **I.** *Blattella germanica* nymph from Tehran (photo credit by: Kian Saidabadi).

Family Pseudophyllodromiidae Hebard, 1929 [3/88 genera]

Genus *Balta* Tepper, 1893 [1/92 species]

Balta ramifera (Walker, 1871)

Blatta ramifera Walker, 1871:27. Holotype ♀ (BMNH).– Indian Subcontinent, Nepal.

Distribution. Afghanistan, Iraq, Iran, Nepal.

Distribution in Iran. Sistan-o Balouchestan (Mirzayans 1986).

Genus *Supella* Shelford, 1911 [1/10 species]

Supella (Supella) longipalpa (Fabricius, 1798)

Blatta longipalpa Fabricius, 1798:185. Type material and type locality uncertain.

Syn. *Blatta cubensis* Saussure, 1862. *Blatta extenuata* Walker, 1868. *Blatta incisa* Walker, 1868. *Blatta phalerata* Saussure, 1863. *Ischnoptera quadriplaga* Walker, 1868. *Blatta subfasciata* Walker, 1871. *Blatta supellectilium* Serville, 1838. *Blatta transversalis* Walker, 1871. *Ischnoptera vacillans* Walker, 1868.

Fig. 2G

Distribution. Cosmopolitan.

Distribution in Iran. Hamedan (Salehzadeh 1995), Isfahan (Fathpour et al. 2003), Southwest Iran (Shahraki et al. 2010), Tehran (Bey-Bienko 1950).

Remarks. Extant species of the genus *Supella* are otherwise confined to the Afrotropical region (Africa, the Arabian Peninsula, and the Comoro Islands), with *S. longipalpa* as the only species that can be found nearly worldwide through human-mediated dispersal. Fossil congeners are known from Central America (Mexican amber, Chiapas) (Vršanský et al. 2011), the West Indies (Dominican amber) (Poinar 2022), and Europe (Rovno amber, Ukraine) (Anisyutkin & Perkovsky 2023).

Genus *Supellina* Chopard, 1921 [1/3 species]***Supellina buxtoni* Chopard, 1921**

Supellina buxtoni Chopard, 1921:48. Holotype (BMNH).– Iraq.

Distribution. Iraq, Iran, Afghanistan, the United Arab Emirates, and Egypt.

Distribution in Iran. Sistan-o Baluchestan (Bey-Bienko 1950).

Order Mantodea Latreille, 1802

Family Amorphoscelidae Stål, 1877 [1/2 subfamilies]

Subfamily Amorphoscelinae Stål, 1877 [1/5 genera]

Genus *Amorphoscelis* Stål, 1871 [1/50 species]

***Amorphoscelis pantherina* Roy, 1966**

Amorphoscelis pantherina Roy, 1966:268–270. Holotype ♂ (USNM).– Iraq.

Fig. 4F

Distribution. Iraq, Iran, Turkey.

Distribution in Iran. Fars (Mirzaee & Sadeghi 2021), Hormozgan (Ghahari & El-Den Nasser 2014).

Remark. This species was first recorded by Ghahari and El-Den Nasser (2014) from Hormozgan province, Minab (1 adult male, collected in April 2007), though the depository of the specimen remains unknown.

Family Nanomantidae Brunner von Wattenwyl, 1893 [1/4 subfamilies]

Subfamily Hapalomantinae Beier, 1964 [1/2 tribes]

Tribe Nilomantini Ehrmann & Roy, 2002 [1/9 genera]

Genus *Nilomantis* Werner, 1907 [1/2 species]

***Nilomantis floweri* Werner, 1907**

Nilomantis floweri Werner, 1907:234. Holotype ♀ (NHMW).– Sudan.

Syn. *Cryptomantis tenella* Giglio-Tos, 1915:45–46. *Nilomantis arabica* Beier, 1930:438–439. *Nilomantis [Cryptomantis] tenella* (Giglio-Tos, 1915); Beier, 1935:55. *Nilomantis arabica* Beier, 1930; Beier, 1935:55. *Cryptomantis tenella* Giglio-Tos, 1915; Roy, 1975:311–316. *Nilomantis arabica* Beier, 1930; Roy, 1975:311–316.

Fig. 4E

Distribution. Chad, Ethiopia (North), Iran, Mauritania, Niger, Oman, Saudi Arabia, Sudan, Yemen.

Distribution in Iran. Bushehr, Fars, Hormozgan, and Sistan-o Baluchistan (Mirzaee et al. 2023).

Family Gonypetidae Westwood, 1889 [1/2 subfamilies]

Subfamily Gonypetinae Westwood, 1889 [2/2 tribes]

Tribe Armenini Schwarz & Roy, 2019 [1/1 genus]

Genus *Armene* Stål, 1877 [1/7 species]

***Armene pusilla* Eversmann, 1854**

Mantis pusilla Eversmann, 1859:124–125. Holotype ♂ (ZIN RAS).– Kazakhstan.

Syn. *Gonypeta?* [*Mantis*] *pusilla* (Eversmann, 1859); Saussure, 1871:55. *Ameles alata* Saussure, 1872:51–52. *Ameles alata* Saussure, 1872; Saussure, 1874:13–14. *Armene* [*Ameles*] *alata* (Saussure, 1872); Stål, 1877:49. *Armene* [*Ameles*] *alata* (Saussure, 1872); Jacobson & Bianchi, 1905:145–146. *Armene* [*Ameles*] *alata* (Saussure, 1872); Kirby, 1904:225.

Fig. 3E

Distribution. Afghanistan, Iran, Mongolia, Russia, Tajikistan, Turkmenistan, Uzbekistan.

Distribution in Iran. Kurdistan (Sakenin et al. 2011).

Remark. The online Mantodea species catalogue (Otte et al. 2023) lists the type locality as “Mongolia: Kirghis Steppes (Südliche Kirgisensteppe östlich vom Ural-Fluss, Mongolei – Ehrmann 2002)”, but this formulation conflates the historical German locality name with modern geography and incorrectly assigns the site to Mongolia. In fact, “Kirgisensteppe” (Kirgiz/Kyrgyz Steppe) refers to steppe regions situated in present-day Kazakhstan and adjacent areas south and east of the Ural River, rather than to any locality in Mongolia.

Tribe Gonypetini Westwood, 1889 [2/4 subtribes]

Subtribe Gonypetina Westwood, 1889 [2/9 genera]

Genus *Elaea* Stål, 1877 [1/7 species]

***Elaea richteri* Beier, 1956**

Elaea richteri Beier, 1956:69–70. Holotype ♂ (SMNS).– Iran.

Fig. 3B

Distribution. (Iran).

Distribution in Iran. Southern and southeastern Iran (Beier 1956).

Genus *Holaptilon* Stål, 1877 [5/6 species]

***Holaptilon abdullahii* Mirzaee & Battiston 2024**

Holaptilon abdullahii Mirzaee & Battiston 2024:103. Holotype ♂ (SDEI), Paratypes ♀♀ (SDEI).– Iran.

Distribution. Iran.

Distribution in Iran. South of Iran, Bushehr (Mirzaee et al. 2024b).

***Holaptilon brevipugilis* Kolnegari, 2018**

Holaptilon brevipugilis Kolnegari, 2018:9–17. Holotype ♂, Allotype ♀ (IAUA), paratypes ♂♀ (SDEI).– Iran.

Syn. *Holaptilon yagmur* Yılmaz & Sevgili, 2023:18.

Distribution. Turkey, Iran.

Distribution in Iran. Centre and north-east of Iran, Markazi (Kolnegari & Vafaei-Shoushtari 2018), Khorasan-e Razavi (Mirzaee et al. 2024b)

***Holaptilon iranicum* Mirzaee & Sadeghi, 2024**

Holaptilon iranicum Mirzaee & Sadeghi, 2024:104–105. Holotype ♂ (SDEI).– Iran.

Distribution. Iran.

Distribution in Iran. Fars, Kohgiluyeh va Boyer-Ahmad (Mirzaee et al. 2024b).

***Holaptilon khozestani* Mirzaee & Battiston, 2024**

Holaptilon khozestani Mirzaee & Battiston 2024:105–106. Holotype ♂ (SDEI).– Iran.

Fig. 3D

Distribution. Iran.

Distribution in Iran. Khuzestan (Mirzaee et al. 2024b).

***Holaptilon tadovaniense* Mirzaee & Sadeghi, 2024**

Holaptilon tadovaniensis Mirzaee & Sadeghi, 2024:106–107. Holotype ♂ (SDEI).– Iran.

Distribution. Iran.

Distribution in Iran. Fars (Tadovan) (Mirzaee et al. 2024b).

Remarks. At the time this species was described, it was unintentionally given the name *tadovaniensis*. However, because *Holaptilon* is a neuter noun, the correct form of the specific epithet is *tadovaniense*. Therefore, we here adopt the corrected combination *Holaptilon tadovaniense*.

Family Rivetiniidae Ehrmann & Roy, 2002 [1/2 subfamilies]**Subfamily Rivetiniinae Ehrmann & Roy, 2002 [1/2 tribes]****Tribe Rivetiniini Ehrmann & Roy (2002) [3/7 genera]****Genus *Rivetina* Berland & Chopard, 1922 [5/34 species]*****Rivetina caucasica caucasica* (Saussure, 1871)**

Iris (Fischeria) caucasica Saussure, 1871:110–112. Holotype ♂, paratypes ♀ ♂ (NHMW).– Caucasus.

Syn. *Iris (Fischeria) caucasica* Saussure, 1871; Saussure 1871:149–362. *Fischeria [Iris (Fischeria)] caucasica* Saussure, 1871; Bolivar 1899:587. *Fischeria [Iris (Fischeria)] caucasica* Saussure, 1871; Uvarov 1912:8; *Eufischeriella [Iris (Fischeria)] caucasica* (Saussure, 1871); Giglio-Tos 1927:485. *Kinzelbachia ragnari* Harz, 1988:209. *Kinzelbachia ragnari* Harz, 1988; Ehrmann 2000:2–3.

Distribution. Iran, Caucasus, Syria, Tajikistan, and Turkey (Battiston et al. 2010).

Distribution in Iran. Lorestan (Mirzaee & Sadeghi 2019), West Azerbaijan (Sakenin et al. 2011).

***Rivetina dolichoptera* (Schulthess-Rechberg, 1894)**

Bolivaria dolichoptera Schulthess-Rechberg, 1894:71. Holotype ♂ (ETHZ).– S. Yemen.

Syn. *Fischeria [Bolivaria] dolichoptera* (Schulthess-Rechberg, 1894); Giglio-Tos 1916:21–22. *Eufischeriella [Bolivaria] dolichoptera* (Schulthess-Rechberg, 1894); Giglio-Tos 1927:485.

Distribution. Iran, Saudi Arabia, Yemen (Ehrmann 2002).

Distribution in Iran. Khorasan-e Jonubi, and Sistan-o Baluchistan (Beier 1956).

***Rivetina excellens* La Greca & Lombardo, 1982**

La Greca & Lombardo, 1982:374. Holotype ♂ (SMNS).– SE Iran.

Syn. *Rivetina excellens* Beier, 1956:73–74; Ehrmann, 2000:7, 2002:312; Otte & Spearman, 2005:235.

Distribution. Iran, Iraq, United Arab Emirates, Yemen (Beier 1956).

Distribution in Iran. Sistan-o Baluchistan (Beier 1956).

***Rivetina inermis* (Uvarov, 1923)**

Fischeria inermis Uvarov, 1923:721–722. Holotype ♂ (BMNH).– Saudi Arabia.

Distribution. Iran, Saudi Arabia, Yemen.

Distribution in Iran. Fars and Bushehr (Mirzaee & Sadeghi 2021).

***Rivetina rhombicollis* La Greca & Lombardo, 1982**

Rivetina rhombicollis La Greca & Lombardo, 1982:384. Holotype ♂ (NHMW).– Iran.

Distribution. Afghanistan, Iran, Pakistan.

Distribution in Iran. Sistan-o Baluchistan (La Greca & Lombardo 1982).

Genus *Bolivaria* Stål, 1877 [1/3 species]***Bolivaria brachyptera* (Pallas, 1773)**

Mantis brachyptera Pallas, 1773:728. Lectotype ♂ (ZMHB) (Alcohol).– Western Asia (Iran, Iraq).

Syn. *Mantis brachyptera* Pallas, 1773; Goeze 1778:32. *Mantis brachyptera* Pallas, 1773; Olivier 1792:635. *Mantis brachyptera* Pallas, 1773; Fischer-Waldheim, 1846:103. *Mantis brachyptera* Pallas, 1773; Kittary, 1849:447. *Mantis commutata* Fieber, 1853:95. *Mantis brachyptera* Pallas, 1773; Eversmann, 1859:125. *Iris pallasii* Saussure, 1869:65. *Iris* [*Mantis*] *brachyptera* (Pallas, 1773); Saussure, 1871:112–114. *Fischeria* [*Mantis*] *brachyptera* (Pallas, 1773); Saussure, 1874:18.

Fig. 3I

Distribution. Afghanistan, Armenia, Crete, Iran, Mongolia, Palestine, Turkey.

Distribution in Iran. Fars (Mirzaee & Sadeghi 2021), Guilan (Moradzadeh et al. 2021), Hamedan (Sadeghi & Sadeghi 2015), Isfahan (Beier 1956), Khuzestan (Ghahari & El-Den Nasser 2014), Kurdistan (Rouhani et al. 2015), Lorestan (Mirzaee & Sadeghi 2019), Mazandaran (Beier 1956), South Khorasan (Rabieh et al. 2016), Tehran (Beier 1956; Mirzaee & Pashaei Rad 2017), West Azerbaijan (Sakenin et al. 2011).

Genus *Microthespis* Werner, 1908 [1/2 species]***Microthespis dmitriewi* Werner, 1908**

Microthespis dmitriewi Werner, 1908:120. Holotype ♂, Paratype: ♂ (probably ZIN RAS).– Ethiopia.

Fig. 4C

Distribution. Bahrain, Egypt, Ethiopia, Iran, Jordan, Oman, Pakistan, Palestine/Israel, Saudi Arabia, Somalia, United Arab Emirates, Yemen.

Distribution in Iran. Fars (Mirzaee & Sadeghi 2021), Lorestan (Mirzaee & Sadeghi 2019), Sistan-o Baluchistan (Beier 1956).

Remarks. The fifth author, Leonid Anisyutkin, examined the Mantodea collection at ZIN RAS but was unable to locate the type material of this species. He noted that the type of *M. dmitriewi* may be in the collection, but is not properly designated. Therefore, further research is needed to identify the type material of this species.

Family Amelidae Westwood, 1889 [1/2 tribes]**Tribe Amelini Westwood, 1889** [1/3 genera]**Genus *Ameles* Burmeister, 1838** [1/24 species]***Ameles persa* Bolivar, 1911**

Fig. 3G

Ameles persa Bolivar, 1911:2–3. Holotype ♂, Allotype ♀ (MNMS).– Iran.

Distribution. Afghanistan, Armenia, Iran, Turkmenistan.

Distribution in Iran. Fars (Mirzaee & Sadeghi 2021), Isfahan (Ghahari & El-Den Nasser 2014), Kurdistan (Rouhani et al. 2015), Lorestan (Mirzaee & Sadeghi 2019), Sistan-o Baluchistan (Beier 1956), Khorasan-e Jonubi (Rabieh et al. 2016), Tehran (Mirzaee & Pashaei Rad 2017).

Family Eremiaphilidae Saussure, 1869 [2/4 subfamilies]

Subfamily Iridinae Westwood, 1889 [1/4 tribes]

Tribe Iridini Westwood, 1889 [1/3 genera]

Genus *Iris* Saussure, 1869 [4/14 species]

***Iris nana* Uvarov, 1930**

Fig. 3F

Iris nana Uvarov, 1930:632. Holotype ♂ (BMNH).– Iraq.

Syn. *Iris radians* Uvarov, 1931: 234–235.

Distribution. Afghanistan, Iran, Iraq, Pakistan, India.

Distribution in Iran. Fars (Uvarov 1938; Mirzaee & Sadeghi 2021), Tehran (Mirzaee & Pashaei Rad 2017).

***Iris oratoria* (Linné, 1758)**

Gryllus (*Mantis*) *oratorius* Linné, 1758:426. Holotype ♂ (UZIUI).– Africa.

Syn. *Mantis dentata* Goeze, 1778:36–37. *Mantis bella* Germar, 1817:6. *Mantis minima* Charpentier, 1825:91.

Distribution. Cosmopolitan.

Distribution in Iran. Fars (Mirzaee & Sadeghi 2021), Kerman (Ramme 1951), Kurdistan (Rouhani et al. 2015), Lorestan (Mirzaee & Sadeghi 2019), Khorasan-e Shomali (Sakenin et al. 2011), Khorasan-e Jonubi (Rabieh et al. 2016), Tehran (Mirzaee & Pashaei Rad 2017).

***Iris persa* Uvarov, 1922**

Iris persa Uvarov, 1923:723. Syntype ♂ (BMNH).– SW Iran.

Distribution. Iran.

Distribution in Iran. Fars (Ebner 1963), Bushehr (Uvarov 1922), Sistan-o Baluchistan (Beier 1956).

***Iris polystictica* (Fischer-Waldheim, 1846)**

Mantis polystictica Fischer-Waldheim, 1846:102. Holotype ♂ (probably ZIN RAS).– Russia.

Syn. *Iris tiflisina* Giglio-Tos, 1915:74. *Iris tiflisina* Giglio-Tos, 1915; Giglio-Tos, 1927:333. *Iris polystictica mongolica* Sjöstedt, 1933:2–3.

Distribution. Afghanistan, Armenia, Azerbaijan, Caucasus, China, Georgia, Iran, Kazakhstan, Russia, Tajikistan, Turkey, Turkmenistan, Uzbekistan.

Distribution in Iran. Ardabil (Morshedi Aghbolagh et al. 2012), Fars (Mirzaee & Sadeghi 2021), Tehran (Mirzaee & Pashaei Rad 2017), Mazandaran (Jamali & Mofidi-Neyestanak 2013).

Remark. The fifth author, Leonid Anisyutkin, examined the Mantodea collection at ZIN RAS but was unable to locate the type material of this species. He noted that the ZIN RAS collection holds numerous specimens of *I. polystictica*, including very old material. Therefore, further research is needed to identify the type that is not simple for old material.

Subfamily Eremiaphilinae Saussure, 1869 [1/2 genera]

Genus *Eremiaphila* Lefebvre, 1835 [2/64 species]

***Eremiaphila genei* Lefebvre, 1835**

Eremiaphila genei Lefebvre, 1835:486–489. Holotype ♂, Paratypes ♂♀ (MNHN).– Syria.

Syn. *Eremophila* [*Eremiaphila*] *genei* (Lefebvre, 1835); Burmeister, 1838:525. *Eremophila* [*Eremiaphila*] *zetterstedtii* (Lefebvre, 1835); Burmeister, 1838:525. *Eremiaphila burmeisteri* Saussure, 1871:252. *Eremiaphila genei* var. *laevipennis* Werner, 1905:386–387. *Eremiaphila hauensteini* Werner, 1905:387. *Eremiaphila burmeisteri* Saussure, 1871; Giglio-Tos, 1927:52.

Distribution. Afghanistan, Armenia, Egypt, Iran, Jordan, Lebanon, Palestine/Israel, Saudi Arabia, Syria, Turkey, Yemen.

Distribution in Iran. East Azerbaijan (Sakenin et al. 2011), Golestan (Sakenin et al. 2011), Isfahan (Ghahari & El-Den Nasser 2014), Khorasan-e Jonubi (Rabieh et al. 2016).

***Eremiaphila persica* Werner, 1905**

***Eremiaphila persica persica* Werner, 1905**

Eremiaphila persica persica Werner, 1905:388–389. Syntypes ♀♂♂ (ZIN RAS).– Iran.

Distribution. Azerbaijan, Iran, Iraq, Turkey.

Distribution in Iran. Isfahan (Beier 1956), Kerman (Ghahari & El-Den Nasser 2014), Khorasan-e Jonubi (Beier 1956).

Remarks. The online Mantodea species catalogue (Otte et al. 2023) states “syntypes male and female (ZIN St. Petersburg – holotype male, allotype female, Ehrmann 2002)”. However, examination of the ZIN RAS Mantodea collection revealed four specimens (one female and three males), all labelled as part of the type series and originally mentioned by Werner, with no specimen explicitly designated as holotype. Consequently, the current catalogue wording is somewhat misleading, because a formal holotype has not been fixed on any of the existing specimens.

***Eremiaphila persica sjostedti* Werner, 1930**

Eremiaphila persica sjostedti Werner, 1930:1–2. Holotype ♀, paratype ♂ (NHRS).– Iran.

Fig. 3C

Distribution. Iran, Pakistan.

Distribution in Iran. Bushehr (Beier 1956).

Family Toxoderidae Saussure, 1869 [2/4 subfamilies]

Subfamily Oxythespinae Giglio-Tos, 1916 [1/2 tribes]

Tribe Oxythespini Giglio-Tos, 1916 [4/7 genera]

Genus *Sinaiella* Uvarov, 1924 [1/4 species]

***Sinaiella azadi* Kolnegari & Schwarz, 2025**

Sinaiella azadi Kolnegari & Schwarz, 2025:102. Holotype ♂ (NHMB), paratype: juvenile (NMNHGR).– Iran.

Distribution. Armenia, Iran.

Distribution in Iran. Markazi (Kolnegari et al. 2025).

Genus *Severinia* Finot, 1902 [1/8 species]

***Severinia persica* Bolivar, 1913**

Oxythespis persica Bolivar, 1913:605–606. Holotype ♂, Allotype: ♀ (MNMS).– Iran.

Distribution. Iran, Afghanistan.

Distribution in Iran. Fars (Mirzaee & Sadeghi 2021), Isfahan (Ghahari & El-Den Nasser 2014), Lorestan (Mirzaee & Sadeghi 2019), Sistan-o Baluchistan (Beier 1956), Khorasan-e Jonubi (Rabieh et al. 2016).

Remarks. Kaltenbach (1982), in his contribution to the *Fauna of Saudi Arabia*, treated the species under the combination *Amblythespis persica* (Bolivar, 1913). Later, Lombardo (1991), in his revision “Remarks on the genus *Severinia*”, placed *Amblythespis* in synonymy with *Severinia*, thereby automatically transferring all its species to *Severinia* and making *Severinia persica* the correct combination under both taxonomic usage and zoological nomenclature. Despite this, the valid name *Severinia persica* is still frequently overlooked in the Mantodea literature, where the older combination as *Oxythespis persica* is often retained.

Genus *Lobothespis* La Greca & Lombardo, 1987 [1/1 species]***Lobothespis vignai* La Greca & Lombardo, 1987**

Lobothespis vignai La Greca & Lombardo, 1987:239–242. Holotype ♂ (INER).– Iran.

Distribution. Iran.

Distribution in Iran. Sistan-o Baluchistan (La Greca & Lombardo 1987).

Subfamily Toxoderinae Saussure, 1869 [1/4 tribes]**Tribe Aethalochroini Giglio-Tos, 1914** [1/3 genera]**Genus *Aethalochroa* Wood-Mason, 1877** [1/6 species]***Aethalochroa ashmoliana* (Westwood, 1841)**

Vates ashmoliana Westwood, 1841:272. Type ♂ (OUMNH).– India.

Syn. *Vates ashmoliana* Westwood, 1841; Westwood, 1843:52. *Popa* [*Vates*] *ashmoliana* (Westwood, 1841); Saussure, 1871: 161. *Arsacia* [*Vates*] *ashmoliana* (Westwood, 1841); Stål, 1877:75.

Distribution. Iran, India, Pakistan, Sri Lanka, and West Bengal.

Distribution in Iran. Sistan-o Baluchistan (Beier 1956).

Family Empusidae Burmeister, 1838 [2/2 subfamilies]**Subfamily Blepharodinae Giglio-Tos, 1919** [1/2 genera]**Genus *Blepharopsis* Rehn, 1902** [1/1 species]***Blepharopsis mendica* (Fabricius, 1775)**

Mantis mendica Fabricius, 1775:275. Types ♂♂, ♀♀ (ZMB).– Egypt.

Syn. *Gryllus monstrosus* Forskål, 1775:82. *Mantis mendica* Fabricius, 1775; Fabricius, 1781:347. *Mantis marmorata* Olivier, 1792:641. *Empusa* [*Mantis*] *mendica* (Fabricius, 1775); Latreille, 1807:90. *Gongylus* [*Mantis*] *mendica* (Fabricius, 1775); Thunberg, 1815:295. *Blepharis* [*Mantis*] *mendica* (Fabricius, 1775); Serville, 1831:47; *Mantis dilaticollis* Gistel, 1856:427. *Chersomantis picta* Gistel, 1856; Giglio-Tos, 1927:649.

Fig. 4A

Distribution. Afghanistan, Algeria, Canary Islands, Chad, Cyprus, Egypt, Ethiopia, NW India, Iran, Jordan, Lebanon, Libya, Mauritania, Morocco, Niger, Oman, Pakistan, Palestine/Israel, Somalia, Sudan, Tunisia, Turkey, United Arab Emirates, Uzbekistan, Yemen.

Distribution in Iran. Fars (Mirzaee & Sadeghi 2021), Lorestan (Mirzaee & Sadeghi 2019), Khorasan-e Shomali (Sakenin et al. 2011), Sistan-o Baluchistan (Ghahari & El-Den Nasser 2014, Beier 1956), Khorasan-e Jonubi (Rabieh et al. 2016).

Subfamily Empusinae Burmeister, 1838 [1/2 tribes]**Tribe Empusini Burmeister, 1838** [1/2 subtribes]**Subtribe Empusina Burmeister, 1838** [1/4 genera]**Genus *Empusa* Illiger, 1798** [3/11 species]***Empusa fasciata* Brullé, 1832**

Empusa fasciata Brullé, 1832:83. Probably lost.– Jerusalem.

Syn. *Empusa longicollis* Ramme, 1951:134–135.

Fig. 3A

Distribution. Afghanistan, Croatia, Cyprus, Egypt, Greece, India, Iran, Iraq, Jordan, Palestine/Israel, Romania, Russia, Slovenia, Turkey.

Distribution in Iran. Fars (Mirzaee & Sadeghi 2021), Guilan (Moradzadeh et al. 2021), Kurdistan (Rouhani et al. 2015), Khorasan-e Rzaei (Sakenin et al. 2011), Khorasan-e Jonubi (Rabieh et al. 2016), Tehran (Mirzaee & Pashaei Rad 2017).

Empusa hedenborgii Stål, 1877

Empusa hedenborgii Stål, 1877:77. Holotype ♂ (NHRS).– Sudan.

Syn. *Empusa* [*Mantis*] *pectinicornis* (Drury, 1770); Burmeister, 1838:546. *Empusa* [*Mantis*] *pectinicornis* (Drury, 1770); Stål, 1871:383. *Empusa egena* Charpentier, 1841; Rehn, 1901:288. *Empusa stollii* Saussure, 1871; Werner, 1907:247.

Distribution. Cameroon, North Somalia, Eritrea, Ethiopia, Senegal, Sudan, Yemen, Egypt, Iran, Saudi Arabia, United Arab Emirates.

Distribution in Iran. Khuzestan (Ghahari & El-Den Nasser 2014), Lorestan (Mirzaee & Sadeghi 2019), Sistan-o Baluchistan (Beier 1956), Tehran (Mirzaee & Pashaei Rad 2017).

Empusa pennicornis Lindt, 1978

Mantis pennicornis Pallas, 1773: 728. Type: ♂ (ZMB).– Syria.

Syn. *Mantis pennicornis* Pallas, 1773; Goeze, 1778:31–32. *Mantis pennicornis* Pallas, 1773; Gmelin-Linné, 1790:2055. *Mantis pennicornis* Pallas, 1773; Olivier, 1792:635. *Mantis pallasiana* Olivier, 1792:637–638. *Mantis pectinicornis* Drury, 1770; Lichtenstein, 1802:22–23. *Mantis pennicornis* Pallas, 1773; Stoll, 1813:30–31. *Gongylus marginatus* Thunberg, 1815:294. *Empusa* (*Empusa*) *orientalis* Burmeister, 1838:546–547. *Empusa orientalis* Burmeister, 1838; Fischer Waldheim, 1846:97. *Empusa* [*Mantis*] *pauperata* (Fabricius, 1781); Kittary, 1849:443. *Empusa orientalis* Burmeister, 1838; Fischer, 1853:132–133. *Empusa stollii* Saussure, 1871:188. *Empusa attenuata* Ramme, 1951:135–136.

Distribution. Afghanistan, China, Georgia, Iran, Iraq, Kazakhstan, Russia, Tajikistan, Turkey, Turkmenistan, Ukraine, Uzbekistan.

Distribution in Iran. Ardabil (Sakenin et al. 2011), Fars (Mirzaee & Sadeghi 2019), Kurdistan (Ghahari & El-Den Nasser 2014), Sistan-o Baluchistan (Beier 1956), Khorasan-e Jonubi (Beier 1956), Tehran (Mirzaee & Pashaei Rad 2017).

Family Mantidae Latreille, 1802 [2/11 subfamilies]

Subfamily Mantinae Latreille, 1802 (1/3 genera)

Genus *Mantis* Linnaeus, 175 [1/14 species]

Mantis religiosa Linné, 1758

Gryllus (*Mantis*) *religiosus* Linné, 1758:426. Holotype ♂, Paratypes: ♀♀♀ (LSUK).– Africa.

Syn. *Gryllus religiosus* Linné, 1758; Scopoli 1763:105. *Mantis religiosa*: Linné 1767:690. *Mantis oratoria* Fabricius, 1775:276–277. *Mantis sancta* Fabricius, 1787:228. *Mantis sancta* Fabricius, 1787; Olivier 1792:628–629. *Mantis religiosa* var. *striata* Fabricius, 1793:20. *Mantis oratoria* Fabricius, 1775; Lichtenstein 1802:28–29. *Mantis maroccana* Thunberg, 1815:287–299. *Mantis pia* Audinet-Serville, 1839:193. *Mantis prasina* Audinet-Serville, 1839:195. *Mantis radiata* Motchoulsky: Fischer-Waldheim 1846:101. *Mantis capensis* Saussure, 1872:46–47. *Mantis capensis*: Stål 1877:60–61; *Mantis religiosa* var. *major* Gerstaecker, 1873:12. *Mantis prasina* Audinet-Serville, 1839: Stål 1877:61. *Mantis pia* Audinet-Serville, 1839; Kirby 1899:348. *Mantis macroura* Brunner de Wattenwyl, 1882:60. *Mantis carinata* Cosmovici, 1888:172–173. *Mantis religiosa flava* Padewieth, 1900:20. *Mantis religiosa brunnea* Padewieth, 1900:20.

Distribution. Africa, Asia, Europe, North America (introduced) (Ehrmann 2002).

Distribution in Iran. East Azerbaijan (Sakenin et al. 2011), Fars (Sadeghi & Sadeghi 2015; Mirzaee & Sadeghi 2021; Ghahari & El-Den Nasser 2014), Hormozgan (Ghahari & El-Den Nasser 2014), Kurdistan (Rouhani et al. 2015; Ghahari & El-Den Nasser 2014), Lorestan (Mirzaee & Sadeghi 2019), Mazandaran (Beier 1956), Khorasan-e Razavi (Sakenin et al. 2011), Khorasan-e Jonubi (Rabieh et al. 2016), Sistan-o Baluchistan (Beier 1956), Tehran (Mirzaee & Pashaei Rad 2017), West Azerbaijan (Sakenin et al. 2011).

Subfamily Hierodulinae Brunner von Wattenwyl, 1893 [1/2 tribes]

Tribe Hierodulini Brunner von Wattenwyl, 1893 [1/13 genera]

Genus *Hierodula* Burmeister, 1838 [2/104 species]

***Hierodula coarctata* Saussure, 1869**

Hierodula coarctata Saussure. 1869:67. Holotype ♂ (MHNG).– India.

Syn. *Hierodula (Hierodula) macrostigmata* Deeleman-Reinhold, 1957:57–58.

Distribution. Iran, India, Java, Nepal, Pakistan.

Distribution in Iran. Hormozgan (Deeleman-Reinhold 1957), Sistan-o Baluchistan (Mirzaee et al. 2026).




Figure 3. Representative Mantodea of Iran: **A.** *Empusa fasciata*, female from Bushehr: Jam; **B.** *Elaea richteri*, male from Sistan-o Baluchistan; **C.** *Eremiaphila persica*, female from Bushehr; **D.** *Holaptilon khozestani*, male from Khuzestan; **E.** *Armene pusilla*, male from Mashahd; **F.** *Iris nana*, female from Bushehr: Kangan; **G.** *Ameles persa*, female from Fars; **H.** *Severinia* sp., subadult female from Fars; **I.** *Bolivarina brachyptera*, female from Bushehr. Photos by: Zohreh Mirzaee (A–D, F–I); Mehdi Ghafarnia (E).



Figure 4. Representative Mantodea of Iran: **A.** *Blepharopsis mendica*, female from Bushehr: Jam; **B.** *Hierodula tenuidentata*, female from Fars; **C.** *Microthespis dmitriewi*, female from Bushehr; **D.** *Rivetina* sp., female from Bushehr: Jam; **E.** *Nilomantis floweri*, male from Hormozgan; **F.** *Amorphoscelis pantherina*, male from Fars; Photos by: Zohreh Mirzaee (A-E); Mohsen Kiani (F).

Hierodula tenuidentata Saussure, 1869

Hierodula tenuidentata Saussure, 1869:68. Holotype ♂ (MHNG).– India.

Syn. *Hierodula* [*Mantis*] *simulacrum* (Fabricius, 1793); Westwood 1889:34. *Hierodula heterodera* Westwood, 1889:34. *Sphodromantis* [*Hierodula*] *tenuidentata* (Saussure, 1869); Kirby 1904:244. *Hierodula westwoodi* Kirby, 1904:247.

Fig. 4B

Distribution. Borneo, Bosnia and Herzegovina, Bulgaria, China, Germany, India, Iran, Italy, Kazakhstan, Mongolia, Nepal, Romania, Serbia, Sunda Islands, Tajikistan, Turkmenistan, Turkey.

Distribution in Iran. North, Northeast, Central, and Southwest of Iran (Mirzaee et al. 2026).

Order Phasmatodea Jacobson & Bianchi, 1902

Suborder Euphasmatodea Bradler, 1999

Superfamily Bacilloidea Brunner von Wattenwyl, 1893 [1/3 families]

Family Bacillidae Brunner von Wattenwyl, 1893 [1/9 tribes]

Tribe Gratidiini Cliquennois, 2005 [2/16 genera]

Genus *Adelungella* Brunner von Wattenwyl, 1907 [1/1 species]

***Adelungella insignis* Brunner von Wattenwyl, 1907**

Adelungella insignis Brunner von Wattenwyl, 1907:229. Syntypes (2): ♀♀ (ZIN).– Iran.

Distribution. Iran.

Distribution in Iran. Kerman (Brunner von Wattenwyl 1907).

Remark. So far only documented only from the type locality, but the distribution is assumed to be wider.

Genus *Clonaria* Stål, 1875 [4/127 species]

***Clonaria inconspicua* (Brunner von Wattenwyl, 1907)**

Gratidia inconspicua Brunner von Wattenwyl, 1907:223. Syntypes (8): 2♀♀, 1♂ (NHMW). Uzbekistan: Buchara; 2♂♂ (ZIN). Iran: Kerman; 1♂ (ZIN). Iran: Khorasan; 1♂ (ZIN). Iran: Road from Trauscaspian (Turkmenia) to Kuchan (Persia **Syn.** Iran); 1♂ (ZIN). Uzbekistan: Buchara.

Syn. *Ramulus inconspicua* (Brunner von Wattenwyl, 1907); Brock, 1991:39.

Fig. 5A

Distribution. Turkmenistan, Iran.

Distribution in Iran. Kerman, Khorasan-e Razavi (Brock et al. 2025).

***Clonaria predtetshenkyi* (Bey-Bienko, 1946)**

Gratidia predtetshenkyi Bey-Bienko, 1946:74. Holotype ♂ (ZIN RAS). Iran.

Syn. *Ramulus predtetshenkyi* (Bey-Bienko, 1946); Brock, 1991:39.

Distribution. Iran.

Distribution in Iran. Qazvin (Bey-Bienko 1946).

Remark. So far only documented only from the type locality, but the distribution is assumed to be wider.

***Clonaria spinulosa* (Brunner von Wattenwyl, 1907)**

Gratidia spinulosa Brunner von Wattenwyl, 1907:228. Holotype ♀ (ZIM). Iran.

Syn. *Ramulus spinulosa* (Brunner von Wattenwyl, 1907); Brock, 1991:39.

Distribution. Iran.

Distribution in Iran. Sistan-o Baluchistan (Brunner von Wattenwyl 1907).

Remark. So far only documented only from the type locality, but the distribution is assumed to be wider.

***Clonaria uvaroviana* (Mistshenko, 1937)**

Gratidia uvaroviana Mishchenko, 1937:119. Holotype ♀ (ZIN RAS). Turkmenistan: V. Keltetshinar; paratype ♂ (ZIM). Iran: Khorasan.

Syn. *Ramulus uvaroviana* (Mishchenko, 1937); Brock, 1991:39.

Distribution. Turkmenistan, Iran.

Distribution in Iran. Khorasan (Brock 2007).

Remark. So far only been documented in Iran from the type localities, but the distribution is assumed to be wider.




Figure 5. Representative Phasmatodea of Iran, Most likely the majority are new species and hardly any doubled species: **A.** Adult female *Clonaria* cf. *inconspicua* from Mashhad, photo credit by: Amin Ghaffarian; **B.** Adult male of *Clonaria* sp. from Yazd, photo credit by: Mehdi Zare Khormizi; **C.** immature female *Adelungella* sp. from Semnan photo credit by: Alireza Salehi; **D.** Young nymph *Clonaria* sp. from Semnan, photo credit by: Alireza Salehi; **E.** Adult female *Adelungella* sp. from, photo credit by: Milad; **F.** Adult female *Clonaria* sp. from Yazd, photo credit by: Hamid Dehoi; **G.** Adult male *Clonaria* sp. from Isfahan, **H.** Adult female *Clonaria* sp. from Fars, photo credit by: Amir Hossein Abdullahi; **I.** Adult female *Adelungella* sp. from Yazd, photo credit by: Moosa Ghorbaniorjanli; **J.** Adult female *Clonaria* sp. from Golestan; **K.** Nymph *Clonaria* sp. from Fras: Shiraz, photo credit by: Mobin.

DISCUSSION

The study of three polyneopteran orders in Iran reveals a diverse yet unevenly explored insect fauna compared to global standards. This disparity is problematic given Iran's biogeographic position at the intersection of the Palaearctic, Afrotropical, and Oriental realms, which suggests a potentially richer and under-documented fauna (Zohary 1973; Olson et al. 2001). Based on current knowledge, approximately 23% of Iranian Blattodea species, 33% of Mantodea, and 60% of Phasmatodea are known only from Iran.

Research on Blattodea in Iran has been modest but is improving with molecular data and checklists. Although several cockroach and termite species have been reported, sampling biases in remote and arid provinces likely underestimate true diversity. Termite taxonomy is hindered by cryptic species complexes, while synanthropic species such as *Blattella germanica* and *Periplaneta* spp. are well-documented in urban areas. In contrast, xeric and soil-dwelling taxa remain poorly collected and studied (Bey-Bienko 1950; Ghayourfar 1998; Hashemi-Aghdam & Oshaghi 2015). Iran's distribution patterns and endemism emphasize its biogeographic importance for polyneopterans (Mehrabian & Rad 2021; Moradi et al. 2024). Species-rich termite genera that include several Iranian endemics, such as *Microcerotermes* and *Amitermes*, also highlight local diversification, though many cockroach taxa once thought monotypic have since revealed hidden species upon further investigation (Mirzayans 1986). Widespread cosmopolitan species like *Periplaneta americana* and *Blattella germanica* contrast sharply with lesser-known xerophilous and subterranean species.

Over the past two decades, Mantodea research has notably expanded due to increased faunistic and systematic studies by Iranian entomologists, whereas investigations into Blattodea and Phasmatodea remain limited in both depth and width. A significant knowledge gap persists for all three groups. Early studies were based on opportunistic collections, creating a fragmented understanding of Mantodea diversity (Uvarov 1938; Beier 1956; Kaltenbach 1982; Ehrmann 2002), a pattern even more pronounced among Blattodea and Phasmatodea (Mirzayans 1986; Hashemi-Aghdam & Oshaghi 2015; Brock & Büscher 2022). Despite recent progress in Mantodea taxonomy and distribution (Mirzaee & Pashaei Rad 2017; Kolnegari & Vafaei-Shoushtari 2018; Mirzaee & Sadeghi 2019, 2021; Moradzadeh et al. 2021; Kolnegari 2023; Mirzaee et al. 2023, 2024a, 2024b; Kolnegari et al. 2025), vast regions of Iran, particularly deserts and mountain zones, remain insufficiently surveyed. The continuous discovery of new species and records highlights the need for comprehensive field studies in unexplored provinces and ecologically unique habitats (Olson et al. 2001; Mirzaee et al. 2023, 2024a, 2024b). On iNaturalist, there are currently 242 and 182 observations from Iran of Mantodea (iNaturalist:48112) and Blattodea (iNaturalist:81769), respectively, across the whole country. Within Mantodea, widespread species like *Blepharopsis mendica* exhibit broad ecological tolerance, while microendemic taxa, such as species of *Holaptilon* (Beier 1964), are highly localized and considered endangered due to their narrow ranges and vulnerability to habitat alteration (Mirzaee et al., 2024b). *Holaptilon* species are particularly useful as bioindicators, reflecting habitat stability and ecological integrity (Battiston et al. 2020).

Phasmatodea research is minimal, relying on historical records (Brunner von Wattenwyl 1907; Bey-Bienko 1946; Otte & Brock 2005; Brock & Büscher 2022). Only a handful of species, five across two genera, are known, representing a small fraction of the likely existing fauna. Additionally, yet-undetected taxa are expected in the Hyrcanian forests, Zagros woodlands, and montane refugia, where sampling remains limited, and the cryptic behavior and camouflage of phasmids further complicate detection. Throughout Iran, documented observations are scarce for Phasmatodea, although their diversity is expected to be somewhat high. Examples from other countries have recently shown that sampling in such undersampled regions has frequently multiplied documented species numbers in a short time (Brock & Büscher 2022). There are currently twelve observations of phasmids from Iran on iNaturalist (iNaturalist:47198), which are scattered across the whole country, and excluding biodiversity-rich areas, none of those have been identified to species or genus due to a deficiency of taxonomical data. Most likely, the actual phasmid species-richness is drastically underrepresented across all documentations for Iran. This is further supported by molecular evidence on gratidiine stick insects in general, which appear to be rather diverse, but often undescribed or poorly diagnosed (Bank & Bradler 2022).

Taxonomic complexity across Blattodea, Mantodea, and Phasmatodea stems from morphological variability, outdated descriptions, and limited molecular data. In cockroach and termite taxonomy, suffer from synonymies and cryptic diversity, with gene markers like *cox1* proving valuable for species delimitation (Ren et al. 2025). Similarly, mantis genera such as *Rivetina*, *Iris*, and *Ameles* present particular challenges, requiring integrative approaches that combine morphology, molecular markers, and ecology (Mirzaee et al., 2023, 2024a, 2024b). For Phasmatodea, resolving synonymies and lost type material demands modern DNA-based revisions and museum re-examinations. Furthermore, the basis for approaching the taxonomy of phasmids from Iran is conflicted by the resolution of the phylogenetic relationships in regard to Gratidiini (e.g., Büscher et al. 2024). Presently, the taxonomic treatment of Bacillidae in general and Gratidiini in particular is challenged by phylogenetic studies using molecular sequences, indicating that both Bacillidae and Gratidiini are not monophyletic and several lineages within Gratidiini are actually not directly related to each other and nested in different clades (Bank & Bradler 2022). This particularly affects the two genera represented in Iran, both likely representatives of a subordinate lineage within Clitumninae (Gao et al. 2025). The Iranian taxa are, so far, not included in any phylogenetic analysis. However, they are most likely closely related to other West Asian *Clonaria* spp. analysed in Gao et al. (2025) that form a subordinate clade in Clitumninae together with the genera *Sceptrophasma* Brock & Seow-Choen, 2000, and *Macellina* Uvarov, 1940.

Several Iranian provinces, including Chaharmahal and Bakhtiari, Kohgiluyeh va Boyer-Ahmad, Qazvin, and Zanjan, remain little studied despite hosting ecologically diverse landscapes. Western and southern mountains, along with semi-desert oases and plateaus, hold great promise for new discoveries

(Noori et al. 2024). Southern and highland regions especially require renewed field and molecular efforts to uncover hidden diversity (Mirzayans 1986; Hashemi-Aghdam & Oshaghi 2015). Cockroach species such as *P. americana* and *B. germanica* exemplify the role of synanthropic fauna in biosecurity risks. Although stick insects are less likely to disperse through trade, the ornamental plant and pet industries warrant monitoring for possible introductions, as frequently observed in other countries (e.g., Great Britain: Lee, 1993). Currently, all mantis species recorded in Iran are native, with no established alien taxa. Nevertheless, global trends of exotic mantis introductions linked to human trade raise the possibility of future invasions (Battiston et al. 2018, 2020; Moulin 2020; Shcherbakov & Govorov 2020; Sevgili & Yilmaz 2022).

Habitat destruction is a major threat, particularly for mantises specialized to forest margins or arid steppe habitats and phasmids in general. Agricultural intensification, urban expansion, and deforestation, especially in the Zagros and Alborz ranges, have caused widespread habitat loss (Jowkar et al. 2016). Overgrazing and desertification further reduce viable habitats for specialist taxa (Valizadeh 2010; Mahmoudi et al. 2011). While these pressures threaten native Blattodea, Mantodea, and Phasmatodea, they simultaneously create opportunities for synanthropic and invasive species. Climate change poses an additional, long-term threat to Iranian biodiversity. Projected temperature and precipitation shifts could particularly endanger montane and forest-adapted species, while desert-dwelling taxa may show greater resilience. These climatic shifts could drive range contractions, endanger endemic populations, and reshape faunal communities (La Sorte & Jetz 2010; Ansari et al. 2023). This risk is particularly high for phasmids, as desert species are rare (Brock & Büscher 2022) and temperature increase likely poses major threats to the majority of stick insects (Grosman 2022).

This study emphasizes the need for coordinated efforts to preserve Iran's polyneopteran diversity. It highlights the importance of systematic field surveys in poorly studied regions and the adoption of integrative taxonomic approaches that merge morphological, molecular, and ecological data. Additionally, revisiting and digitizing historical museum collections is advised to enhance taxonomic resolution. On the policy level, it is recommended to prioritize the conservation of critical habitats by integrating climate change scenarios, particularly for microendemic and specialist species.

AUTHOR'S CONTRIBUTION

The authors confirm their contribution to the paper as follows: Z. Mirzaee: Study conception and design, data analysis, interpretation of results, drafting the manuscript, visualization; T.H. Büscher: revision of Phasmatodea part, data analysis, drafting the manuscript; D. Evangelista: revision of Blattodea part, drafting the manuscript; Paul Eggleton: revision of Blattodea part; L. Anisyutkin: revision of Blattodea part, data analysis, drafting the manuscript; Frédéric Legendre: revision of Blattodea part, data analysis, drafting the manuscript. The authors read and approved the final version of the manuscript.

FUNDING

No funding was used for this study. Participation by Leonid Anisyutkin was supported in the framework of the state research project 125012901042-9 (Systematization of insect diversity in taxonomic, ecophysiological, and evolutionary aspects).

AVAILABILITY OF DATA AND MATERIAL

Depositories for the specimens listed in this study are listed in the Material and methods section.

ETHICS APPROVAL AND CONSENT TO PARTICIPATE

This study only included arthropod material, and all required ethical guidelines for the treatment and use of animals were strictly adhered to in accordance with international, national, and institutional regulations. No human participants were involved in any studies conducted by the authors for this article.

CONSENT FOR PUBLICATION

Not applicable.

CONFLICT OF INTERESTS

The authors declare that there is no conflict of interest regarding the publication of this paper.

GENERATIVE AI STATEMENT

The authors declare that no generative AI tools were used in the preparation of this manuscript.

ACKNOWLEDGMENTS

We would like to thank Yaser Bakhshi, Alireza Salehi, Hamid Dehoi, Mehdi Ghafarnia, Amin Ghafarian, Mehdi Zare Khormizi, Mohsen Kiani, Amir Hossein Abdullahi, Zahed Aram, Kian Saidabadi, Moosa Ghorbaniorjanli, Houman Doroudi, and Daryoosh Mansouri for generously sharing their photographs, which greatly enriched this work. Sincere thanks are also extended to Evgeny Shcherbakov (Lomonosov Moscow State University, Russia) for his thoughtful and constructive comments on the Mantodea section and Paul D. Brock (Natural History Museum, London, UK) for his comments on the Iranian Phasmatodea.

REFERENCES

- Abedi, F. & Dayer, M. (2006) Evaluation of the effect of the fungus *Metarhizium anisopliae*, as a biological control agent, on German cockroaches *Blattella germanica*. *Modares Journal of Medical Sciences: Pathobiology*, 8 (1), 31–36 [In Persian]
- Akhtar, M.S. (1974) New termites from Pakistan. *Biologia* (Lahore), 20 (1), 23–61.
- Al-Alawy, S.A., Abdul-Rassoul, M.S. & Al-Azawi, A.F. (1990) Description of new species of termites (Insecta, Isoptera) from Iraq. *Bulletin of the Iraq Natural History Museum*, 8 (3), 25–33.
- Anisyutkin, L.N. & Perkovsky, E.E. (2023) Two new cockroaches (Dictyoptera: Ectobiidae) from Rovno amber. *Biologia*, 78 (6), 1721–1730. <https://doi.org/10.1007/s11756-023-01333-6>
- Ansari, A., Ghorbanpour, M., Kazemi, A. & Kariman, K. (2023) Ecological assessment of Iran's terrestrial biomes for wildlife conservation. *Scientific Reports*, 13 (1), 17761. <https://doi.org/10.1038/s41598-023-45120-4>
- Austin, J.W. (2006) *Phylogeny and Genetic Variation of Reticulitermes (Isoptera: Rhinotermitidae)*. Ph.D. Thesis, Middle East Technical University, Ankara, Turkey, 116 pp.
- Bank, S. & Bradler, S. (2022) A second view on the evolution of flight in stick and leaf insects (Phasmatodea). *BMC Ecology & Evolution* 22, 62. <https://doi.org/10.1186/s12862-022-02018-5>
- Basset, Y. (2001) Invertebrates in the canopy of tropical rain forests: How much do we really know? *Plant Ecology*, 153, 87–107. https://doi.org/10.1007/978-94-017-3606-0_8
- Battiston, R., Picciau, L., Fontana, P. & Marshall, J. (2010) *Mantids of the Euro-Mediterranean Area*. Verona: World Biodiversity Association, Verona. 239 pp.
- Battiston, R., Leandri, F., di Pietro, W. & Andria, S. (2018) The Giant Asian Mantis *Hierodula tenuidentata* Saussure, 1869 spreads in Italy: a new invasive alien species for the European fauna? (Insecta, Mantodea). *Biodiversity Journal*, 9 (4), 399–404. <https://doi.org/10.31396/Biodiv.Jour.2018.9.4.399.404>
- Battiston, R., Amerini, R., Di Pietro, W., Guariento, L.A., Bolognin, L. & Moretto, E. (2020) A new alien mantis in Italy: Is the Indochina mantis *Hierodula patellifera* chasing the train for Europe? *Biodiversity Data Journal*, 8, e50779. <https://doi.org/10.3897/BDJ.8.e50779>
- Beccaloni, G. (2024) *Cockroach Species File*. Available from <https://cockroach.speciesfile.org/> [Accessed November 27, 2025]
- Beier, M. (1956) Mantiden aus dem Iran 1954 (Orthoptera). Ergebnisse der Entomologischen Reisen Willi Richter, Stuttgart, im Iran 1954 und 1956, Nr. 2. *Jahreshefte des Vereins für vaterländische Naturkunde in Württemberg*, 111, 68–75.
- Beier, M. (1964) Ein neues Mantiden-Genus aus Israel. *Israel Journal of Zoology*, 13, 184–186.
- Beier, M. (1968) 12. Mantodea (Fangheuschrecken). In: Helmcke, J.G., Starck, D. & Wermuth, H. (eds) *Handbuch der Zoologie. IV. Band: Arthropoda, 2: Insecta*. Walter de Gruyter & Co., Berlin, pp. 1–47.
- Belyaeva, N.V. (1974) About species belonging [to] the Turkmenian termites *Amitermes rhizophagus* sp. n. (Isoptera: Termitidae). In: Zolotarev, E.K. (ed) *Termites* (collected articles). Transactions of the Entomological Division 5, Biodeterioration Research Laboratory, Faculty of Biology, Moscow Lomonosov University Publishing House, Moscow, pp. 62–73. [in Russian]
- Bey-Bienko, G.Y. (1946) A new species of the genus *Gratidia* Stål (Phasmoidea) from Iran, with a list of Palearctic species of this genus. *Proceedings of the Royal Entomological Society of London. Series B, Taxonomy*, 15 (5–6), 74–76. <http://doi.org/10.1111/j.1365-3113.1946.tb00826.x>
- Bey-Bienko, G.Y. (1950) Cockroach insects. Fauna of the USSR. New Series 40. Nauka Publishers, Leningrad and Moscow. 343 pp. [in Russian]
- Bignell, D.E. & Eggleton, P. (2000) Termites in ecosystems. In: Abe, T., Bignell, D.E. & Higashi, E. (eds) *Termites: Evolution, Sociality, Symbioses, Ecology*. Springer, Dordrecht, Netherlands, pp. 363–387. https://doi.org/10.1007/978-94-017-3223-9_17
- Babmorad, M., Ghayourfar, R., Saleheh Shoshtari, M.H. & Zeinali, S. (2014) The first report of *Postelectrotermes pasniensis* Akhtar (Isop.: Kalotermitidae) from south of Iran. *Journal of Field Crop Entomology*, 6 (1), 59–61.
- Baniardalani, M., Nourjah, N. & Changani, F. (2006) Susceptibility of cockroaches to Lambdacyhalothrin and Cypermethrin in hospitals of Tehran University of Medical Science. *Journal of Qazvin University of Medical Sciences*, 10 (1), 95–100. [In Persian]
- Bohn, H. (2013) Another species of European Ectobiinae travelling north—the new genus *Planuncus* and its relatives (Insecta: Blattodea: Ectobiinae). *Arthropod Systematics & Phylogeny*, 71 (3), 139–168. <https://doi.org/10.3897/asp.71.e31774>

- Bohn, H. (2019) Revision of the genus *Dziriblatia* Chopard, 1936 (Blattodea, Ectobiidae, Ectobiinae) from North Africa, Spain, and the Macaronesian islands. I. The nine subgenera of the genus. *Zootaxa*, 4610 (1), 1–73. <https://doi.org/10.11646/zootaxa.4610.1.1>
- Bohn, H. (2024) The spine armament of the legs as an important means for the characterisation of the genera of Corydiinae and their relationships (Blattodea, Corydiidae). *Zootaxa*, 5482 (1), 1–79. <https://doi.org/10.11646/zootaxa.5482.1.1>
- Bohn, H. & Chladek, F. (2011) Revision of the maculata-group of *Phyllodromica*: species from Central Europe (Insecta: Blattodea: Blattellidae: Ectobiinae). *Arthropod Systematics & Phylogeny*, 69, 3–54. <https://doi.org/10.3897/asp.69.e31735>
- Bolivar, I. (1911) Description de cinq espèces nouvelles d'Orthoptères [Description of five new species of Orthoptera]. *Bulletin de la Société des Amis des Sciences Naturelles de Rouen*, 1911, 1–3.
- Bolivar, I. (1913) Orthoptères. Expedition to the central western Sahara by Ernst Hartert. *Novitates Zoologicae*, 20, 603–607.
- Brock, P.D. (1991) *Stick-insects of Britain, Europe and the Mediterranean*. The Amateur Entomologists' Society. London. 50 pp.
- Brock, P.D. (2007) The stick-insects (Phasmatodea) of the Crimea. *Zootaxa*, 1398 (1), 51–56.
- Brock, P.D. & Büscher, T.H. (2022) *Stick and Leaf-insects of the World: Phasmids*. NAP éditions, NAP éditions, Verrières-le-Buisson, France. 612 pp.
- Brock, P.D., Büscher, T.H. & Baker, E. (2025) *Phasmida Species File Online*. Available from: <https://phasmida.speciesfile.org/> [Accessed October 13, 2025]
- Brunner von Wattenwyl, C. (1878) Naturwissenschaftliche Beiträge zur Kenntnis der Kaukasusländer auf Grund seiner Sammelbeute. *Naturwissenschaftliche Beiträge zur Kaukasus*, 5, 87–90.
- Brunner von Wattenwyl, C. (1907) Die Insektenfamilie der Phasmiden. II. Phasmidae Anareolatae (Clitumnini, Lonchodini, Bacunculini). *Phasmiden*, 2, 181–340.
- Büscher, T.H., Reck, L.M. & Gorb, S.N. (2024) Functional surface structures on the eggs of stick and leaf insects (Insecta: Phasmatodea). *Zoologica*, 166, 1–340.
- Chhotani, O.B. & Bose, G. (1979) Insects of Saudi Arabia. Isoptera. *Fauna of Saudi Arabia*, 1, 75–83.
- Chopard, L. (1921) Report on the Orthoptera of Mesopotamia and Persia collected by Messrs. P.A. Buxton M.A. and W.E. Evans, B. So. *Journal of the Bombay Natural History Society*, 27 (4), 47–59.
- Chopard, L. (1929) Orthoptera Plaeartica critica. VII. Les Polyphagiens de la faune paléarctique (Orth., Blatt.). *Eos*, 5, 215–358.
- Dedeine, F., Dupont, S., Bagnères, A.-G., Mantovani, B., Dronnet, S. & Bouchon, D. (2016) From speciation to introgressive hybridization: the phylogeographic structure of an island subspecies of the termite *Reticulitermes lucifugus*. *Molecular Phylogenetics and Evolution*, 94, 267–277.
- Deeleman-Reinhold, C.L. (1957) Notes sur les Mantides du Musée de Leyde avec la description d'un nouveau genre et de quelques espèces nouvelles. *Zoologische Mededeelingen*, 35, 55–67.
- Deng, W., Luo, X., Ho, S. Y., Liao, S., Wang, Z. & Che, Y. (2023) Inclusion of rare taxa from Blattidae and Anaplectidae improves phylogenetic resolution in the cockroach superfamily Blattoidea. *Systematic Entomology*, 48 (1), 23–39. <https://doi.org/10.1111/syen.12560>
- Deng, W.B., Drager, K., Li, X. R., Han, W., Djernæs, M., Wang, Y.S., Martin, J., Schwartz, J., Zhang, Z.Z., Kovacs, T.G.L., Lo, N.T., Helleman, S., Che, Y.L., Bourguignon, T., Wang, Z.Q. & Evangelista, D. (2026) Blattodea systematics in the genomic age. *Insect Systematics and Diversity*, 10 (1), ixaf042. <https://doi.org/10.1093/isd/ixaf042>
- Desneux, J. (1906) Variétés termitologiques. *Annales de la Société Entomologique de Belgique*, 49 (12), 336–360.
- Djernæs, M. (2018) Biodiversity of Blattodea-the cockroaches and termites. *Insect biodiversity: science and society*, 2, 359–387. <https://doi.org/10.1002/9781118945582.ch14>
- Djernæs, M. & Muriene, J. (2022) Phylogeny of Blattoidea (Dictyoptera: Blattodea) with a revised classification of Blattidae. *Arthropod Systematics and Phylogeny*, 80, 209–228. <https://doi.org/10.3897/asp.80.e75819>
- Doroodgar, A., Arbabi, M. & Asadi, M.A. (2006) Survey on parasitic and fungal agents of Hospital cockroaches in Kashan-2002. *Feyz Medical Sciences Journal*, 10 (1), 1–6.
- Ebner, V.R. (1963) Orthopteroidea und Dictyoptera der Österreichischen Iranexpedition 1949/1950. *Annalen des Naturhistorischen Museums in Wien*, 67, 395–403.
- Ehrmann, R. (2002) *Mantodea: Gottesanbeterinnen der Welt*. Natur und Tier-Verlag, Münster, Germany. 519 pp.
- Enayati, A.A. & Motevali, H.F. (2007) Biochemistry of pyrethroid resistance in German cockroach (Dictyoptera, Blattellidae) from hospitals of Sari, Iran. *Iranian Biomedical Journal*, 11 (4), 251–258.
- Evangelista, D.A., Wipfler, B., Béthoux, O., Donath, A., Fujita, M., Kohli, M.K., Legendre, F., Liu, S., Machida, R., Misof, B., Peters, R.S., Podsiadlowski, L., Rust, J., Schuette, K., Tollenaar, W., Ware, J.L., Wappler, T., Zhou, X., Meusemann, K. & Simon, S. (2019) An integrative phylogenomic approach illuminates the evolutionary history of cockroaches and termites (Blattodea). *Proceedings of the Royal Society B*, 286 (1895), 20182076. <https://doi.org/10.1098/rspb.2018.2076>
- Evangelista, D., Simon, S., Wilson, M.M., Kawahara, A.Y., Kohli, M.K., Ware, J.L., Wipfler, B., Béthoux, O., Grandcolas, P. & Legendre, F. (2021) Assessing support for Blaberoidea phylogeny suggests optimal locus quality. *Systematic Entomology*, 46 (1), 157–171. <https://doi.org/10.1111/syen.12454>
- Evangelista, D.A., Nelson, D., Varadínová, Z.K. & Legendre, F. (2023) Phylogenomics and deep convergence in cockroach hind-wing morphology. *Organisms Diversity & Evolution*, 23 (4), 929–940. <https://doi.org/10.1007/s13127-023-00609-8>

- Evangelista, D.A., Nelson, D., Varadínová, Z.K., Kotyk, M., Rousseaux, N., Shanahan, T., Grandcolas, P. & Legendre, F. (2024) Phylogenomic analyses of Blattodea combining traditional methods, incremental tree-building, and quality-aware support. *Molecular Phylogenetics and Evolution*, 200, 108177. <https://doi.org/10.1016/j.ympev.2024.108177>
- Fahiminia, M., Paksa, A., Zarei, A., Shams, M., Fazlzadeh-davil, M., Bakhtiari, H. & Norouzi, M. (2010) Survey of optimal methods for the control of cockroaches in sewers of Qom City. *Iranian Journal of Health and Environment*, 3 (1), 19–26 [In Persian].
- Fakoorziba, M.R., Eghbal, F., Hassanzadeh, J. & Moemenbellah-fard, M.D. (2010) Cockroaches (*Periplaneta americana* and *Blattella germanica*) as potential vectors of the pathogenic bacteria found in nosocomial infections. *Annals of Tropical Medicine & Parasitology*, 104 (6), 521–528. <https://doi.org/10.1179/136485910X12786389891326>
- Fallahzadeh, M., Asadi, R. & Talebi, A.A. (2008) The first record of *Anastatus (Anastatus) tenuipes* (Hym.: Eupelmidae), a parasitoid of *Bellatella germanica* (Blattaria: Blattellidae) in Iran. *Journal of Entomological Society of Iran*, 28, 99–100.
- Fathpour, H., Emtiazi, G. & Ghasemi, E. (2003) Cockroaches as reservoirs and vectors of drug resistant *Salmonella* spp. *Iranian Biomedical Journal*, 7 (1), 35–38.
- Fuller, C. (1922) The termites of South Africa. *South African Journal of Natural History*, 3 (2), 70–131.
- Gao, H.R., Wang, C., Chen, W.X., Li, J.J., Zhang, H.R. & Li, Y.J. (2025) Across desert and island: Phylogeny of two thin stick insects from China with taxonomic insights into the clade “Gratidiini II” (Phasmatoidea, Clitumninae). *ZooKeys*, 1260, 149. <https://doi.org/10.3897/zookeys.1260.143038>
- Ghahari, H. & El-Den Nasser, M.G. (2014) A contribution to the knowledge of the Mantodea (Insecta) fauna of Iran. *Linzer Biologische Beiträge*, 46 (1), 665–673.
- Ghavami, M.B., Mohammadi, J. & Ladonni, H. (2005) Susceptibility of *Blattella germanica* (Blattaria: Blattidae) to permethrin in Shahid Beheshti hospital, Zanjan. *Journal of Zanjan University of Medical Sciences and Health Services*, 13, 8–16. [In Persian]
- Ghayourfar, R. (1994) Termite fauna of Iran and their economical importance. *Zeitoon*, 123, 150–153. [in Persian]
- Ghayourfar, R. (1996) Geographical distribution, morphological and micrometrical comparison of species *Anacanthotermes vagans* Hagen, *A. turkestanicus* Jacobson and *A. ahngerianus* Jacobson in Iran. *Applied Entomology and Phytopathology*, 63 (1–2), 12–20. [in Persian]
- Ghayourfar, R. (1998) Two new termite species from Iran, *Anacanthotermes esmailii* sp. nov. and *Microcerotermes chhotanii* sp. nov. (Isoptera: Hodotermitidae and Termitidae). *Journal of Entomological Society of Iran*, 16–17, 11–21, 27–29.
- Ghayourfar, R. (1999) Three new termite species from Iran: *Microcerotermes varaminicus* sp. n., *Amitermes iranicus* sp. n., and *Amitermes kharrazii* sp. n. (Isoptera, Termitidae, Termitinae). *Journal of Entomological Society of Iran*, 18 (1–2), 16–31.
- Ghayourfar, R. (2001) Report of *Amitermes belli* (Desneux) (Isop.: Termitidae) from Iran. *Journal of Entomological Society of Iran*, 20 (2), 97–98.
- Ghayourfar, R. (2005) Three new species of termite from Iran (Isoptera, Termitidae). *Zoology in the Middle East*, 34 (1), 61–66. <https://doi.org/10.1080/09397140.2005.10638083>
- Ghayourfar, R. (2006) Zoogeographical study of *Reticulitermes* sp. near *lucifugus* (Isoptera: Rhinotermitidae) in Iran. *Iranian Journal of Forest and Range Protection Research*, 4 (2), 61–70.
- Ghayourfar, R. & Akhtar, M.S. (2005) Report of a new termite species from Iran, *Microcerotermes sistaniensis* sp. n. (Isoptera: Termitidae). *Journal of Entomological Society of Iran*, 24 (2), 155–163.
- Ghayourfar, R. & Esmaili, M. (1993) Introduction of *Microcerotermes buettikeri* Chhotani & Bose and its morphological comparison with *Microcerotermes diversus* Silvestri and *Microcerotermes gabrielis* Weidner. In: Ahoonmanesh, A., Kharazi, A., Rahimian, H. & Baiat, H. (eds) *Proceedings of the 11th Iranian Plant Protection Congress*. 28 August–2 September, 1993. Rasht, Iran. Ministry of Agriculture, Tehran, Vol I, p. 259.
- Grosman, A. (2022) *Thermal Sensitivity and Plasticity in Metabolic Rate and Cuticular Water Loss Across New Zealand Stick Insects (Order: Phasmatoidea)*. Doctoral dissertation, The University of Auckland. 215 pp.
- Habibpour, B., Ekhtelat, M., Khocheili, F. & Mossadegh, M.S. (2010) Foraging population and territory estimates for *Microcerotermes diversus* (Isoptera: Termitidae) through mark-release-recapture in Ahwaz (Khouzestan, Iran). *Journal of Economic Entomology*, 103 (6), 2112–2117. <https://doi.org/10.1603/EC09338>
- Hagen, H.A. (1858) Monographie der Termiten. *Linnaea Entomologica*, 12, I–III + 4–342 + 459.
- Han, W., Qiu, L., Zhang, J., Wang, Z. & Che, Y. (2024) Phylogenetic reconstruction of Corydioidea (Dictyoptera: Blattodea) provides new insights on the placement of Latindiinae and supports the proposal of the new subfamily Ctenoneurinae. *Systematic Entomology*, 49 (1), 156–172. <https://doi.org/10.1111/syen.12610>
- Hanafi-Bojd, A.A. & Sadaghiani, S. (2009) *Cockroaches of Iran*. Tehran University of Medical Sciences Press, Tehran. 148 pp. [In Persian]
- Harris, W.V. (1957) Isoptera. *British Museum (Natural History) report of expedition to South-West Arabia*, 1 (28), 421–433.
- Harris, W.V. (1967) Termites of the genus *Anacanthotermes* in North Africa and the Near East. *Proceedings of the Royal Entomological Society of London (B)*, 36 (5–6), 79–88. <https://doi.org/10.1111/j.1365-3113.1967.tb00540.x>
- Hashemi-Aghdam, S.S. & Oshaghi, M.A. (2015) A checklist of Iranian cockroaches (Blattodea) with description of *Polyphaga* sp. as a new species in Iran. *Journal of Arthropod-Borne Diseases*, 9 (2), 161–171.
- Hellemans, S., Rocha, M.M., Wang, M., Romero Arias, J., Aanen, D.K., Bagnères, A.-G., Buček, A., Carrijo, T.F., Chouvenec, T., Cuzzo, C., Constantini, J.P., Constantino, R., Dedeine, F., Deligne, J., Eggleton, P., Evans, T.A., Hanus, R., Harrison, M.C., Harry, M., Josens, G., Jouault, C., Kalleshwaraswamy, C.M., Kaymak, E., Korb, J., Lee, C.-Y., Legendre, F., Li, H.-F., Lo, N., Lu, T., Matsuura, K., Maekawa, K., McMahon, D.P., Mizumoto, N., Oliveira, D.E., Poulsen, M., Sillam-Dussès,

- D., Su, N.-Y., Tokuda, G., Vargo, E.L., Ware, J. L., Šobotník, J., Scheffrahn, R.H., Canello, E., Roisin, Y., Engel, M.S. & Bourguignon, T. (2024) Genomic data provide insights into the classification of extant termites. *Nature Communications*, 15 (1), 6724. <https://doi.org/10.1038/s41467-024-51028-y>
- Holmgren, N. (1912) Termites from British India (Bombay) collected by Dr. J. Assmuth, S.J. *Journal of the Bombay Natural History Society*, 21 (3), 774–793.
- Holmgren, N. (1913) Termites from British India (near Bombay, in Gujerat and Bangalore) collected by Dr. J. Assmuth, S.J. Part II. *Journal of the Bombay Natural History Society*, 22 (1), 101–117.
- Holt, B.G., Lessard, J.P., Borregaard, M.K., Fritz, S.A., Araújo, M.B., Dimitrov, D., Fabre, P.H., Graham, C.H., Graves, G.R., Jönsson, K.A., Nogués-Bravo, D., Wang, Z., Whittaker, R.J., Fjeldså, J. & Rahbek, C. (2013). An update of Wallace's zoogeographic regions of the world. *Science*, 339 (6115), 74–78. <https://doi.org/10.1126/science.1228282>
- Hopkins, H. & Beccaloni, G. (2023) Cockroach species file online. Available from: <http://cockroach.speciesfile.org/> [Accessed November 27, 2025]
- Hurd, L.E. (1999) Ecology of praying mantids. In: Prete, F.R., Wells, H., Wells, P.H. & Hurd, L.E. (eds) *The Praying Mantids*. Johns Hopkins University Press, Charles Village, Baltimore, MA, pp. 43–60
- Inward, D., Beccaloni, G. & Eggleton, P. (2007) Death of an order: A comprehensive molecular phylogenetic study confirms that termites are eusocial cockroaches [Comparative Study Research Support, Non-U.S. Gov't]. *Biology Letters*, 3 (3), 331–335. <https://doi.org/10.1098/rsbl.2007.0102>
- Jacobson, G. (1905) Zur Kenntnis der Termiten Russlands. *Annuaire du Musée Zoologique de l'Académie Impériale des Sciences de St.-Petersbourg*, 9, 57–107.
- Jamali, M. & Mofidi-Neyestanak, M. (2013) An introduction to the fauna of mantids (Insecta: Mantodea) of northern slopes of Elburz Mountains within Mazandaran Province, Iran. *Journal of Field Crop Entomology*, 3, 67–76.
- Jiang, S., Marco, H.G., Scheich, N., He, S., Wang, Z., Gäde, G. & McMahon, D.P. (2023) Comparative analysis of adipokinetic hormones and their receptors in Blattodea reveals novel patterns of gene evolution. *Insect Molecular Biology*, 32 (6), 615–633. <https://doi.org/10.1111/imb.12861>
- Jowkar, H., Ostrowski, S., Tahbaz, M. & Zahler, P. (2016) The conservation of biodiversity in Iran: Threats, challenges and hopes. *Iranian Studies*, 49 (6), 1065–1077. <https://doi.org/10.1080/00210862.2016.1241602>
- Kaltenbach, A.P. (1963) Kritische Untersuchungen zur Systematik, Biologie und Verbreitung der europäischen Fangheuschrecken (Dictyoptera Mantodea). *Zoologische Jahrbücher. Abteilung für Systematik, Geographie und Biologie der Tiere*, 90, 521–598.
- Kaltenbach, A.P. (1982) Insects of Saudi Arabia. Mantodea. *Fauna of Saudi Arabia*, 4, 29–72.
- Kamyabi, F., Vatandoost, H., Aboulhasani, M., Aghasi, M., Telmadarraiy, Z. & Abaei, M.R. (2006) Sensitivity level of German cockroach (*Blattella germanica*) to three pirimphos methyl, lambda-cyhalothrin and propoxur insecticides in three Kerman hospitals. *Journal of Mazandaran University of Medical Sciences*, 16 (52), 98–108. [In Persian]
- Krishna, K., Grimaldi, D.A., Krishna, V. & Engel, M.S. (2013) Treatise on the Isoptera of the world. *Bulletin of the American Museum of Natural History*, 377, 1–2704. <https://doi.org/10.1206/377.1>
- Kolnegari, M. (2023) Mantodea of Iran. *Journal of Orthoptera Research*, 32 (2), 177–188. <https://doi.org/10.3897/jor.32.96916>
- Kolnegari, M. & Vafaei-Shoushtari, R. (2018) The first Persian boxer mantid, a new species of *Holaptilon* Beier, 1964 from Haftad-gholeh Protected Area, Iran (Mantodea, Mantidae). *Journal of Entomological Research*, 10 (3), 9–17.
- Kolnegari, M., Schwarz, C.J., Ghrejan, T. & Borer, M. (2025) *Sinaielli azadi* sp. nov. (Mantodea: Toxoderidae); a new praying mantis species from Western Asia (Iran and Armenia). *Zootaxa*, 5594 (1), 98–110. <https://doi.org/10.11646/zootaxa.5594.1.4>
- Ladonni, H. (1997) Susceptibility of different field strains of *Blattella Germanica* to four pyrethroides (Orthoptera: Blattellidae). *Iranian Journal of Public Health*, 26 (3–4), 35–40.
- La Greca, M. & Lombardo, F. (1982) Le specie Mediterranee e dell'Asia occidentale del gen. *Rivetina*. *Animalia*, 9, 345–393.
- La Greca, M. & Lombardo, F. (1987) Mantodei del Sistan, con descrizione di un nuovo genere di Oxyothespinae. *Animalia*, 14, 237–246.
- La Sorte, F.A. & Jetz, W. (2010) Projected range contractions of montane biodiversity under global warming. *Proceedings of the Royal Society B: Biological Sciences*, 277 (1699), 3401–3410. <https://doi.org/10.1098/rspb.2010.0612>
- Latifian, M., Rad, B. & Habibpour, B. (2018) Termites of Iranian date palm orchards and their spatial and temporal distribution. *Sociobiology*, 65 (1), 24–30. <https://doi.org/10.13102/sociobiology.v65i1.1784>
- Lee, M. (1993) A survey into the distribution of the stick insects of Britain. *Phasmid Studies*, 4 (1), 15.
- Li, Y., Luo, X.X., Zhang, J.W., Wang, Z.Q. & Che, Y.L. (2022) A new species of *Bundoksia* Lucañas, 2021 with comments on its subfamilial placement, based on morphological and molecular data. *ZooKeys*, 1085, 145–163. <https://doi.org/10.3897/zookeys.1085.72927>
- Limoe, M., Enayati, A.A., Ladonni, H., Vatandoost, H., Baseri, H. & Oshaghi, M.A. (2006) Various mechanisms responsible for permethrin metabolic resistance in seven field-collected strains of the German cockroach from Iran, *Blattella germanica* (L.) (Dictyoptera: Blattellidae). *Pesticide Biochemistry and Physiology*, 87, 138–146. <https://doi.org/10.1016/j.pestbp.2006.07.003>
- Limoe, M., Enayati, A.A., Khassi, K., Salimi, M. & Ladonni, H. (2011) Insecticide resistance and synergism of three field collected strains of the German cockroach *Blattella germanica* (L.) (Dictyoptera: Blattellidae) from hospitals in Kermanshah, Iran. *Tropical Biomedicine*, 28 (1), 111–118.

- Liu, J.L., Zhang, J.W., Han, W., Wang, Y.S., He, S.L. & Wang, Z.Q. (2023) Advances in the understanding of Blattodea evolution: insights from phylotranscriptomics and spermathecae. *Molecular Phylogenetics and Evolution*, 182, 107753. <https://doi.org/10.1016/j.ympev.2023.107753>
- Lombardo, F. (1991) Remarks on the genus *Severinia* Finot 1902 and a description of two new genera of Oxythospinae (Insecta Mantodea). *Tropical Zoology*, 4 (2), 203–207. <https://doi.org/10.1080/03946975.1991.10539490>
- Lo, N., Tokuda, G., Watanabe, H., Rose, H., Slaytor, M., Maekawa, K., Bandi, C. & Noda, H. (2000) Evidence from multiple gene sequences indicates that termites evolved from wood-feeding cockroaches. *Current Biology*, 10, 801–804. [https://doi.org/10.1016/S0960-9822\(00\)00561-3](https://doi.org/10.1016/S0960-9822(00)00561-3)
- Luchetti, A., Marini, M. & Mantovani, B. (2007) Filling the European gap: biosystematics of the eusocial system *Reticulitermes* (Isoptera, Rhinotermitidae) in the Balkanic Peninsula and Aegean area. *Molecular Phylogenetics and Evolution*, 43, 377–383. <https://doi.org/10.1016/j.ympev.2007.07.015>
- Luo, X.X., Li, Q.Q., Zamani, A., Che, Y.L. & Wang, Z.Q. (2023) Redescription of *Periplaneta arabica* (Bey-Bienko, 1938) (Blattodea, Blattellidae), with a comparative analysis of three species of *Periplaneta* Burmeister, 1838 (sensu stricto). *ZooKeys*, 1146, 165. <https://doi.org/10.3897/zookeys.1146.90817>
- Luo, X., Deng, W., Han, W., Lo, N., Cai, J.-Z., Che, Y.-L. & Wang, Z.Q. (2025) Revision of the cockroach subfamily Blattellinae based on morphological and molecular analyses. *Systematic Entomology*, 50 (4), 836–854. <https://doi.org/10.1111/syen.12680>
- Ma, J., Liu, J., Shen, Y., Fan, Z., Yue, B. & Zhang, X. (2019) Population genetic structure and intraspecific genetic distance of *Periplaneta americana* (Blattodea: Blattellidae) based on mitochondrial and nuclear DNA markers. *Ecology and Evolution*, 9 (22), 12928–12939. <https://doi.org/10.1002/ece3.5777>
- Mahjoub, M., Nejati, J. & Keyhani, A. (2010) Evaluation of bacterial infection of external surface and digestive system of cockroach species. *Hormozgan Medical Journal*, 14 (1), 80–86.
- Mahmoudi, P., Kalim, D. & Amirmoradi, M. R. (2011) Investigation of Iran vulnerability trend to desertification with approach of climate change. In *Proceedings of the Second International Conference on Environmental Science and Development IPCBEE*, 4, 63–67.
- Malem, J., Robillard, T., Cluzel, D., Bellier, L., Nattier, R., Grandcolas, P. & Legendre, F. (2023) Origins of old lineages in New Caledonia: A geologically informed test of the island-hopping hypothesis. *Journal of Biogeography*, 50 (9), 1587–1601. <https://doi.org/10.1111/jbi.14673>
- Marques, N., Soares, C.D.M., Casali, D.M., Guimarães, E.C., Fava, F. G., Abreu, J.M.D.S., Moras, L.M., da Silva, L.G., Matias, R., de Assis, R.L., Fraga, R., Almeida, S.M., Lopes, V.G., Oliveira, V., Missagia, R., Carvalho, E.C., Carneiro, N.J., Alves, R., Souza-Filho, P., Oliveira, G., Miranda, M. & Tavares, V.D.C. (2024) Retrieving biodiversity data from multiple sources: Making secondary data standardised and accessible. *Biodiversity Data Journal*, 12, e133775. <https://doi.org/10.3897/BDJ.12.e133775>
- Marshall, J.A. (1983) The orthopteroid insects described by Linnaeus, with notes on the Linnaean collection. *Journal of the Linnean Society of London, Zoology*, 78, 375–396. <https://doi.org/10.1111/j.1096-3642.1975.tb02266.x>
- Mehrabian, A. & Rad, M.A. (2021) Distribution patterns and priorities for conservation of Iranian Endemic Monocots: determining the Areas of Endemism (AOEs). *Journal of Wildlife and Biodiversity*, 5 (2), 69–87.
- Mirzaee, Z. (2024) *Ecology and Phylogeny of Mantodea of Iran and Adjacent Areas*. Doctoral dissertation, University of Potsdam. 187 pp. <https://doi.org/10.25932/publishup-65273>
- Mirzaee, Z. & Pashaei Rad, S. (2017) Seven new records of Mantids (Insecta: Mantodea) for Alborz Mountains, (Tehran Province) Iran. *Iranian Journal of Animal Biosystematics*, 13 (2), 221–228. <https://doi.org/10.22067/ijab.v13i2.61900>
- Mirzaee, Z. & Sadeghi, S. (2019) On a summer collection of mantids (Insecta: Mantodea) from Lorestan Province with nine new records. *Iranian Journal of Animal Biosystematics*, 15 (2), 175–187. <https://doi.org/10.22067/ijab.v15i2.79314>
- Mirzaee, Z. & Sadeghi, S. (2021) A survey on mantids fauna (Insecta: Mantodea) from Fars Province, Iran. *First Fars Biodiversity Conference*. March 2–3, 2021. Shiraz, Iran. Shiraz University. p. 25.
- Mirzaee, Z., Sadeghi, S., Ballarin, F., Schmitt, T., Simões, M. & Wiemers, M. (2023) Life history and biogeography of the enigmatic mantid *Nilomantis floweri* (Mantodea, Nanomantidae). *ZooKeys*, 1173, 275–287. <https://doi.org/10.3897/zookeys.1173.107204>
- Mirzaee, Z., Simões, M.V., Battiston, R., Sadeghi, S., Wiemers, M. & Schmitt, T. (2024a) Biology, ecology, and biogeography of eremic praying mantis *Blepharopsis mendica* (Insecta: Mantodea). *PeerJ*, 12, e16814. <https://doi.org/10.7717/peerj.16814>
- Mirzaee, Z., Battiston, R., Ballarin, F., Sadeghi, S., Simões, M., Wiemers, M. & Schmitt, T. (2024b) The six dwarfs of the Middle East: Revision of the enigmatic praying mantis genus *Holaptilon* (Mantodea: Gonypetidae: Gonypetinae) with the description of four new species under integrative taxonomy. *Arthropod Systematics & Phylogeny*, 82 (1), 89–117. <https://doi.org/10.3897/asp.82.e112834>
- Mirzaee, Z., Wiemers, M., Schmitt, T., Govorov, V. & Shcherbakov, E. (2026) Review of the genus *Hierodula* Burmeister (Mantodea: Mantidae) in Iran. *Frontiers in Insect Science*, 6, 1757219. <https://doi.org/10.3389/finsc.2026.1757219>
- Mirzayans, H. (1986) Fauna of Iranian cockroaches (Orthopteroidea: Blattaria). *Journal of the Entomological Society of Iran*, 4 (Suppl.), 1–34. [In Persian]
- Mistshenko, L. (1937) A study of the genus *Gratidia* Stål (Phasmodea, Clitumnidae). *Proceedings of the Royal Entomological Society of London. Series B* 6, 119–120. <https://doi.org/10.1111/j.1365-3113.1937.tb00452.x>

- Mohammadi, J. (1998) Determining the active genera of cockroaches living in the hospitals and residential houses of Zanjan; their seasonal activities, and bacterial pollutions. *Proceedings of the 1st Congress on Iran Medical Entomology*. Pasteur Institute, Tehran, Iran. 139.
- Moradi, M., Joger, U., Shafiei Bafti, S., Sharifi, A. & Sehhati Sabet, M.E. (2024) Biogeography of the Iranian snakes. *Plos One*, 19 (10), e0313751. <https://doi.org/10.1371/journal.pone.0313751>
- Moradzadeh, S., Pashaei Rad, S. & Shahbazi, F. (2021) Introducing four new records of Mantis for the fauna of Lahijan city Guilan Province. *Experimental Animal Biology*, 9 (3), 55–61.
- Morshedi Aghbolagh, A., Mofidi-Neyestanak, M. & Taghizadeh, M. (2012) Fauna of Mantodea (Insecta) of Dasht-e Moghan in Ardabil Province, Iran. *Field Crop Entomology*, 1, 85–94.
- Mousavi, S.B., Vatandoust, H. & Rafinezhad, J. (2004) Evaluation of the resistance level of different strains of German cockroach, *Blattella germanica* (L.), to delta-methrin, propoxur and pirimiphos-methyl insecticides in Tehran, 2000. *Pathobiology Research (Modares journal of medical sciences)*, 7 (1), 97–106. [In Persian]
- Motavali-Haghi, F., Gholami, S.H. & Sedaghat, M.M. (1997) Survey of susceptibility of cockroaches to insecticides in Sari hospitals in 1996. *Journal of Mazandaran University of Medical Sciences*, 7 (16), 1–6.
- Moulin, N. (2020) When citizen science highlights alien invasive species in France: The case of Indochina mantis, *Hierodula patellifera* (Insecta, Mantodea, Mantidae). *Biodiversity Data Journal*, 8, e46989. <https://doi.org/10.3897/BDJ.8.e46989>
- Naderinasab, M. & Moravvej, G.H. (2012) Bacterial and fungal contamination of cockroaches at two seasons in public habitants of Mashhad, Iran. *Proceedings of the 13th Iranian and the Second International Congress of Microbiology*, July 14–16, 2012. Ardabil University of Medical Sciences, Ardabil, Iran, p. 81.
- Nasirian, H. (2007) Duration of fipronil and imidacloprid gel baits toxicity against *Blattella germanica* strains of Iran. *Iranian Journal of Arthropod-Borne Diseases*, 1 (2), 40–47.
- Nejati, J., Keyhani, A., Moosa-Kazemi, S.S., Mohammadi, M., Mahjoob, M. & Boostanbakhsh, A. (2011) Cockroaches' bacterial infections in wards of hospitals, Hamedan City, west of Iran. *Asian Pacific Journal of Tropical Disease*, 2 (5), 381–384. [https://doi.org/10.1016/S2222-1808\(12\)60083-8](https://doi.org/10.1016/S2222-1808(12)60083-8)
- Noori, S., Zahiri, R., Yusefi, G. H., Rajabzadeh, M., Hawlitschek, O., Rakhshani, E., Husemann, M. & Rajaei, H. (2024) Patterns of zoological diversity in Iran—A review. *Diversity*, 16 (10), 621. <https://doi.org/10.3390/d16100621>
- Olson, D.M., Dinerstein, E., Wikramanayake, E. D., Burgess, N. D., Powell, G.V.N., Underwood, E.C., D'Amico, J.A., Itoua, I., Strand, H.E., Morrison, J.C., Loucks, C. J., Allnutt, T.F., Ricketts, T.H., Kura, Y., Lamoreux, J.F., Wettengel, W.W., Hedao, P. & Kassem, K.R. (2001) Terrestrial ecoregions of the world: A new map of life on Earth. *Bioscience*, 51 (11), 933–938. [https://doi.org/10.1641/0006-3568\(2001\)051\[0933:TEOTWA\]2.0.CO;2](https://doi.org/10.1641/0006-3568(2001)051[0933:TEOTWA]2.0.CO;2)
- Otte, D. & Brock, P.D. (2005) *Phasmida Species File. Catalog of stick and leaf insects of the world*. The Insect Diversity Association at the Academy of Natural Sciences, <https://phasimida.speciesfile.org/> [Accessed September 16, 2025]
- Otte, D., Spearman, L. & Stiewe, M.B.D. (2023) *Mantodea Species File Online*. Retrieved from <http://Mantodea.SpeciesFile.org/> [Accessed September 28, 2025]
- Pérez-Gómez, Á., León-Osper, M., Pareja, D. & Robla, J. (2023) Flower visits of cockroaches (Insecta: Blattodea) in the Iberian Peninsula: Are they neglected pollinators? *Journal of Applied Entomology*, 147 (8), 565–576. <https://doi.org/10.1111/jen.13157>
- Poinar, G. (2022) *Supella dominicana*, a new species of cockroach (Blattida: Ectobiidae) with developed spermatids in Dominican amber. *Biologia*, 78 (6), 1731–1738. <https://doi.org/10.1007/s11756-022-01271-9>
- Pringle, R.M., Kartzinel, T.R., Palmer, T.M., Thurman, T.J., Fox-Dobbs, K., Xu, C.C.Y., Hutchinson, M.C., Coverdale, T.C., Daskin, J.H., Evangelista, D.A., Gotanda, K.M., Man in 't Veld, N.A., Wegener, J.E., Kolbe, J.J., Schoener, T.W., Spiller, D. A., Losos, J.B. & Barrett, R.D.H. (2019) Predator-induced collapse of niche structure and species coexistence. *Nature*, 570 (7759), 58–64. <https://doi.org/10.1038/s41586-019-1264-6>
- Rabieh, M.M., Rouhani, A. & Battiston, R. (2016) Preliminary study of Mantodea fauna of South Khorasan Province. In: Talei-Hansanloo, R. (ed.) *Proceedings of the 22nd Iranian Plant Protection Congress*. August 27–30, 2016. University of Tehran, College of Agriculture & Natural Resources, Karaj, Iran. P. 459.
- Ramme, W. (1951) Zur Systematik, Faunistik und Biologie der Orthopteren von Süd-Ost-Europa und Vorder-Asien. *Mitteilungen aus dem Zoologischen Museum in Berlin*, 27, 1–431. <https://doi.org/10.1002/mmnz.4830270108>
- Ravan, S. (2010) Evaluation of the toxicity and repellency of Dimilin, Sevin and Dimecron against *Postelectrotermes pasniensis*, 1276–1279.
- Ravan, S. & Akhtar, M.S. (1993) A new termite from Iran. *Proceedings of the Pakistan Congress of Zoology*, 13, 555–560.
- Ravan, S. & Akhtar, M.S. (1998) Feeding preferences and diversity of termites of Iran. *Punjab University Journal of Zoology*, 13, 9–26
- Ravan, S. & Akhtar, M.S. (1999) New termites from Iran. *Punjab University Journal of Zoology*, 14, 65–85.
- Ravan, S., Khan, I.A., Manzoor, F. & Khan, Z. (2015) Feeding habitats and wood preferences of termites in Iran. *Journal of Entomology and Zoology Studies*, 3, 20–23.
- Rouhani, A., Pashaei Rad, S. & Maleki, L. (2015) The mantis fauna (Insecta: Mantodea) of Southern part of Kordestan province, Iran. *1st Iranian International Congress of Entomology*. August 29–31, 2015. Iranian Research Institute of Plant Protection, Tehran, Iran. p. 15
- Roonwal, M.L. & Bose, G. (1960) A new termite, *Psamotermes rajasthanicus* sp. nov. from Rajasthan, India. *Science and Culture*, 26 (1), 38–39.

- Ren, Y.M., Han, W., Che, Y.L. & Wang, J.R. (2025) Four new species of the genus *Pseudoeupolyphaga* Qiu & Che, 2024 (Blattodea, Corydioidea, Corydiinae) from Yunnan, China. *Zookeys*, 261, 337–358. <https://doi.org/10.3897/zookeys.1261.168015>
- Sadeghi, N. & Sadeghi, S. (2015) *A faunal study of suborder Mantodea in Fars Province (Iran)*. MSc thesis. University of Shiraz, 101 pp.
- Sakenin, H., Samin, N., Shakouri, M. J., Mohebbi, H. R., Ezzatpanah, S. & Moemen Beitollahi, S. (2011) A faunistic survey of the insect predators in some regions of Iran. *Calodema*, 142, 1–10.
- Salehzadeh, A. (1995) The study of Blattaria and their susceptibility to Ficam and Lindane in Hamadan. *Avicenna Journal of Clinical Medicine*, 2 (2), 32–36.
- Saussure, H. (1870) Additions au Système des Mantides. *Mitteilungen der Schweizerischen Entomologischen Gesellschaft*, 3 (7), 221–244.
- Schwarz, C.J. & Ehrmann, R. (2018) Invasive Mantodea species in Europe. *Articulata*, 33, 73–90.
- Schwarz, C.J. & Roy, R. (2019) The systematics of Mantodea revisited: An updated classification incorporating multiple data sources (Insecta: Dictyoptera). *Annales de la Société Entomologique de France (N.S.)*, 55 (2), 101–196. <https://doi.org/10.1080/00379271.2018.1556567>
- Sevgili, H. & Yılmaz, K. (2022) Contributions of citizen scientists to monitoring alien species: The case study on Giant Asian Mantids, *Hierodula tenuidentata* and *H. patellifera* (Mantodea: Mantidae). *Zoology in the Middle East*, 68 (4), 350–358. <https://doi.org/10.1080/09397140.2022.2145802>
- Shahi, M., Hanafi, B.A. & Vatandoust, H. (2008) Evaluation of five local formulated insecticides against German cockroach (*Blattella germanica* L.) in Southern Iran. *Journal of Arthropod-Borne Diseases*, 2 (1), 21–27.
- Shahraki, G.H., Noor, H.M., Rafinejad, J., Shahar, M.K. & Bin-Ibrahim, Y. (2010) Biorational control programme for the German cockroach (Blattaria: Blattellidae) in selected urban communities. *Tropical Biomedicine*, 27 (2), 226–235.
- Shahraki, G.H., Noor, H.M., Rafinejad, J., Shahard, M.K. & Bin-Ibrahim, Y. (2011) Field efficacy of siege gel bait in an IPM program on life stages of German cockroach (Blattaria, Blattellidae) in a residential building. *Asian Biomed*, 5 (5), 619–624.
- Shcherbakov, E. & Govorov, V. (2020) *Statilia maculata* (Thunberg, 1784)—the first invasive praying mantis (Mantodea, Mantidae) in the fauna of Russia. *Annales de la Société Entomologique de France (N.S.)*, 56 (3), 189–202. <https://doi.org/10.1080/00379271.2020.1785941>
- Sugimoto, A., Bignell, D.E. & MacDonald, J.A. (2000) Global impact of termites on the carbon cycle and atmospheric trace gases. In Abe, T., Bignell, D.E. & Higashi, M. (eds) *Termites: Evolution, Sociality, Symbioses, Ecology*. Springer, Dordrecht. pp. 409–435. https://doi.org/10.1007/978-94-017-3223-9_19
- Tarli, V.D., Pequeno, P.A.C.L., Franklin, E., Oliveira, J. & Guilherme, D.R. (2014) Multiple Environmental Controls on Cockroach Assemblage Structure in a Tropical Rain Forest. *Biotropica*, 46 (5), 598–607.
- Uvarov, B.P. (1922) Records and descriptions of Orthoptera from S.W. Asia. *Journal of the Bombay Natural History Society*, 28, 719–725.
- Uvarov, B.P. (1938) Orthoptera from Iraq and Iran. *Zoological series of Field Museum of Natural History*, 20 (32), 439–451.
- Uvarov, B.P. (1939) Twenty-four new generic names in Orthoptera. *Annals and Magazine of Natural History*, Series 11, 3 (16), 457–459. <https://doi.org/10.1080/03745481.1939.9723626>
- Vahabi, A., Rafinezhad, J., Mohammadi, P. & Biglarian, F. (2007) Regional evaluation of bacterial contamination in hospital environment cockroaches. *Iranian Journal of Environmental Health Science & Engineering*, 4 (1), 57–60.
- Vaissi, S. & Mohammadi, A. (2024) Climate-driven distribution shifts of Iranian amphibians and identification of refugia and hotspots for effective conservation. *Scientific Reports*, 14, 31610. <https://doi.org/10.1038/s41598-024-79293-3>
- Valizadeh, R. (2010) Iranian sheep and goat industry at a glance. In Karim, S.A. & Joshi, A. (eds) *Climate Change and Stress Management: Sheep and Goat Production*. Satish Serial Publishing House, Mashhad, pp. 547–551.
- Vazirianzadeh, B., Mehdinejad, M. & Dehghani, R. (2011) Identification of bacteria which possible transmitted by *Polyphaga aegyptica* (Blattodea: Blattidae) in the region of Ahvaz, SW Iran. *Jundishapur Journal of Microbiology*, 2 (1), 36–40.
- Vršanský, P., Cifuentes-Ruiz, P., Vidlička, L., Čiampor, F. & Vega F.J. (2011) Afro-Asian cockroach from Chiapas amber and the lost Tertiary American entomofauna. *Geologica Carpathica*, 62 (5), 463–475. <https://doi.org/10.2478/v10096-011-0033-8>
- Wang, L., Xiong, Q., Saelim, N., Wang, L., Nong, W., Wan, A.T.Y., Shi, M., Liu, X., Cao, Q., Hui, J.H.L., Sookrung, N., Leung, T.F., Tungtrongchitr, A. & Tsui, S.K.W. (2023) Genome assembly and annotation of *Periplaneta americana* reveal a comprehensive cockroach allergen profile. *Allergy*, 78 (4), 1088–1103. <https://doi.org/10.1111/all.15531>
- Weidner, H. (1960) Die termiten von Afghanistan, Iran und Irak (Contribution a l'etude de la fauna d'Afghanistan 29). *Abhandlungen und Verhandlungen des Naturwissenschaftlichen Vereins in Hamburg [N.F.]*, 4, 43–70.
- Werner, F. (1905) Ergebnisse einer zoologischen Forschungsreise nach Ägypten und dem ägyptischen Sudan. I.- Die Orthopterenfauna Ägyptens mit besonderer Berücksichtigung der *Eremiaphila*. *Sitzungsberichte der Mathematisch-Naturwissenschaftlichen Klasse der Österreichischen Akademie der Wissenschaften*, 114 (1), 357–436.
- Werner, F. (1930) Über asiatische Mantidae aus dem naturhistorischen Reichsmuseum in Stockholm. *Arkiv för Zoologi*, 21A (1), 1–10. <https://doi.org/10.5962/bhl.part.1498>
- Zarchi, A.A.K. & Vatani, H. (2009) A survey on species and prevalence rate of bacterial agents isolated from cockroaches in three hospitals. *Vector-Borne and Zoonotic Diseases*, 9 (2), 197–200. <https://doi.org/10.1089/vbz.2007.0230>
- Zohary, M. (1973) *Geobotanical Foundations of the Middle East*. Volumes 1–2. Fischer, Stuttgart: Gustav Fischer Verlag; Amsterdam, Swets & Zeitlinger. 739 pp.

تنوع بندپایان در ایران: رده حشرات – گروه پلی نئوترها، بخش اول. Blattodea Brunner von Phasmatodea Jacobson & Bianchi, و Mantodea Brumeister, 1838, Wattenwyl, 1882 1902

زهره میرزایی^{۱*}، تیس اچ. بوشر^۲، دومینیک اوانجلیستا^۳، پل اگلتون^۴، لئونید آنیسووتکین^۵، فردریک لجندر^۶

- ۱ موسسه حشره‌شناسی سنکنبرگ، خیابان ابرسوالدر ۹۰، مونچبرگ، آلمان.
- ۲ بخش مطالعات ریخت‌شناسی عملکردی و بیومکانیک، دانشگاه کیل، باغ گیاه‌شناسی، کیل، آلمان
- ۳ سالن ۴۶۵ موریل، دانشگاه ایلینوی، اوربانا-شمپاین، اوربانا، ایلینوی، ایالات متحده.
- ۴ گروه تنوع زیستی خاک، بخش حشره‌شناسی، موزه تاریخ طبیعی، لندن، انگلستان.
- ۵ آزمایشگاه سیستماتیک حشرات، موسسه جانورشناسی آکادمی علوم روسیه، سن پترزبورگ، روسیه.
- ۶ موسسه مطالعات سیستماتیک، تکامل و تنوع زیستی، موزه ملی تاریخ طبیعی، پاریس، فرانسه.

نویسندهٔ مسئول: زهره میرزایی | zmirzaee1988@gmail.com

ویراستار علمی

علیرضا زمانی

دریافت: ۱۹ مهر ۱۴۰۴

ویرایش: ۲۵ بهمن ۱۴۰۴

پذیرش: ۲۶ بهمن ۱۴۰۴

انتشار: ۰۸ اسفند ۱۴۰۴

چکیده: این پژوهش، بر راسته‌های حشرات Mantodea (آخوندک‌ها)، Blattodea (سوسری‌ها و موربانه‌ها)، و Phasmatodea (حشرات چوبک‌مانند) در ایران ارایه می‌دهد که بر اساس منابع علمی، فهرست‌های ملی، و پایگاه‌های داده‌های طبقه‌بندی جهانی تهیه شده است تا فهرستی به‌روز از گونه‌های موجود تهیه شود. با پیروی از طبقه‌بندی‌های پذیرفته‌شده اخیر، این مرور در مجموع سه راسته، ۲۰ خانواده، ۴۴ جنس، و ۱۰۴ گونه را ثبت می‌کند که شامل ۹ خانواده و ۱۹ جنس و ۳۳ گونه Mantodea، ۱۰ خانواده و ۲۳ جنس و ۶۵ گونه Blattodea (سوسری‌ها و موربانه‌ها)، و یک خانواده با دو جنس و ۵ گونه Phasmatodea است. بیشترین مطالعات در ایران روی راسته Mantodea، و کمترین بررسی روی راسته Phasmatodea صورت گرفته به نحوی که تنها ۵ گونه از آنها تا کنون ثبت شده است. ادغام داده‌های طبقه‌بندی به‌روز برای رفع ناسازگاری‌های موجود در گزارش‌های قدیمی یا تأیید نشده و ایجاد مبنای قابل اعتماد برای تحلیل‌های زیست‌جغرافیایی، برنامه‌ریزی حفاظت، و مدیریت گونه‌های بومی و مهاجم ضروری است. به‌ویژه برای Phasmatodea، مشخص است که طبقه‌بندی این گروه جانوری هم برای آشکارسازی تنوع زیستی و همچنین به منظور مشخص کردن جایگاه این آرایه‌های در سطوح بالاتر تکاملی، نیاز به مطالعه بیشتر دارد. ادامه بررسی‌های میدانی و مطالعات تاکسونومیک تلفیقی برای مستندسازی کامل تنوع این گروه‌های حشره در زیستگاه‌های متنوع ایران، ضروری است.

واژگان کلیدی: تنوع زیستی، چک‌لیست، سوسری‌ها، تورینه‌بالان، چوبک‌مانندها، تاکسونومی، موربانه‌ها