

Updated Checklist of Vespidae (Hymenoptera: Vespoidea) in Iran

Zahra Rahmani¹, Ehsan Rakhshani^{1*} & James Michael Carpenter²

¹ Department of Plant Protection, College of Agriculture, University of Zabol, P.O. Box 98615-538, I.R. Iran.

² Division of Invertebrate Zoology, American Museum of Natural History, Central Park West at 79th Street, New York, NY 10024, USA.

ABSTRACT. 231 species of the family Vespidae (Hymenoptera, Vespoidea) of Iran, in 55 genera belonging to 4 subfamilies Eumeninae (45 genera, 184 species), Masarinae (5 genera, 24 species), Polistinae (2 genera, 17 species) and Vespinae (3 genera, 6 species) are listed. An overall assessment of the distribution pattern of the vespid species in Iran indicates a complex fauna of different biogeographic regions. 111 species are found in both Eastern and Western Palaearctic regions, while 67 species were found only in the Eastern Palaearctic region. Few species (14 species – 6.1%) of various genera are known as elements of central and western Asian area and their area of distribution is not known in Europe (West Palaearctic) and in the Far East. The species that were found both in the Oriental and Afrotropical Regions comprises 11.7 and 15.6% the Iranian vespid fauna, respectively. Many species (48, 20.8%) are exclusively recorded from Iran and as yet there is no record of these species from other countries. The highest percentage of the vespid species are recorded from Sistan-o Baluchestan (42 species, 18.2%), Alborz (42 species, 18.2%), Fars (39 species, 16.9%) and Tehran provinces (38 Species 16.5%), representing the fauna of the Southeastern, North- and South Central of the country.

Received:
01 January, 2020

Accepted:
17 January, 2020

Published:
31 January, 2020

Subject Editor:
Volker Mauss

Key words: Catalogue, Distribution, Hornets, Potter wasps, Paper wasps, Pollen wasps, Yellow jackets

Citation: Rahmani, Z., Rakhshani, E. & Carpenter, J.M. (2020) Updated Checklist of Vespidae (Hymenoptera: Vespoidea) in Iran. *Journal of Insect Biodiversity and Systematics*, 6 (1), 27–86.

Introduction

Vespidae is one of the largest and most diverse families of the order Hymenoptera with more than 5000 species worldwide in six subfamilies: Euparagiinae (single genus, 10 species), Masarinae (14 genera, 374 species), Eumeninae (204 genera, 3844 species), Stenogastrinae (7 genera, 72 species), Polistinae (25 genera, 1024 species), and Vespinae (4 genera, 69 species) (Pickett & Carpenter, 2010; Carpenter, unpubl.). Species of the family Vespidae play important roles in ecosystems, as biological control agents, since they feed their larvae with caterpillars and other insects (Ross & Matthews, 1991). They also serve as pollinators of various plants (Fateryga, 2010). Adult vespids are commonly black or brown

Corresponding author: Ehsan Rakhshani, E-mail: rakhshani@uoz.ac.ir

Copyright © 2020, Rahmani et al. This is an open access article distributed under the terms of the Creative Commons Attribution License (CC BY NC 4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

in color, but often marked with yellow or white patterns. Most species are solitary, but many species are social (Goulet & Huber, 1993). The social species belong to the subfamilies Stenogastrinae, Polistinae and Vespinae, while all members of the Euparagiinae, Masarinae and Eumeninae live solitarily (Ross & Matthews, 1991). In solitary species, the larvae feed normally on larvae of other insects, especially caterpillars, within a cell constructed and provisioned by the adult female, but in the Masarinae the larva is supplied with a mixture of pollen and nectar instead. In social species the larva is progressively fed by the adult females on masticated insects (Ross & Matthews, 1991; Goulet & Huber, 1993).

Scattered data on Vespidae of Iran were partly reviewed in Farahbakhsh (1961) and Esmaili & Rastegar (1974). Subsequent faunistic or taxonomic works (Pashaei Rad et al., 2006, 2010; Abbasi et al., 2008; Ebrahimi & Carpenter, 2008, 2012; Salehi Fard et al., 2013; Khoobdel et al., 2014; Nematy et al., 2016; Hamzavi et al., 2019) contributed to increasing the knowledge on the Vespidae of Iran. The first comprehensive study of the vespid fauna of Iran was done by Blüthgen & Gusenleitner (1970), in which numerous species were recorded from various regions. Many more vespid species of Iran were treated by Gusenleitner (1972, 1973b, 1985b, 1986a, 1986b, 1992, 1995a, 1998c, 2000c, 2001, 2002, 2004, 2006a, 2010a, 2011a, 2012a, 2012b, 2014, 2016, 2018), in his long series of publications. Various papers including data on the recorded Vespidae from Iran were also published, elsewhere (e.g. Morawitz, 1885; Kostylev, 1937, 1940; Giordani Soika, 1961, 1970; Guiglia, 1976, 1977; Archer, 1981; Castro & Dvořák, 2009; Dvořák et al., 2012; Gusenleitner & Ockermüller, 2018; Selis, 2019).

In the last catalogue of Vespidae of Iran (Ebrahimi & Carpenter, 2008), 51 genera and 182 species were recorded with relevant data about distribution in various parts of the country. Since then, many other species have been recorded from Iran or their distribution in other parts of the country is documented. On the other hand, the status of some recorded species was changed from that considered in Ebrahimi & Carpenter (2008). Here we present an updated checklist of Vespidae from Iran with details based on the original references indicating the recent advances of the knowledge on the fauna of Vespidae in various parts of the country.

Material and methods

The known vespid species recorded from Iran were compiled and listed. Data about general distribution of the recorded species were compiled from the selected relevant literature. The provincial distribution of each species was also based on the locality records in the published data. Classification and nomenclature followed Carpenter (1996, 2001), Carpenter & Kojima (1997), Gusenleitner (2013, 2019) and Schmid-Egger et al. (2017). The occurrence of species in Iran as originally indicated the junior synonyms under which the species were previously recorded from Iran are ignored. The valid name of each species is presented according to the last taxonomic modification. Other taxonomic changes and occurrence of the subspecies were also highlighted as separate remarks. Within the subfamilies the species are listed and presented alphabetically.

Results

231 species of the family Vespidae in 55 genera belonging to four subfamilies recorded from Iran are listed.

Family: Vespidae
Subfamily: Eumeninae Leach, 1815

Genus: *Acanthodynerus* Gusevleitner, 1969

Acanthodynerus Gusevleitner, 1969: 13. Type species: *Acanthodynerus giordanii* Gusevleitner, 1969, by original designation.

***Acanthodynerus multimaculatus* Gusevleitner, 2000**

Acanthodynerus multimaculatus Gusevleitner, 2000: 932, ♀. – Turkey.

Distribution in Iran: Isfahan ([Gusevleitner, 2011a](#)).

General distribution: Eastern (Iran) and Western (Israel, Turkey) Palaearctic ([Gusevleitner, 2013](#)).

Genus: *Alastor* Lepeletier, 1841

Alastor Lepeletier, 1841: 668. Type species: *Alastor atropos* Lepeletier, 1841. Designated by Ashmead, 1902.

***Alastor (Alastor) darius* Gusevleitner, 1986**

Alastor (Alastor) darius Gusevleitner, 1986: 30, ♀♂: Holotype ♀. – Iran.

Distribution in Iran: Hormozgan, Fars ([Gusevleitner, 1986b](#)).

General distribution: Eastern Palaearctic (Iran) ([Gusevleitner, 2013](#)).

***Alastor (Alastor) iranus* Blüthgen, 1956**

Alastor (Alastor) iranus Blüthgen, 1956: 125, ♀. – Iran.

Distribution in Iran: Sistan-o Baluchestan ([Blüthgen, 1956](#); [Blüthgen & Gusevleitner, 1970](#)).

General distribution: Eastern Palaearctic (Iran) ([Gusevleitner, 2013](#)).

***Alastor (Alastor) xerxes* Gusevleitner, 1986**

Alastor (Alastor) xerxes Gusevleitner, 1986b: 33, ♀. – Iran.

Distribution in Iran: Hormozgan ([Gusevleitner, 1986b](#)).

General distribution: Eastern Palaearctic (Iran) ([Gusevleitner, 2013](#)).

***Alastor (Alastor) zoroaster* Gusevleitner, 1986**

Alastor (Alastor) zoroaster Gusevleitner, 1986: 35, ♀. – Iran.

Distribution in Iran: Hormozgan ([Gusevleitner, 1986b](#)).

General distribution: Eastern Palaearctic (Iran) ([Gusevleitner, 2013](#)).

***Alastor (Megalastor) esfandiarii* Giordani Soika, 1970**

Alastor (Megalastor) esfandiarii Giordani Soika, 1970: 58, ♂. – Iran.

Distribution in Iran: Sistan-o Baluchestan ([Giordani Soika, 1970](#)).

General distribution: Eastern (Iran - [Giordani Soika, 1970](#)) and Western (Turkey - [Yildirim & Kojima, 1999](#)) Palaearctic.

Genus: *Allodynerus* Blüthgen, 1938

Allodynerus Blüthgen, 1938: 280. Type species: “*Lionotus floricola* Sauss, 1852” [= *Odynerus floricola* de Saussure, 1853].

***Allodynerus delphinalis* (Giraud, 1866)**

Odynerus (Lionotus) delphinalis Giraud, 1866: 464, ♀. – Grenoble [Paris].

Distribution in Iran: East Azerbaijan ([Ebrahimi & Carpenter, 2008](#)).

General distribution: Eastern (China, Japan, Iran, Kazakhstan, Korea, Russian Far East) and Western (Lebanon, North Africa, Russia, Turkey, Europe) Palaearctic ([Fateryga, 2018](#)).

***Allodynerus dignotus* (Morawitz, 1895)**

Odynerus (Lionotus) dignotus Morawitz, 1895: 457, ♀♂. – Transcaucasia: Tschemachlinskaja.

Distribution in Iran: Khorasan-e Razavi ([Salehi Fard et al., 2013](#)), Sistan-o Baluchestan ([Blüthgen & Guseinleitner, 1970](#)).

General distribution: Afrotropical (Oman), Eastern (Iran, Kazakhstan, Pakistan, Turkmenistan) and Western (Iraq, Greece, Israel, Jordan, Syria, Turkey, UAE) Palaearctic ([Rafi et al., 2017](#)).

***Allodynerus rossii* (Lepeletier, 1841)**

Odynerus rossii Lepeletier, 1841: 633, ♀♂. – Environs de Paris. (destroyed).

Distribution in Iran: Alborz ([Giordani Soika, 1970](#)), East Azarbaijan ([Ebrahimi & Carpenter, 2008](#)).

General distribution: Eastern (Iran, Turkmenistan) and Western (Caucasus, Europe, Northwest Africa, Russia, Turkey) Palaearctic ([Fateryga, 2018](#)).

Genus: *Ancistrocerus* Wesmael, 1836

Ancistrocerus Wesmael, 1836: 45. Type species: *Vespa parietum* L., 1758. Designated by Girard, 1879.

***Ancistrocerus auctus* (Fabricius, 1793)**

Vespa aucta Fabricius, 1793: 272, ♀. – Kiliae [København].

Distribution in Iran: East Azarbaijan ([Ebrahimi & Carpenter, 2008](#)), Zanjan ([Abbasi et al., 2008; Pashaei Rad et al., 2010](#)).

General distribution: Eastern (Iran, Kazakhstan) and Western (Europe, Israel, Jordan, Russia, Syria, Turkey) Palaearctic ([Fateryga, 2018](#)).

***Ancistrocerus biphaleratus* (de Saussure, 1852)**

Odynerus (Ancistrocerus) biphaleratus de Saussure, 1852: 134, ♀♂. – Egypt.

Distribution in Iran: Hormozgan ([Guseinleitner, 2010a](#)).

General distribution: Eastern (Iran) and Western (Europe, Northwest Africa, Turkey) Palaearctic ([Guseinleitner, 2013](#)).

Remarks: The subspecies *Ancistrocerus biphaleratus triphaleratus* (de Saussure, 1855) has been recorded from Iran ([Guseinleitner, 2010a](#)).

***Ancistrocerus claripennis* Thomson, 1874**

Ancistrocerus claripennis Thomson, 1874: 76, ♀♂. – Sweden.

Distribution in Iran: Ardabil ([Ebrahimi & Carpenter, 2008](#)), Mazandaran ([Guseinleitner, 2011a](#)).

General distribution: Eastern (China, Tajikistan, Iran) and Western (Caucasus, Europe, Russia, Turkey) Palaearctic ([Antropov & Fateryga, 2017](#)).

Remarks: The subspecies *Ancistrocerus claripennis claripennis* has been recorded from Iran ([Ebrahimi & Carpenter, 2008; Guseinleitner, 2011a](#)).

***Ancistrocerus gazella* (Panzer, 1798)**

Vespa gazella Panzer, 1798: 10, ♂. – Austria (destroyed).

Distribution in Iran: No locality cited ([Gusenleitner, 1995b](#)).

General distribution: Eastern Palaearctic (Afghanistan, Iran, Pakistan), Nearctic (Canada, U.S.A – exotic), Oceanic (New Zealand – exotic), Western Palaearctic (Azerbaijan, Cyprus, Europe, Israel, North Africa, Russia, Turkey) ([Fateryga et al., 2019c](#)).

***Ancistrocerus nigricornis* (Curtis, 1826)**

Odynerus nigricornis Curtis, 1826: 137. – Great Britain.

Distribution in Iran: No locality cited ([Gusenleitner, 1995b](#)).

General distribution: Oriental (India), Eastern (China, Iran, Japan, Kazakhstan, Mongolia, Russian Far East) and Western (Europe, North Africa, Russia, Turkey) Palaearctic ([Antropov & Fateryga, 2017](#)).

***Ancistrocerus parietinus* (Linnaeus, 1761)**

Vespa parietina Linnaeus, 1761: 418. – Sweden.

Distribution in Iran: No locality cited ([Gusenleitner, 1995b](#)).

General distribution: Eastern (Iran, Japan, Kazakhstan, Korea, Mongolia, Russia Far East) and Western (Caucasus, Europe, North Africa, Russia, Turkey) Palaearctic ([Fateryga, 2018](#)).

***Ancistrocerus parietum* (Linnaeus, 1758)**

Vespa parietum Linnaeus, 1758: 572.

Distribution in Iran: Qazvin ([Morice, 1921](#)); No locality cited ([Kurzenko, 1978](#)).

General distribution: Eastern (Iran, Japan, Kazakhstan, Korea, Mongolia) and Western (Armenia, Azerbaijan, Europe, North Africa, Russia, Turkey) Palaearctic ([Fateryga et al., 2019](#)).

Remarks: This species has been recorded from Iran as *Odynerus parietum* (Linnaeus, 1758) ([Morice, 1921](#)).

***Ancistrocerus scoticus* (Curtis, 1826)**

Odynerus scoticus Curtis, 1826: 137. – Scotland.

Distribution in Iran: No locality cited ([Ebrahimi & Carpenter, 2008](#)).

General distribution: Eastern (China, Iran, Kazakhstan, Korea, Mongolia, Russia) and Western (Europe, Georgia, North Africa, Russia, Turkey) Palaearctic ([Antropov & Fateryga, 2017](#)).

***Ancistrocerus tenebrosus* Gusenleitner, 2010**

Ancistrocerus tenebrosus Gusenleitner, 2010: 1349, ♀♂, Holotype ♂. – Iran.

Distribution in Iran: Mazandaran ([Gusenleitner, 2010a](#)).

General distribution: Eastern Palaearctic (Iran) ([Gusenleitner, 2013](#)).

Genus: *Antepipona* de Saussure, 1855

Antepipona Saussure, 1855: 244. Type species: *Odynerus silaos* Saussure, 1853. Designated by ICZN, 1970.

***Antepipona barrei* (Radoszkovski, 1893)**

Odynerus barrei Radoszkovski, 1893: 76, ♀. – Turkmenistan.

Distribution in Iran: Tehran ([Giordani Soika, 1970](#)); No locality cited ([Castro & Dvořák, 2009](#)).

General distribution: Eastern (Afghanistan, Iran, Turkmenistan) and Western (Armenia) Palaearctic ([Giordani Soika, 1970](#); [Gusenleitner, 2013](#)).

Remarks: The subspecies *Antepipona barrei vastifica* (Morawitz, 1895) has been recorded from Iran ([Giordani Soika, 1970](#)).

Antepipona deflenda (S. S. Saunders, 1853)

Ancistrocerus deflendus S. S. Saunders, 1853: 141, ♀♂. – Albania.

Distribution in Iran: East Azarbaijan, North Khorasan, Tehran ([Ebrahimi & Carpenter, 2008](#)), Kerman ([Pashaei Rad et al., 2010](#)), Khorasan-e Razavi ([Giordani Soika, 1970](#)), Sistan-o Baluchestan ([Hamzavi et al., 2019](#)), Zanjan ([Abbasi et al., 2008](#); [Pashaei Rad et al., 2010](#)).

General distribution: Eastern (China, Iran, Kazakhstan, Tajikistan, Turkmenistan, Uzbekistan) and Western (Armenia, Azerbaijan, Cyprus, Europe, Georgia, Iraq, Israel, Jordan, Lebanon, North Africa, Russia, Turkey) Palaearctic ([Fateryga et al., 2019](#)).

Remarks: The subspecies *Antepipona deflenda deflenda* has been recorded from Iran ([Pashaei Rad et al., 2010](#)).

Antepipona iconia (Blüthgen, 1951)

Odontodynerus iconius Blüthgen, 1951: 169, 182, Holotype ♀. – Turkey.

Distribution in Iran: Mazandaran ([Gusenleitner, 2012a](#)).

General distribution: Eastern (Iran) and Western (Israel, Syria, Turkey) Palaearctic ([Gusenleitner, 2013](#)).

Antepipona karadgensis Giordani Soika, 1970

Antepipona karadgensis Giordani Soika, 1970: 113, ♀. – Iran.

Distribution in Iran: Alborz ([Giordani Soika, 1970](#)).

General distribution: Eastern Palaearctic (Iran) ([Gusenleitner, 2013](#)).

Antepipona laevigata (Blüthgen, 1951)

Odontodynerus laevigatus Blüthgen, 1951: 169, 180, ♂. – Armenia.

Distribution in Iran: Alborz, Mazandaran, Semnan ([Gusenleitner, 1986a](#)).

General distribution: Eastern (Iran, Pakistan) and Western (Armenia, Iraq, Lebanon, Turkey) Palaearctic ([Gusenleitner, 2013](#)).

Antepipona osmania Gusenleitner, 1986

Antepipona osmania Gusenleitner, 1986: 361, ♀. – Turkey.

Distribution in Iran: East Azarbaijan ([Gusenleitner, 2012a](#)).

General distribution: Eastern (Iran) and Western (Turkey) Palaearctic ([Gusenleitner, 2012a](#)).

Antepipona praecincta Giordani Soika, 1970

Antepipona praecincta Giordani Soika, 1970: 117, ♀. – Iran.

Distribution in Iran: Khuzestan ([Giordani Soika, 1970](#)).

General distribution: Eastern Palaearctic (Iran, Pakistan) ([Gusenleitner, 2013](#)).

Antepipona prompta Giordani Soika, 1970

Antepipona prompta Giordani Soika, 1970: 123, ♀. – Iran.

Distribution in Iran: Khuzestan ([Giordani Soika, 1970](#)).

General distribution: Eastern Palaearctic (Iran) ([Gusenleitner, 2013](#)).

***Antepipona specifica* (Morawitz, 1895)**

Odynerus (Lionotus) specificus Morawitz, 1895: 464, ♂♀. – Turkmenistan.

Distribution in Iran: Markazi ([Ebrahimi & Carpenter, 2008](#)).

General distribution: Eastern (Iran, Kyrgyzstan, Turkmenistan, Uzbekistan) and Western Palaearctic (Azerbaijan, Iraq) ([Fateryga et al., 2019](#)).

***Antepipona vagabunda* (Dalla Torre, 1889)**

Odynerus vagabundus Dalla Torre, 1889: 125.

Distribution in Iran: Kerman ([Pashaei Rad et al., 2010](#)), Tehran ([Giordani Soika, 1970](#)), Zanjan ([Abbasi et al., 2008](#); [Pashaei Rad et al., 2010](#)).

General distribution: Eastern Palaearctic (Afghanistan, Iran, Turkmenistan) ([Gusenleitner, 2013](#)).

***Antepipona varentzowi* (Morawitz, 1895)**

Odynerus varentzowi Morawitz, 1895: 474, ♀. – Iran.

Distribution in Iran: North Iran ([Morawitz, 1895](#)).

General distribution: Eastern Palaearctic (Iran - [Morawitz, 1895](#); Kazakhstan, Uzbekistan - [Gusenleitner, 1986a](#)).

Remarks: This species has been recorded from Iran as *Odynerus varentzowi* Morawitz, 1895 ([Morawitz, 1895](#)).

***Antepipona vescovilis* Giordani Soika, 1970**

Antepipona vescovilis Giordani Soika, 1970: 116, ♀. – Iran.

Distribution in Iran: Alborz, Khorasan-e Razavi ([Giordani Soika, 1970](#)).

General distribution: Eastern Palaearctic (Iran - [Gusenleitner, 2013](#); Pakistan - [Gusenleitner, 2006c](#)).

Genus: *Antodynerus* de Saussure, 1855

Antodynerus Saussure, 1855: 242, 287. Type species: *Vespa flavescens* Fabricius, 1775 ("*Odynerus punctum* (Fabricius)" *sensu* Saussure, 1853). Designated by van der Vecht, 1959.

***Antodynerus incomparabilis* Giordani Soika, 1970**

Antodynerus incomparabilis Giordani Soika, 1970: 138, ♀. – Iran.

Distribution in Iran: Fars ([Giordani Soika, 1970](#)).

General distribution: Eastern (Iran) and Western (Turkey) Palaearctic ([Gusenleitner, 2013](#)).

Genus: *Brachyodynerus* Blüthgen, 1938

Brachyodynerus Blüthgen, 1938: 450, 459. Type species: *Odynerus magnificus* Morawitz, 1867. Original designation.

***Brachyodynerus zhelochvtzevi* (Kostylev, 1929)**

Odynerus (Lionotus) zhelochvtzevi Kostylev, 1929: 112, ♂. – Turkestan.

Distribution in Iran: Golestan ([Gusenleitner, 2011b](#)).

General distribution: Eastern Palaearctic (Iran, Kazakhstan) ([Fateryga et al., 2017](#)).

Genus: *Brachypipona* Gusenleitner, 1967

Brachypipona Gusenleitner, 1967: 671. Type species: *Pseudepipona schmidti* Gusenleitner, 1967, by original designation.

***Brachypipona orientalis* Gusenleitner, 2004**

Brachypipona orientalis Gusenleitner, 2004: 1102, Paratype ♀. – Iran.

Distribution in Iran: Hamadan ([Gusenleitner, 2004](#)).

General distribution: Eastern (Iran - [Gusenleitner, 2004](#)) and Western (Turkey - [Gusenleitner, 2013](#)) Palaearctic.

Genus: *Cephalochilus* Blüthgen, 1939

Cephalochilus Blüthgen, 1939: 13. Type species: *Pterochilus grandis* Lepeletier, 1841 (= *Vespa labiata* Fabricius 1798), by original designation.

***Cephalochilus draco* Giordani Soika, 1970**

Cephalochilus draco Giordani Soika, 1970: 40, ♀♂. – Turkey.

Distribution in Iran: Markazi ([Ebrahimi & Carpenter, 2008](#)).

General distribution: Eastern (Iran - [Ebrahimi & Carpenter, 2008](#)) and Western (Turkey, Syria - [Gusenleitner, 2013](#)) Palaearctic.

Genus: *Chlorodynerus* Blüthgen, 1951

Chlorodynerus Blüthgen, 1951: 67, 75 (as subgenus of "Euodynerus Blüthgen"). Type species: *Odynerus chloroticus* Spinola, 1838, by original designation.

***Chlorodynerus araxanus* (Blüthgen, 1956)**

Euodynerus (Chlorodynerus) araxanus Blüthgen, 1956: 281.

Distribution in Iran: Tehran ([Giordani Soika, 1970](#)).

General distribution: Eastern (Iran) and Western (Armenia) Palaearctic ([Gusenleitner, 2013](#)).

***Chlorodynerus flavoferrugineus* Giordani Soika, 1970**

Chlorodynerus flavoferrugineus Giordani Soika, 1970: 170, ♀♂. – Iran.

Distribution in Iran: Khuzestan ([Giordani Soika, 1970](#)).

General distribution: Eastern Palaearctic (Iran) ([Gusenleitner, 2013](#)).

***Chlorodynerus gratus* (Kostylev, 1940)**

Odynerus (Pareudynerus) gratus Kostylev, 1940: 31, ♂. – Iran.

Distribution in Iran: Kerman (Kostylev, 1940).

General distribution: Eastern Palaearctic (Iran) ([Gusenleitner, 2013](#)).

Remarks: This species has been recorded from Iran as *Odynerus (Pareudynerus) gratus* Kostylev, 1940 ([Kostylev, 1940](#)).

***Chlorodynerus infrenis* Giordani Soika, 1970**

Chlorodynerus infrenis Giordani Soika, 1970: 172, ♀♂. – Iran.

Distribution in Iran: Khuzestan ([Giordani Soika, 1970](#)), Sistan-o Baluchestan ([Ebrahimi & Carpenter, 2008](#)).

General distribution: Eastern Palaearctic (Iran) ([Gusenleitner, 2013](#)).

Chlorodynerus loeffleri Giordani Soika, 1958

Chlorodynerus loeffleri Giordani Soika, 1958: 144, ♀. – Iran.

Distribution in Iran: Fars ([Giordani Soika, 1958](#)).

General distribution: Eastern Palaearctic (Iran, Pakistan) ([Gusenleitner, 2013](#)).

Chlorodynerus schulthessianus (Kostylev, 1937)

Odynerus schulthessianus Kostylev, 1937: 226, ♀♂. – Armenia.

Distribution in Iran: Khuzestan, Tehran ([Giordani Soika, 1970](#)).

General distribution: Easten (Iran) and Western (Armenia, Israel) Palaearctic ([Gusenleitner, 2013](#)).

Genus: *Cyphodynerus* van der Vecht, 1971

Cyphodynerus van der Vecht, 1971: 127. Type species: *Odynerus dimidiatus* Spinola, 1838, by original designation.

Remarks: It is not *Odynerus dimidiatus* Guérin, 1834; replaced by *Odynerus canaliculatus* Saussure, 1855).

Cyphodynerus sculpturatus (Dover, 1925)

Odynerus sculpturatus Dover, 1925: 300, ♀. – Karachi [Pakistan].

Distribution in Iran: Kerman ([Kostylev, 1937](#)).

General distribution: Eastern Palaearctic (Iran - [Kostylev, 1937](#), Pakistan - [Dover, 1925](#)).

Remarks: This species has been recorded from Iran as *Odynerus canaliculatus nigroflammeus* Kostylev, 1937 ([Kostylev, 1937](#)).

Genus: *Cyrtolabulus* van der Vecht, 1969

Cyrtolabus van der Vecht, 1963: 11. Type species: *Cyrtolabus suavis* van der Vecht, 1963, by original designation. Replaced by *Cyrtolabulus* van der Vecht, 1969.

Cyrtolabulus iranus (Giordani Soika, 1968)

Cyrtolabus iranus Giordani Soika, 1968: 112, ♀. – Iran.

Distribution in Iran: Fars ([Giordani Soika, 1968](#)).

General distribution: Eastern (Iran - [Giordani Soika, 1968](#)) and Western (Turkey - [Gusenleitner, 1988a](#); [Yildirim & Kojima, 1999](#)) Palaearctic.

Cyrtolabulus karachiensis Gusenleitner, 2006

Cyrtolabulus karachiensis Gusenleitner, 2006: 1298, ♀. – Pakistan.

Distribution in Iran: Sistan-o Baluchestan ([Hamzavi et al., 2019](#)).

General distribution: Eastern Palaearctic (Iran - [Hamzavi et al., 2019](#); Pakistan - [Gusenleitner, 2006c](#)).

Cyrtolabulus syriacus (Giordani Soika, 1968)

Cyrtolabus syriacus Giordani Soika, 1968: 120, ♀♂. – Syria.

Distribution in Iran: Sistan-o Baluchestan ([Hamzavi et al., 2019](#)).

General distribution: Eastern (Iran - [Hamzavi et al., 2019](#); Pakistan) and Western (Israel, Syria) ([Gusenleitner, 2006c](#)) Palaearctic.

***Cyrtolabulus zarudnyi* (Kostylev, 1939)**

Labus zarudnyi Kostylev, 1939: 166, 168, ♀. – Iran.

Distribution in Iran: Kerman ([Kostylev, 1939](#)).

General distribution: Eastern (Iran) and Western (Turkey) Palaearctic ([Gusenleitner, 2013](#)).

Genus: *Delta* de Saussure, 1855

Delta Saussure, 1855: 130, 143. Type species: *Vespa maxillosa* DeGeer, 1773 (= *Vespa emarginata* L., 1758). Designated by Bequaert, 1925.

***Delta anomalam* (Zavattari, 1909)**

Eumenes anomala Zavattari, 1909: 4, ♂. – Kashmir.

Distribution in Iran: Sistan-o Baluchestan ([Blüthgen & Gusenleitner, 1970](#)).

General distribution: Eastern (Iran - [Blüthgen & Gusenleitner, 1970](#); Pakistan) and Western (Egypt - [Gusenleitner, 2013](#)) Palaearctic.

Remarks: This species has been recorded from Iran as *Alfieria anomala* (Zavattari, 1909) ([Blüthgen & Gusenleitner, 1970](#)).

***Delta campaniforme* (Fabricius, 1775)**

Vespa campaniformis Fabricius, 1775: 371, Holotype ♀. – Nova Hollandia.

Distribution in Iran: Fars ([Giordani Soika, 1970](#)), Hormozgan ([Khoobdel et al., 2014](#)), Khuzestan ([Giordani Soika, 1970](#)), Sistan-o Baluchestan ([Blüthgen & Gusenleitner, 1970](#); [Giordani Soika, 1979](#)).

General distribution: Asutralasian (Australia), Eastern Palaearctic (China, Iran, Pakistan), Oceanic (Hawaii – exotic, Papua New Guinea), Oriental (Cambodia; India - [Gusenleitner, 2006b](#); Indonesia; Laos - [Gusenleitner, 2011c](#); Malaysia; Myanmar - [Dover, 1929](#); Nepal, Philippines; Singapore - [Dover & Rao, 1922](#); Thailand, Vietnam) ([Nguyen et al., 2014](#)).

Remarks: The subspecies *Delta campaniforme chloroticum* Giordani Soika, 1979 ([Giordani Soika, 1979](#)) has been recorded from Iran.

***Delta dimidiatipenne* (de Saussure, 1852)**

Eumenes dimidiatipennis de Saussure, 1852: 51, ♀, ♂. – Djidda (Arabie), les Indes Orientales, l’Egypte”.

Distribution in Iran: Alborz, Fars ([Giordani Soika, 1970](#)), Ardabil, Kohgiloyeh-va Boyerahmad, Kordestan, Lorestan, Markazi ([Ebrahimi & Carpenter, 2008](#)), Hormozgan, Tehran ([Blüthgen & Gusenleitner, 1970](#); [Ebrahimi & Carpenter, 2008](#)), Kerman ([Blüthgen & Gusenleitner, 1970](#); [Pashaei Rad et al., 2010](#)), Khuzestan ([Blüthgen & Gusenleitner, 1970](#); [Ebrahimi & Carpenter, 2008](#)), Sistan-o Baluchestan ([Giordani Soika, 1970](#); [Ebrahimi & Carpenter, 2008](#); [Hamzavi et al., 2019](#)), South Khorasan ([Blüthgen & Gusenleitner, 1970](#)), Different regions of Iran ([Esmaili & Rastegar, 1974](#)).

General distribution: Afrotropical (Sudan to South Africa - [Carpenter et al., 2010](#); Saudi Arabia; Oman, Qatar - [Guichard, 1986](#); UAE, Yemen), Oriental (India; Nepal - [Gusenleitner, 1987](#)), Eastern (Afghanistan - [Blüthgen, 1961](#); Iran; Pakistan - [Mahmood et al., 2012](#)) and Western (Canary Islands, North Africa) Palaearctic ([Gusenleitner, 2013](#)).

Remarks: Two subspecies *Delta emarginatus dimidiatipennis* (de Saussure, 1852) ([Blüthgen & Gusenleitner, 1970](#)) and *Eumenes maxillosa dimidiatipennis* de Saussure ([Esmaili & Rastegar, 1974](#)) have been recorded from Iran.

***Delta esuriens* (Fabricius, 1787)**

Vespa esuriens Fabricius 1787: 293, Syntype. – India.

Distribution in Iran: Hormozgan ([Blüthgen & Gusenleitner, 1970](#)), Kerman ([Pashaei Rad et al., 2010](#)), Khuzestan ([Blüthgen & Gusenleitner, 1970](#)), Markazi ([Ebrahimi & Carpenter, 2008](#)), Sistan-o Baluchestan ([Ebrahimi & Carpenter, 2008; Hamzavi et al., 2019](#)).

General distribution: Afro-tropical (African continent - [Carpenter et al., 2010](#); Saudi Arabia; Mauritius - [Gusenleitner & Madl, 2009](#); Oman; Qatar - [Giordani Soika, 1957](#); UAE - [Gusenleitner 2010b](#)), Oriental (India; Laos - [Gusenleitner, 2011c](#); Myanmar, Sri Lanka - [Wickwar, 1908](#); Thailand - [Gusenleitner, 1988b](#); Vietnam - [Nguyen et al., 2014](#)), Eastern (Iran; Pakistan - [Dover, 1925](#)) and Western (Iraq - [Giordani Soika, 1970](#); Israel - [Castro & Dvořák, 2009](#); North Africa - [Gusenleitner, 2011c](#)) Palaearctic ([Gusenleitner, 2013](#)).

Remarks: This species has been recorded from Iran as *Delta companioniforme esuriens* (Fabricius, 1787) ([Blüthgen & Gusenleitner, 1970](#); [Giordani Soika, 1970](#)). Two subspecies, *Delta esuriens esuriens* ([Pashaei Rad et al., 2010](#)) and *Delta esuriens gracile* (de Saussure, 1852) ([Ebrahimi & Carpenter, 2008](#)) have been recorded from Iran.

***Delta hottentotta* (de Saussure, 1852)**

Eumenes hottentotta de Saussure, 1852: 63, ♀. – South Africa.

Distribution in Iran: East Azarbaijan, Fars, Khuzestan ([Ebrahimi & Carpenter, 2008](#)), Kerman ([Pashaei Rad et al., 2010](#)), Sistan-o Baluchestan, South Khorasan ([Blüthgen & Gusenleitner, 1970](#)).

General distribution: Afro-tropical (Burkina Faso, Cameroon, Central African Republic, Democratic Republic of Congo, Ethiopia, Gabon, Gambia, Ivory Coast, Kenya, Mozambique, Namibia, Nigeria, Senegal, South Africa, Tanzania - [Carpenter et al., 2010](#); Saudi Arabia, Oman, UAE, Yemen - [Gusenleitner, 2013](#)), Eastern (Iran - [Ebrahimi & Carpenter, 2008](#)) and Western (Egypt, Jordan, Israel, North Africa - [Gusenleitner, 2013](#)) Palaearctic.

Remarks: The subspecies *Delta hottentotum elegans* (de Saussure, 1852) has been recorded from Iran ([Blüthgen & Gusenleitner, 1970](#); [Ebrahimi & Carpenter, 2008](#); [Pashaei Rad et al., 2010](#)).

***Delta unguiculatum* (Villers, 1789)**

Vespa unguiculata Villers, 1789: 282, ♂. – In Galliae Aust. [France]

Distribution in Iran: Guilan, Mazandaran ([Giordani Soika, 1970](#)), Sistan-o Baluchestan ([Giordani Soika, 1970; Ebrahimi & Carpenter, 2008](#)), Tehran ([Blüthgen & Gusenleitner, 1970](#); [Giordani Soika, 1970](#)).

General distribution: Afro-tropical (Saudi Arabia), Eastern (Iran, Turkmenistan) and Western (Caucasus, Cyprus, Europe, Israel, Lebanon, North Africa, Russia, Syria, Turkey) Palaearctic ([Fateryga, 2018](#)).

Genus: *Discoelius* Latreille, 1809

Discoelius Latreille, 1809: 140. Type species: *Vespa zonalis* Panzer, 1801. Monotypic.

***Discoelius dufouri* Lepeletier, 1841**

Discoelius dufouri Lepeletier, 1841: 605, ♀. – Saint Séver [France].

Distribution in Iran: North Iran ([van der Vecht & Fischer, 1972](#)).

General distribution: Eastern (China, Iran, Kazakhstan, Korea, Mongolia, Russian Far East) and Western Palaearctic (Caucasus, Europe, Russia, Turkey) ([Fateryga, 2018](#)).

***Discoelius pictus* Kostylev, 1940**

Discoelius pictus Kostylev, 1940: 140, ♀♂. – Ak-Tach, chaîne Ala-Tau de Talas, près de Tachkent [Uzbekistan].

Distribution in Iran: No locality cited ([Kurzenko, 1978](#)).

General distribution: Eastern (Afghanistan - [Kurzenko, 1978](#); Iran, Kazakhstan, Kyrgyzstan, Tajikistan, Turkmenistan, Uzbekistan) and Western (Russia) Palaearctic ([Fateryga & Mokrousov, 2019](#)).

***Discoelius zonalis* (panzer, 1801)**

Discoelius zonalis panzer 1801: pl. 18, ♀. – Nürnberg [Germany].

Distribution in Iran: North Iran ([Kurzenko, 1977, 1978](#)).

General distribution: Eastern (China, Iran, Japan, Korea, Mongolia, Russia Far East) and Western (Caucasus, Europe, Russia, Turkey) Palaearctic ([Fateryga, 2018](#)).

Genus: *Eumenes* Latreille, 1802

Eumenes Latreille, 1802: 360. Type species: “*Eumenes coarctata*, Fab.” [= *Vespa coarctata* Linnaeus, 1758], by subsequent designation of Latreille, 1810: 438.

***Eumenes coarctatus* (Linnaeus, 1758)**

Vespa coarctata Linnaeus, 1758: 573, ♀. – Linnean Society, London.

Distribution in Iran: Alborz, Khorasan ([Gusenleitner, 1972](#)), East Azarbaijan ([Ebrahimi & Carpenter, 2008](#)), Kermanshah ([Morice, 1921](#)), Kordestan ([Ebrahimi & Carpenter, 2008](#)), Tehran ([Gusenleitner, 1972; Ebrahimi & Carpenter, 2008](#)).

General distribution: Eastern (China, Iran, Japan, Kazakhstan, Mongolia, Russia Far East) and Western (Caucasus, Cyprus, Europe, Israel, Jordan, North Africa, Russia, Syria, Turkey) Palaearctic ([Fateryga, 2018](#)).

Remarks: This species has been recorded from Iran as *Eumenes lunulatus ordubadensis* Blüthgen, 1938 ([Gusenleitner, 1972; Ebrahimi & Carpenter, 2008](#)) and as *Eumenes coarctatus lunulatus* Fabricius, 1804 ([Ebrahimi & Carpenter, 2008](#)).

***Eumenes comberi* Dover, 1925**

Eumenes comberi Dover, 1925: 293, ♀. – Karachi; Shapali, Bombay Pres.

Distribution in Iran: Kerman ([Pashaei Rad et al., 2010](#)), Khuzestan ([Gusenleitner, 1972](#)), Markazi ([Ebrahimi & Carpenter, 2008](#)).

General distribution: Eastern Palaearctic (Iran, Pakistan, Tajikistan), Oriental (India) ([Kumar et al., 2017](#)).

Remarks: This species has been recorded from Iran as *Eumenes mirandus* Kostylev, 1940 ([Gusenleitner, 1972](#)).

***Eumenes coronatus* (Panzer, 1799)**

Vespa coronata, Panzer, 1799: 12, ♂. – Norimbergae [Germany].

Distribution in Iran: Alborz ([Gusenleitner, 1972](#)); No locality cited ([Kurzenko, 1977](#)).

General distribution: Eastern (Central Asia, China, Iran, Japan, Kazakhstan) and Western (Azerbaijan, Caucasus, Europe, Israel, Russia, Turkey) Palaearctic ([Fateryga et al., 2019](#)).

Remarks: The subspecies *Eumenes coronatus detonsus* Blüthgen, 1943 has been recorded from Iran ([Gusenleitner, 1972](#)).

***Eumenes crimensis* Blüthgen, 1938**

Eumenes dubius crimensis Blüthgen, 1938: 463, ♀♂. – Jalta [Crimea].

Distribution in Iran: Tehran ([Gusenleitner, 1972](#)), Zanjan ([Abbasi et al., 2008; Ebrahimi & Carpenter, 2008; Pashaei Rad et al., 2010](#)).

General distribution: Eastern (Iran, Kazakhstan, Turkmenistan, Uzbekistan) and Western (Russia) Palaearctic ([Fateryga, 2018](#)).

Remarks: The occurrence of this species in Iran was originally indicated by two subspecies including *Eumenes dubius crimensis* Blüthgen, 1938 ([Abbasi et al., 2008; Ebrahimi & Carpenter, 2008; Pashaei Rad et al., 2010](#)) and *Eumenes sareptanus scabrosus* Gusenleitner, 1972 ([Gusenleitner, 1972](#)).

***Eumenes dubius* de Saussure, 1852**

Eumenes dubia de Saussure, 1852: 32, ♀♂. – France.

Distribution in Iran: Kordestan, Markazi ([Ebrahimi & Carpenter, 2008](#)); No locality cited ([Gusenleitner, 1972](#)).

General distribution: Eastern (Iran, Kazakhstan, Tajikistan, Turkmenistan) and Western Palaearctic (Caucasus, Cyprus, Europe, Iraq, Israel, Jordan, Lebanon, Russia, Syria, Turkey), Neotropical (Chile - exotic) ([Fateryga, 2017](#)).

Remarks: Two subspecies, *Eumenes dubius dubius* ([Ebrahimi & Carpenter, 2008](#)) and *Eumenes dubius palaestinensis* Blüthgen, 1938 ([Gusenleitner, 1972](#)) are recorded from Iran.

***Eumenes interpositus* Gusenleitner, 1972**

Eumenes interpositus Gusenleitner, 1972: 103, ♀♂: Holotype ♀. – Iran.

Distribution in Iran: Alborz ([Gusenleitner, 1972](#)).

General distribution: Eastern Palaearctic (Iran, Turkmenistan) ([Gusenleitner, 2013](#)).

***Eumenes jarkandensis* Blüthgen, 1938**

Eumenes mediterraneus jarkandensis Blüthgen, 1938: 487, ♀♂. – Uss-Lusch (1600 m) bei Jarkand (Ostturkestan) gesammelt [Turkestan].

Distribution in Iran: Alborz, Khorasan ([Gusenleitner, 1972](#)), East Azarbaijan, Kordestan ([Ebrahimi & Carpenter, 2008](#)).

General distribution: Eastern (Iran, Kyrgyzstan, Mongolia, Tajikistan, Turkmenistan) and Western (Azerbaijan, Lebanon, Turkey) Palaearctic ([Fateryga et al., 2019](#)).

***Eumenes mediterraneus* Kriechbaumer, 1879**

Eumenis mediterranea Kriechbaumer, 1879: 85, ♀♂. – Dalmatia.

Distribution in Iran: Alborz, Fars, Khorasan ([Gusenleitner, 1972](#)), East Azarbaijan, Guilan, Kerman, Khorasan-e Razavi, Kordestan, Markazi, Mazandaran, Qom ([Ebrahimi & Carpenter, 2008](#)), Isfahan ([Blüthgen & Gusenleitner, 1970; Ebrahimi & Carpenter, 2008](#)), Sistan-o Baluchestan, South Khorasan ([Blüthgen & Gusenleitner, 1970](#)), Tehran ([Gusenleitner, 1972; Esmaili & Rastegar, 1974; Ebrahimi & Carpenter, 2008](#)), Zanjan ([Abbasi et al., 2008; Pashaei Rad et al., 2010](#)).

General distribution: Afrotropical (Oman, Saudi Arabia, UAE, Yemen), Oceanic (Tahiti – exotic), Eastern (Afghanistan, China, Iran, Kazakhstan, Korea, Kyrgyzstan, Mongolia, Pakistan, Uzbekistan), and Western (Caucasus, Cyprus, Europe, Iraq, Israel, Jordan, Lebanon, North Africa, Russia, Syria, Turkey) Palaearctic ([Fateryga, 2018](#)).

Remarks: Two subspecies *Eumenes mediterraneus quettaensis* Cameron, 1907 ([Gusenleitner, 1972](#)) and *Eumenis mediterraneus mediterraneus* ([Ebrahimi & Carpenter, 2008](#); [Pashaei Rad et al., 2010](#)) have been recorded from Iran.

Eumenes modestus Gusenleitner, 2006

Eumenes modestus Gusenleitner, 2006: 1361, ♂. – Iran.

Distribution in Iran: Tehran ([Gusenleitner, 2006a](#)).

General distribution: Eastern (Iran - [Gusenleitner, 2013](#)) and Western (Turkey - [Yildirim & Gusenleitner, 2009](#)) Palaearctic.

Eumenes papillarius (Christ, 1791)

Sphex papillaria Christ, 1791: 325 (destroyed).

Distribution in Iran: East Azarbaijan ([Ebrahimi & Carpenter, 2008](#)), Zanjan ([Abbasi et al., 2008](#); [Pashaei Rad et al., 2010](#)).

General distribution: Eastern (Iran, Kazakhstan, Mongolia, Pakistan) and Western (Azerbaijan, Europe, Israel, Lebanon, Russia, Syria, Turkey) Palaearctic ([Fateryga et al., 2019](#)).

Eumenes punctaticeps Giordani Soika, 1943

Eumenes (Eumenes) punctaticeps Giordani Soika, 1943: 29, ♀♂. – Viterbo [Italy].

Distribution in Iran: North Iran ([Kurzenko, 1977](#)).

General distribution: Eastern (Central Asia, Iran, Kazakhstan) and Western (Europe, Russia, Turkey) Palaearctic ([Antropov & Fateryga, 2017](#)).

Remarks: The subspecies *Eumenes punctaticeps kostylevi* Kurzenko, 1976 has been recorded from Iran ([Kurzenko, 1977](#)).

Eumenes tripunctatus (Christ, 1791)

Sphex tripunctata Christ, 1791: 317 (destroyed).

Distribution in Iran: Markazi ([Ebrahimi & Carpenter, 2008](#)); No locality cited ([Kurzenko, 1977](#)).

General distribution: Eastern (China, Iran, Kazakhstan, Mongolia, Turkmenistan) and Western (Ukraine, Russia) Palaearctic ([Fateryga, 2018](#)).

Eumenes sp.

Distribution in Iran: Hormozgan ([Khoobdel et al., 2014](#)).

Genus: *Euodynerus* Dalla Torre, 1904

Euodynerus Dalla Torre, 1904: 38. Type species: *Vespa dantici* Rossi, 1790. Designated by Blüthgen, 1938 (1937): 277.

Euodynerus (Euodynerus) amplus Gusenleitner, 2011

Euodynerus (Euodynerus) amplus Gusenleitner, 2011: 324, ♂. – Iran.

Distribution in Iran: Kerman ([Gusenleitner, 2011a](#)).

General distribution: Eastern Palaearctic (Iran) ([Gusenleitner, 2013](#)).

Euodynerus (Euodynerus) annae (Kostylev, 1937)

Odynerus annae Kostylev, 1937: 223, ♀. – Iran.

Distribution in Iran: Kerman ([Kostylev, 1937](#)).

General distribution: Eastern Palaearctic (Iran) ([Gusenleitner, 2013](#)).

Euodynerus (Euodynerus) caspicus (Morawitz, 1873)

Leionotus caspicus Morawitz, 1873: 295, ♀♂. – Krasnowodsk [Turkmenistan].

Distribution in Iran: No locality cited ([Antropov & Fateryga, 2017](#)).

General distribution: Eastern (China, Iran, Kazakhstan, Kyrgyzstan, Mongolia, Turkmenistan) and Western (Armenia, Azerbaijan, Russia) Palaearctic ([Antropov & Fateryga, 2017](#)).

Euodynerus (Euodynerus) curictensis Blüthgen, 1940

Euodynerus curictensis Blüthgen, 1940: 210, ♀♂. – Insel KrK in Jugoslavien [Croatia].

Distribution in Iran: Alborz ([Giordani Soika, 1970](#)), Sistan-o Baluchestan ([Blüthgen & Guseleitner, 1970](#)).

General distribution: Eastern (Iran, Kazakhstan, Mongolia) and Western (Armenia, Jordan, Europe, Lebanon, North Africa, Russia, Syria, Turkey) Palaearctic ([Fateryga & Mokrousov, 2019](#)).

Euodynerus (Euodynerus) dantici (Rossi, 1790)

Vespa dantici Rossi, 1790: 89, ♀. – Italia.

Distribution in Iran: Kerman ([Pashaei Rad et al., 2010](#)).

General distribution: Oriental (Taiwan, Vietnam), Eastern (Afghanistan, China, Iran, Japan, Kazakhstan, Korea, Russia, Mongolia) and Western (Caucasus, Europe, Israel (?), Jordan, Lebanon, North Africa, Russia, Syria, Turkey) Palaearctic ([Fateryga, 2018](#)).

Remarks: The subspecies *Euodynerus dantici dantici* has been recorded from Iran ([Pashaei Rad et al., 2010](#)).

Euodynerus (Euodynerus) disconotatus (Lichtenstein, 1884)

Odynerus disconotatus Lichtenstein, 1884: L. Montpellier (? destroyed).

Distribution in Iran: Fars, Khorasan-e Razavi, Tehran ([Giordani Soika, 1970](#)), Hormozgan, Kerman, South Khorasan ([Blüthgen & Guseleitner, 1970](#)), Zanjan ([Abbasi et al., 2008](#); [Pashaei Rad et al., 2010](#)).

General distribution: Afrotropical (Saudi Arabia), Eastern (Afghanistan, Iran, Kazakhstan, Pakistan) and Western (Caucasus, Cyprus, Europe, Iraq, Jordan, Russia, Turkey) Palaearctic ([Fateryga, 2018](#)).

Remarks: This species has been recorded from Iran as *Euodynerus sulfuripes* (Morawitz, 1885) ([Giordani Soika, 1970](#); [Blüthgen & Guseleitner, 1970](#)). The subspecies *Euodynerus disconotatus sulfuripes* (Morawitz, 1885) has been recorded from Iran ([Abbasi et al., 2008](#); [Pashaei Rad et al., 2010](#)).

Euodynerus (Euodynerus) effrenatus Giordani Soika, 1970

Euodynerus (Euodynerus) effrenatus Giordani Soika, 1970: 157, ♀. – Iran.

Distribution in Iran: Fars ([Giordani Soika, 1970](#)).

General distribution: Eastern Palaearctic (Iran) ([Guseleitner, 2013](#)).

Euodynerus (Euodynerus) fastidiosus (de Saussure, 1853)

Odynerus (Leionotus) fastidiosus de Saussure, 1853: 154, 189, ♀. – Algeria.

Distribution in Iran: Alborz, Fars, Khorasan-e Razavi ([Giordani Soika, 1970](#)), Zanjan ([Abbasi et al., 2008](#); [Pashaei Rad et al., 2010](#)).

General distribution: Afro tropical (Oman), Eastern (Afghanistan, China, Iran, Kazakhstan) and Western (Caucasus, Europe, Lebanon, North Africa, Russia, Syria, Turkey) Palaearctic ([Fateryga, 2017](#)).

Euodynerus (Euodynerus) khuzestanicus Giordani Soika, 1970

Euodynerus (Euodynerus) khuzestanicus Giordani Soika, 1970: 155, ♀. – Iran.

Distribution in Iran: Khuzestan ([Giordani Soika, 1970](#)).

General distribution: Eastern Palaearctic (Iran) ([Gusenleitner, 2013](#)).

Euodynerus (Euodynerus) rufinus Blüthgen, 1942

Euodynerus (Euodynerus) rufinus Blüthgen, 1942: 303, ♀. – Transcaspien [Turkmenistan].

Distribution in Iran: Khuzestan ([Giordani Soika, 1970](#)), Tehran ([Ebrahimi & Carpenter, 2008](#)).

General distribution: Afro tropical (UAE), Eastern (China, Iran, Kazakhstan, Mongolia, Turkmenistan, Uzbekistan) and Western (Armenia, Israel, North Africa, Russia) Palaearctic ([Fateryga & Mokrousov, 2019](#)).

Remarks: Two subspecies, *Euodynerus rufinus confictus* Giordani Soika, 1970 ([Giordani Soika, 1970](#)) and *Euodynerus rufinus rufinus* ([Ebrahimi & Carpenter, 2008](#)) are recorded from Iran.

Euodynerus (Euodynerus) semidantici (Giordani Soika, 1952)

Pseudepipona (Euodynerus) semidantici Giordani Soika, 1952: 47, ♂. – Palestina.

Distribution in Iran: Unclear locality cited ([Gusenleitner, 2011a](#)).

General distribution: Eastern (Iran) and Western (Israel, Turkey) Palaearctic ([Gusenleitner, 2013](#)).

Euodynerus (Euodynerus) semisaecularis (Dalla Torre, 1889)

Odynerus semisaecularis Dalla Torre, 1889: 125.

Distribution in Iran: Alborz, Khorasan-e Razavi ([Giordani Soika, 1970](#)), No locality cited ([Gusenleitner, 2013](#)).

General distribution: Afro tropical (UAE - [Gusenleitner, 2010a](#)), Eastern (China - [Castro & Dvořák, 2010](#); Iran - [Gusenleitner, 2013](#); Kazakhstan, Pakistan, Tajikistan, Turkmenistan, Uzbekistan - [Castro & Dvořák, 2010](#)) and Western (Turkey - [Gusenleitner, 2013](#)) Palaearctic.

Remarks: Two subspecies, *Euodynerus semisaecularis macedonicus* Blüthgen, 1951 - as *Euodynerus macedonicus* (Blüthgen) ([Giordani Soika, 1970](#)) and *Euodynerus semisaecularis semisaecularis* (Dalla Torre, 1889) ([Gusenleitner, 2013](#)) have been recorded from Iran.

Euodynerus (Euodynerus) setosus Gusenleitner, 1970

Euodynerus (Euodynerus) setosus Gusenleitner, 1970: 11, ♀♂. – Iran.

Distribution in Iran: South Khorasan ([Blüthgen & Gusenleitner, 1970](#)).

General distribution: Afro tropical (Oman - [Gusenleitner, 1994](#)), Eastern Palaearctic (Afghanistan, Iran - [Blüthgen & Gusenleitner, 1970](#)).

Euodynerus (Euodynerus) shirazensis Giordani Soika, 1970

Euodynerus shirazensis Giordani Soika, 1970: 162, ♀. – Iran.

Distribution in Iran: Fars ([Giordani Soika, 1970](#)).

General distribution: Eastern Palaearctic (Iran) ([Gusenleitner, 2013](#)).

Euodynerus (Pareuodynerus) breviventris (Giordani Soika, 1952)

Pseudepipona (Euodynerus) breviventris Giordani Soika, 1952: 383, ♂. – Turkey.

Distribution in Iran: Fars ([Giordani Soika, 1970](#)).

General distribution: Eastern (Iran) and Western (Turkey) Palaearctic ([Gusenleitner, 2013](#)).

Euodynerus (Pareuodynerus) enodatus (Kostylev, 1940)

Odynerus (Pareuodynerus) enodatus Kostylev, 1940: 32, ♀. – Beloutchistan.

Distribution in Iran: Sistan-o Baluchestan ([Hamzavi et al., 2019](#)), No locality cited ([Kostylev, 1940](#)).

General distribution: Eastern Palaearctic (Afghanistan, Iran, Pakistan) ([Gusenleitner, 2013](#)).

Euodynerus (Pareuodynerus) notatus (Jurine, 1807)

Vespa notata Jurine, 1807: 170, ♀. – Geneve [Switzerland].

Distribution in Iran: Mazandaran ([Castro & Dvořák, 2009](#)).

General distribution: Eastern (China, Iran, Korea ([Castro & Dvořák, 2009](#)), Mongolia, Russia) and Western (Europe, Caucasus, North Africa, Russia, Turkey) Palaearctic ([Antropov & Fateryga, 2017](#)).

Remarks: This species has been recorded from Iran as *Euodynerus notatus notatus* ([Castro & Dvořák, 2009](#)).

Euodynerus (Pareuodynerus) posticus (Herrich-Schäffer, 1841)

Odynerus posticus Herrich-Schäffer, 1841: 176, ♀.

Distribution in Iran: Lorestan, Mazandaran, Sistan-o Baluchestan, Tehran ([Ebrahimi & Carpenter, 2008](#)), Guilan ([Blüthgen & Gusenleitner, 1970](#); [Ebrahimi & Carpenter, 2008](#)), Zanjan ([Abbasi et al., 2008](#); [Pashaei Rad et al., 2010](#)).

General distribution: Eastern (Iran) and Western (Armenia, Azerbaijan, Europe, Georgia, Israel, North Africa, Russia, Turkey) Palaearctic ([Fateryga et al., 2019](#)).

Remarks: This species has been recorded from Iran as *Euodynerus posticus punctatissimus* ([Giordani Soika, 1952](#)) ([Blüthgen & Gusenleitner, 1970](#)).

Genus: *Eustenancistrocerus* Blüthgen, 1938

Eustenancistrocerus Blüthgen, 1938: 443,460. Type species: *Odynerus blanchardianus* Saussure, 1855. by original designation.

Eustenancistrocerus askhabadensis (Radoszkowski, 1886)

Odynerus (Lionotus) askhabadensis Radoszkowski, 1886: 47, ♀♂. – Askhabad.

Distribution in Iran: Golestan, Markazi, Sistan-o Baluchestan ([Ebrahimi & Carpenter, 2008](#)), Tehran ([Giordani Soika, 1970](#); [Ebrahimi & Carpenter, 2008](#)).

General distribution: Eastern Palaearctic (Iran, Pakistan, Turkmenistan - [Gusenleitner, 2013](#); Uzbekistan - [Giordani Soika, 1943](#)).

Eustenancistrocerus inconstans (de Saussure, 1863)

Odynerus (Stenancistrocerus) inconstans de Saussure, 1863: 217, ♀. – Abyssinie.

Distribution in Iran: Mazandaran ([Ebrahimi & Carpenter, 2008](#)), Tehran ([Giordani Soika, 1970](#)).

General distribution: Afrotropical (Ethiopia, Oman, Saudi Arabia, UAE, Yemen), Eastern (Central Asia, Iran, Pakistan) and Western (Israel, Jordan, North Africa) Palaearctic ([Gusenleitner, 2006c, 2013](#)).

***Eustenancistrocerus jerichoensis* (Von Schulthess, 1928)**

Odynerus jerichoensis von Schulthess, 1928: 73–74, ♀♂. – Jericho [Israel].

Distribution in Iran: Alborz ([Giordani Soika, 1970](#)), Sistan-o Baluchestan ([Hamzavi et al., 2019](#)).

General distribution: Eastern (Iran) and Western (Azerbaijan, Jordan, Israel, Syria, Russia, Turkey) Palaearctic ([Fateryga & Mokrousov, 2019](#)).

***Eustenancistrocerus malus* Giordani Soika, 1970**

Eustenancistrocerus (Eustenancistrocerus) malus Giordani Soika, 1970: 96, ♂. – Iran.

Distribution in Iran: Fars ([Giordani Soika, 1970](#); [Ebrahimi & Carpenter; 2008](#)), Tehran ([Giordani Soika, 1970](#)).

General distribution: Eastern Palaearctic (Iran) ([Gusenleitner, 2013](#)).

***Eustenancistrocerus spinosissimus* Gusenleitner, 2000**

Eustenancistrocerus (Eustenancistrocerus) spinosissimus Gusenleitner, 2000: 934, ♂. – Iran.

Distribution in Iran: Yazd ([Gusenleitner, 2000c](#)).

General distribution: Eastern Palaearctic (Iran) ([Gusenleitner, 2000c](#); [Gusenleitner, 2013](#)).

***Eustenancistrocerus sulcithorax* Giordani Soika, 1970**

Eustenancistrocerus (Eustenancistrocerus) sulcithorax Giordani Soika, 1970: 93, ♀♂. – Iran.

Distribution in Iran: Fars, Khuzestan, Tehran ([Giordani Soika, 1970](#)).

General distribution: Eastern Palaearctic (Iran) ([Giordani Soika, 1970](#); [Gusenleitner, 2013](#)).

***Eustenancistrocerus tegularis* (Morawitz, 1885)**

Lionotus tegularis Morawitz, 1885: 165, ♀♂. – Transcaucasia [Azerbaijan].

Distribution in Iran: Alborz ([Giordani Soika, 1970](#)), Golestan ([Ebrahimi & Carpenter, 2008](#)), Khorasan-e Razavi ([Giordani Soika, 1970](#); [Pashaei Rad et al., 2006](#)), North Khorasan, Tehran ([Pashaei Rad et al., 2006](#)), Zanjan ([Pashaei Rad et al., 2006](#); [Abbasi et al., 2008](#)), Khorasan ([Kohl & Handlirsch, 1889](#)).

General distribution: Eastern (Iran, Kazakhstan) and Western (Armenia, Azerbaijan, Europe, Israel, Syria, Russia, Turkey) Palaearctic ([Fateryga et al., 2019](#)).

Remarks: This species has been recorded from Iran as *Eustenancistrocerus israelensis* Giordani Soika, 1952 ([Giordani Soika, 1970](#); [Abbasi et al., 2008](#)) and as *Lionotus tegularis* Morawitz, 1885 ([Kohl & Handlirsch, 1889](#)).

***Eustenancistrocerus (Parastenancistrocerus) amadanensis* (de Saussure, 1855)**

Odynerus (Ancistrocerus) amadanensis de Saussure, 1855: 214, ♂. – Amadan, en Perse [Iran].

Distribution in Iran: Alborz ([Giordani Soika, 1970](#)), Bushehr ([Giordani Soika, 1943](#)), East Azarbaijan, Golestan, Markazi, Sistan-o Baluchestan ([Ebrahimi & Carpenter, 2008](#)), Fars ([Giordani Soika, 1970](#)), Khorasan-e Razavi ([Giordani Soika, 1970](#); [Ebrahimi & Carpenter, 2008](#)), Khuzestan ([de Saussure, 1855](#); [Giordani Soika, 1970](#); [Gusenleitner, 2000b](#); [Ebrahimi & Carpenter, 2008](#)), Tehran ([Giordani Soika, 1970](#); [Ebrahimi & Carpenter, 2008](#)), Zanjan ([Abbasi et al., 2008](#); [Pashaei Rad et al., 2010](#)).

General distribution: Afrotropical (Saudi Arabia), Eastern (Central Asia, Iran, Kazakhstan, Kyrgyzstan, Pakistan) and Western (Azerbaijan, Cyprus, Europe, Iraq, Israel, North Africa, Russia, Syria, Turkey) Palaearctic ([Fateryga et al., 2019](#)).

Remarks: This species has been recorded from Iran as *Ancistrocerus iranensis* Giordani Soika, 1943 ([Giordani Soika, 1943](#)), as *Odynerus (Ancistrocerus) amadanensis* de Saussure, 1855 and as *Eustenancistrocerus transitorius* (Moravitz, 1867) ([Giordani Soika, 1970](#)). The subspecies *Eustenancistrocerus amadanensis amadanensis* is recorded from Iran ([Gusenleitner, 2000b](#)).

***Eustenancistrocerus (Parastenancistrocerus) iranicus* Gusenleitner, 2013**
Eustenancistrocerus (Parastenancistrocerus) iranicus Gusenleitner, 2013: 110, ♂. – Iran.

Distribution in Iran: Fars ([Gusenleitner et al., 2013](#)).

Distribution: Eastern Palaearctic (Iran) ([Gusenleitner et al., 2013](#)).

***Eustenancistrocerus (Parancistrocerus) khuzestanicus* Giordani Soika, 1970**
Eustenancistrocerus khuzestanicus Giordani Soika, 1970: 97, ♀♂. – Iran.

Distribution in Iran: Khuzestan ([Giordani Soika, 1970; Ebrahimi & Carpenter, 2008](#)).

Distribution: Eastern Palaearctic (Iran - [Gusenleitner et al., 2013](#); Pakistan - [Gusenleitner, 2006b](#)).

***Eustenancistrocerus* sp.**

Distribution in Iran: South Khorasan ([Blüthgen & Gusenleitner, 1970](#)).

Genus: *Extraepipona* Gusenleitner, 2014

Extraepipona Gusenleitner, 2014: 537. Type species: *Extraepipona occulta* Gusenleitner, 2014, by original designation.

***Extraepipona occulta* Gusenleitner, 2014**

Extraepipona occulta Gusenleitner, 2014: 537, ♀. – Iran.

Distribution in Iran: Fars ([Gusenleitner, 2014](#)).

General distribution: Eastern Palaearctic (Iran) ([Gusenleitner, 2014](#)).

Genus: *Gymnomerus* Blüthgen, 1938

Gymnomerus Blüthgen, 1938 (1937): 286. Type species: *Odynerus laevipes* Shuckard, 1837, by original designation.

***Gymnomerus laevipes* (Shuckard, 1837)**

Odynerus laevipes Shuckard, 1837: 495, ♀♂. – Blackwater, Hampshire [England].

Distribution in Iran: Tehran ([Ebrahimi & Carpenter, 2008](#)).

General distribution: Eastern (Iran, Kazakhstan, Mongolia, Russia ([Antropov & Fateryga, 2017](#))) and Western (Caucasus, Europe, North Africa, Russia ([Antropov & Fateryga, 2017](#)), Turkey, Ukraine ([Yildirim & Kojima, 1999](#)) Palaearctic ([Gusenleitner, 2013](#)).

Genus: *Hemipterochilus* Ferton, 1909

Hemipterochilus Ferton, 1909 (1908): 572. Type species: *Odynerus terricola* Mocsary, 1883 (= *Hemipterochilus bembeciformis terricola*), by monotypy.

***Hemipterochilus hohlbecki* (Kostylev, 1934)**

Pterochilus hohlbecki Kostylev, 1934: 120, ♀. – Transkaspien [Turkmenistan]

Distribution in Iran: Alborz ([Giordani Soika, 1970](#)).

General distribution: Eastern Palaearctic (Afghanistan, Iran, Turkmenistan) ([Gusenleitner, 2013](#)).

***Hemipterochilus punctiventris* (Morawitz, 1885)**

Pterochilus punctiventris Morawitz, 1885: 143, ♀. – Iran.

Distribution in Iran: Semnan ([Morawitz, 1885](#)), Sistan-o Baluchestan ([Blüthgen & Gusenleitner, 1970](#)).

General distribution: Eastern Palaearctic (Iran) ([Gusenleitner, 2013](#)).

Remarks: This species has been recorded from Iran as *Pterochilus punctiventris* Morawitz, 1885 ([Blüthgen & Gusenleitner, 1970](#)). The subspecies *Hemipterochilus punctiventris subquadricolor* Gusenleitner, 1970 has also been recorded from Iran ([Blüthgen & Gusenleitner, 1970](#)).

***Hemipterochilus simplex* Gusenleitner, 2000**

Hemipterochilus simplex Gusenleitner, 2000: 927, ♀♂. – Turkey.

Distribution in Iran: Hamadan ([Gusenleitner, 2000c](#)).

General distribution: Eastern (Iran) and Western (Turkey) Palaearctic ([Gusenleitner, 2013](#)).

Genus: *Ischnogasteroides* Magretti, 1884

Ischnogasteroides Magretti, 1884 (1883): 603. Type species: *Ischnogasteroides flavus* Magretti, 1884 (= *Ischnogasteroides leptogaster flavus*), by monotypy.

***Ischnogasteroides picteti* (de Saussure, 1852)**

Eumenes picteti de Saussure, 1852: 67, ♀♂. – France.

Distribution in Iran: Sistan-o Baluchestan ([Ebrahimi & Carpenter, 2008](#)); No locality cited ([Gusenleitner, 2013](#)).

General distribution: Eastern (Iran - [Ebrahimi & Carpenter, 2008](#); Turkmenistan, Kazakhstan - van der Vecht & Fischer, 1972; Kyrgyzstan - [Gusenleitner, 2011b](#)) and Western (Europe; Turkey - [Gusenleitner, 1988a](#); Yildirim & Kojima, 1999) Palaearctic.

Remarks: The subspecies *Ischnogasteroides picteti tenuis* (Morawitz, 1888) has been recorded from Iran ([Gusenleitner, 2013](#)).

***Ischnogasteroides zarudnyi* (Kostylev, 1939)**

Labus zarudnyi Kostylev, 1939: 166, ♀. – Iran.

Distribution in Iran: Kerman ([Kostylev 1939](#)), Markazi ([Ebrahimi & Carpenter, 2008](#)), South Khorasan ([Blüthgen & Gusenleitner, 1970](#)).

General distribution: Eastern (Afghanistan, Iran - [Gusenleitner, 2013](#)) and Western (Turkey - [Gusenleitner, 1988a](#)) Palaearctic.

Remarks: This species has been recorded from Iran as *Eumenes zarudnyi* Kostylev, 1939 ([Kostylev, 1939](#)).

***Ischnogasteroides* sp.**

Distribution in Iran: Sistan-o Baluchestan ([Blüthgen & Gusenleitner, 1970](#)).

Genus: *Jucancistrocerus* Blüthgen, 1938

Jucancistrocerus Blüthgen, 1938: 442, 460. Type species: *Odynerus jucundus* Mocsary, 1883, by monotypy.

Jucancistrocerus (Eremodynerus) atrofasciatus (Morawitz, 1885)

Lionotus atrofasciatus Morawitz, 1885: 162, ♀♂. – China montibus Alaschanensibus.

Distribution in Iran: No locality cited ([Gusenleitner, 2013](#)).

General distribution: Eastern Palaearctic (Central Asia, Iran - [Gusenleitner, 2013](#); China - [Tan et al., 2018](#)).

Jucancistrocerus caspicus Giordani Soika, 1970

Jucancistrocerus caspicus Giordani Soika, 1970: 102, ♀♂. – Iran.

Distribution in Iran: Alborz ([Giordani Soika, 1970](#)).

General distribution: Eastern (Iran, Kazakhstan) and Western (Armenia, Azerbaijan, Georgia, Europe, Russia, Turkey) Palaearctic ([Fateryga et al., 2019](#)).

Jucancistrocerus citreodecoratus Giordani Soika, 1970

Jucancistrocerus citreodecoratus Giordani Soika, 1970: 104, ♀. – Iran.

Distribution in Iran: Alborz ([Giordani Soika, 1970](#); [Gusenleitner, 2010a](#)), Kordestan, Markazi ([Ebrahimi & Carpenter, 2008](#)), Sistan-o Baluchestan ([Hamzavi et al., 2019](#)).

General distribution: Eastern (Iran) and Western (Turkey, Serbia) Palaearctic ([Gusenleitner, 2013](#)).

Jucancistrocerus jucundus (Mocsary, 1883)

Odyneus (Ancistrocerus) jucundus Mocsary, 1883: 49, ♀♂. – Hungaria centralis (Budapestinum) et Asia minor (Brussa.).

Distribution in Iran: No locality cited ([Ebrahimi & Carpenter, 2008](#)).

General distribution: Eastern (Iran - [Ebrahimi & Carpenter, 2008](#); Turkmenistan) and Western (Europe, Iraq, Turkey) Palaearctic ([Gusenleitner, 2013](#)).

Remarks: This species has originally been recorded from Iran as *Jucancistrocerus jucundus deceptrix* (Morice, 1921) ([Ebrahimi & Carpenter, 2008](#)).

Jucancistrocerus pareumeniformis Giordani Soika, 1973

Jucancistrocerus pareumeniformis Giordani Soika, 1973: 102, ♂ – Iran.

Distribution in Iran: Guilan ([Giordani Soika, 1973](#)).

General distribution: Eastern Palaearctic (Iran) ([Giordani Soika, 1973](#)).

Genus: *Katamenes* Meade-Waldo, 1910

Katamenes Meade-Waldo, 1910: 46. Type species: *Katamenes watsoni* Meade-Waldo, 1910, by monotypy.

Katamenes dimidiatus (Brullé, 1832)

Eumenes dimidiatus Brullé, 1832: 361, ♀. – Plaine de Modon [Greece].

Distribution in Iran: East Azarbaijan, Fars, Markazi ([Ebrahimi & Carpenter, 2008](#)), Tehran ([Giordani Soika, 1970](#)), Zanjan ([Abbasi et al., 2008](#); [Pashaei Rad et al., 2010](#)).

General distribution: Oriental (India - [Kumar et al., 2019](#)), Eastern (Afghanistan, Central Asia, Iran, Kazakhstan, Mongolia, Pakistan) and Western (Caucasus, Cyprus, Europe, Israel, Russia, Syria, Turkey) Palaearctic ([Fateryga, 2018](#)).

Remarks: The subspecies *Katamenes dimidiatus dimidiatus* ([Pashaei Rad et al., 2010](#)) and *Katamenes dimidiatus montanus* ([Nurse, 1904](#)) ([Giordani Soika, 1970](#)) have been recorded from Iran.

Katamenes flavigularis (Blüthgen, 1951)

Eumenes (*Katamenes*) *arbustorum flavigularis* Blüthgen, 1951:176.

Distribution in Iran: Qazvin, Tehran ([Ebrahimi & Carpenter, 2008](#)).

General distribution: Oriental (India - [Kumar et al., 2019](#)), Eastern (Afghanistan, Iran, Tajikistan) and Western (Europe, Caucasus; Israel - [Kumar et al., 2019](#); Lebanon, Russia, Syria, Turkey) Palaearctic ([Antropov & Fateryga, 2017](#)).

Katamenes niger (Brullé, 1839)

Eumenes *niger* Brullé, 1839: 89, ♀♂. – Canaries [Canary Islands].

Distribution in Iran: Unclear locality cited ([Giordani Soika, 1970](#)).

General distribution: Afrotropical (Saudi Arabia), Eastern (Iran - [Giordani Soika, 1970](#)) and Western (Algeria - [Saunders, 1905](#); Canary Islands, Egypt, Jordan, Israel, Morocco) Palaearctic ([Gusenleitner, 2013](#)).

Katamenes sichelii (de Saussure, 1852)

Eumenes *sichelii* de Saussure, 1852: 36, ♀. – Albania.

Distribution in Iran: Alborz ([Giordani Soika, 1949, 1970](#)), Lorestan ([Giordani Soika, 1949](#)), Markazi ([Ebrahimi & Carpenter, 2008](#)), Sistan-o Baluchestan ([Blüthgen & Gusenleitner, 1970](#)), Tehran ([Castro & Dvořák, 2009](#)).

General distribution: Afrotropical (Saudi Arabia, UAE), Eastern (Afghanistan, China, Iran, Kazakhstan, Mongolia, Turkmenistan - [Giordani Soika, 1966](#); Uzbekistan - [Castro & Dvořák, 2009](#)) and Western (Armenia, Azerbaijan, Europe, Iraq, India, Israel, Jordan, North Africa, Russia, Turkey) Palaearctic ([Fateryga et al., 2019](#)).

Remarks: This species has been recorded from Iran as *Eumenes sichelii* var. *baeri* (Radoszkowski, 1865) ([Giordani Soika, 1949](#)). Three subspecies *Katamenes sichelii baerii* (Radoszkowski, 1865) ([Giordani Soika, 1970](#)), *Katamenes sichelii fulvus* (Eversmann, 1854) as *Eumenes* (*Katamenes*) *sichelii* var. *Luristanensis* Giordani Soika, 1949 ([Giordani Soika, 1949](#)) and *Katamenes sichelii tauriae* (Giordani Soika, 1960) ([Blüthgen & Gusenleitner, 1970](#); [Castro & Dvořák, 2009](#)) have been recorded from Iran.

Katamenes tauricus (de Saussure, 1855)

Eumenes *tauricus* de Saussure, 1855: 137, Crimée.

Distribution in Iran: Alborz ([Giordani Soika, 1970](#)), Tehran ([Ebrahimi & Carpenter, 2008](#)); No locality cited ([Giordani Soika, 1949](#)).

General distribution: Oriental (India), Eastern (Afghanistan, China, Iran, Kazakhstan, Kyrgyzstan, Mongolia) and Western (Russia) Palaearctic ([Fateryga, 2018](#)).

Remarks: The subspecies *Katamenes tauricus viratus* (Giordani Soika, 1949) has been recorded from Iran, as *Katamenes sesquicinctus viratus* (Giordani Soika, 1949) ([Giordani Soika, 1970](#)) and as *Eumenes* (*Katamenes*) *latipes* var. *viratus* Giordani Soika, 1949 ([Giordani Soika, 1949](#)).

Genus: *Knemodynerus* Blüthgen, 1940

Knemodynerus Blüthgen, 1940: 43. Type species: *Odynerus excellens* Perez, 1907, by original designation.

***Knemodynerus excellens* (Pérez, 1907)**

Odynerus (Lionotus) excellens Pérez, 1907: 493, ♀♂. – Dibba [Oman], ♀; Bahrein, ♂.

Distribution in Iran: Kerman ([Pashaei Rad et al., 2010](#)), Khuzestan ([Blüthgen & Gusenleitner, 1970](#); [Giordani Soika, 1970](#)), Markazi ([Ebrahimi & Carpenter, 2008](#)), Sistan-o Baluchestan ([Blüthgen & Gusenleitner, 1970](#)), Zanjan ([Abbasi et al., 2008](#); [Pashaei Rad et al., 2010](#)).

General distribution: Afrotopical (Oman, Qatar, Saudi Arabia, UAE - [Gusenleitner, 2013](#)), Eastern Palaearctic (Iran, Turkmenistan - [Gusenleitner, 2013](#); Pakistan - [Gusenleitner, 2006c](#)), Oriental (India) ([Gusenleitner, 2013](#)).

Remarks: This species has been recorded from Iran as *Euodynerus excellens* (Pérez, 1907) ([Giordani Soika, 1970](#)).

Genus: *Labochilus* Blüthgen, 1939

Labochilus Blüthgen, 1939: 12. Type species: *Pterochilus lingarius* Saunders, 1905, by monotypy.

***Labochilus orientalis* Gusenleitner, 2001**

Labochilus orientalis Gusenleitner, 2001: 680, ♀♂: Holotype ♀. – Iran.

Distribution in Iran: Hormozgan ([Gusenleitner, 2001](#)).

General distribution: Eastern (Iran) and Western (Jordan) Palaearctic ([Gusenleitner, 2001](#)).

Genus: *Leptochilus* de Saussure, 1853

Leptochilus Saussure, 1853: 233. Type species: *Pterochilus mauritanicus* Lepeletier, 1841. Designated by Ashmead, 1902.

***Leptochilus (Lionotulus) callidus* (Kostylev, 1940)**

Odynerus (Lionotulus) callidus Kostylev, 1940: 33, ♀♂. – Syr-Daria [Kazakhstan].

Distribution in Iran: Kerman ([Gusenleitner, 2012a](#)).

General distribution: Eastern Palaearctic (Central Asia, Iran, Mongolia) ([Gusenleitner, 2013](#)).

***Leptochilus (Lionotulus) fuscipes* Gusenleitner, 1985**

Leptochilus (Lionotulus) fuscipes Gusenleitner, 1985: 98, ♂. – Iran.

Distribution in Iran: Hamadan, Qazvin ([Gusenleitner, 1985b](#)).

General distribution: Eastern (Iran) and Western (Turkey) Palaearctic ([Gusenleitner, 1985b](#)).

***Leptochilus (Lionotulus) lorestanicus* Gusenleitner, 1995**

Leptochilus (Lionotulus) lorestanicus Gusenleitner, 1995: 174, ♀. – Iran.

Distribution in Iran: Lorestan ([Gusenleitner, 1995a](#)).

General distribution: Eastern Palaearctic (Iran) ([Gusenleitner, 2013](#)).

***Leptochilus (Lionotulus) membranaceus* (Morawitz, 1867)**

Odynerus (Leionotus) membranaceus Morawitz, 1867: 135, ♀. – Gouvernement von Saratov.

Distribution in Iran: Alborz ([Giordani Soika, 1970](#)).

General distribution: Eastern (Iran, Kazakhstan, Turkmenistan, Uzbekistan) and Western (Caucasus, Israel, Europe, Russia, Turkey) Palaearctic ([Fateryga, 2018](#)).

Leptochilus (Lionotulus) neutralis (Giordani Soika, 1943)

Odynerus (Leptochilus) neutralis Giordani Soika, 1943: 12, ♀. – Asia minore, Amasia [Turkey].

Distribution in Iran: Khuzestan ([Giordani Soika, 1961](#)).

General distribution: Eastern (Iran - [Giordani Soika, 1961](#)) and Western (Israel - [Giordani Soika, 1952](#); Syria - [Giordani Soika, 1970](#); Turkey - [Gusenleitner, 2013](#)) Palaearctic.

Leptochilus (Lionotulus) ressli Gusenleitner, 1985

Leptochilus (Lionotulus) ressli Gusenleitner, 1985: 88, ♂. – Iran.

Distribution in Iran: Lorestan ([Gusenleitner, 1985b](#)).

General distribution: Eastern (Iran) and Western (Turkey - [Gusenleitner, 1988a](#)) Palaearctic.

Leptochilus (Lionotulus) simurgh Gusenleitner, 2016

Leptochilus (Lionotulus) simurgh Gusenleitner, 2016: 91, ♀♂. – Iran.

Distribution in Iran: Alborz ([Gusenleitner, 2016](#)).

General distribution: Eastern Palaearctic (Iran) ([Gusenleitner, 2016](#)).

Leptochilus (Neoleptochilus) regulus (de Saussure, 1855)

Odynerus (Odynerus div. Parodynerus) regulus de Saussure, 1855: 247, ♀♂. – Algeria.

Distribution in Iran: East Azarbaijan ([Ebrahimi & Carpenter, 2008](#)).

General distribution: Eastern (Iran) and Western (Caucasus, Cyprus, Jordan, Europe, North Africa, Russia, Syria, Turkey) Palaearctic ([Fateryga, 2017](#)).

Genus: *Microdynerus* Thomson, 1874

Microdynerus Thomson, 1874: 58. Type species: *Odynerus exilis* Herrich-Schäffer, 1839. Designated by Jones, 1937.

Microdynerus (Alastorynerus) microdynerus (Dalla Torre, 1889)

Odynerus microdynerus Dalla Torre, 1889: 124.

Distribution in Iran: No locality cited ([Gusenleitner, 1997a](#)).

General distribution: Eastern (Iran, Kazakhstan, Tajikistan) and Western (Armenia, Azerbaijan, Europe, Georgia, Jordan, Russia, Turkey) Palaearctic ([Fateryga et al., 2019](#)).

Microdynerus (Microdynerus) atriceps Morawitz, 1895

Microdynerus atriceps Morawitz, 1895: 485, ♀. – Transcaucasia, Helenendorf [Azerbaijan].

Distribution in Iran: Qazvin ([Gusenleitner, 1979](#)).

General distribution: Eastern (Iran, Turkmenistan) and Western (Azerbaijan - [Morawitz, 1895](#); Greece - [Gusenleitner, 2008](#); Israel, Turkey) Palaearctic ([Gusenleitner, 2013](#)).

Microdynerus (Pseudomicrodynerus) anatolicus (Blüthgen, 1938)

Pseudomicrodynerus (Pachymicrodynerus) anatolicus Blüthgen, 1938: 456, ♀. – Kleinasiens [without details].

Distribution in Iran: No locality cited ([Ebrahimi & Carpenter, 2008](#)).

General distribution: Eastern (Iran - [Ebrahimi & Carpenter, 2008](#)) and Western Palaearctic (Israel, Syria, Turkey) ([Gusenleitner, 2013](#)).

Microdynerus (Pseudomicromynerus) parvulus (Herrich-Schäffer, 1838)

Odynerus parvulus Herrich-Schäffer, 1838: 19, ♀♂. – bei Regensburg [Germany].

Distribution in Iran: No locality cited ([Gusenleitner, 1979](#)).

General distribution: Eastern (Iran, Kazakhstan, Uzbekistan) and Western (Caucasus, Europe, Russia, Turkey) Palaearctic ([Fateryga, 2018](#)).

Genus: *Odynerus* Latreille, 1802

Odynerus Latreille, 1802: 362. Type species: *Vespa spinipes* L., 1758. Designated by Shuckard, 1837.

Odynerus (Odynerus) ezechiae von Schulthess, 1924

Odynerus (Hoplopus) ezechiae von Schulthess, 1924: 290, ♂. – Constantinople [Turkey], Jerusalem [Israel].

Distribution in Iran: No locality cited ([Gusenleitner, 1998a](#)).

General distribution: Eastern (Iran, Uzbekistan) and Western (Greece, Israel, Jordan, Turkey) Palaearctic ([Gusenleitner, 2013](#)).

Odynerus (Odynerus) femoratus de Saussure, 1856

Odynerus (Epipona div. Hoplopus) femoratus de Saussure, 1856: 310, ♀♂. – Paris [France].

Distribution in Iran: Alborz ([Giordani Soika, 1970](#)).

General distribution: Eastern (Iran) and Western (Caucasus, Europe, Israel, Russia, Syria, Turkey) Palaearctic ([Antropov & Fateryga, 2017](#)).

Odynerus (Odynerus) leucopus (Blüthgen, 1941)

Hoplomerus (Hoplomerus) leucopus Blüthgen, 1941: 310, 331, ♀♂. – Iran.

Distribution in Iran: Bushehr ([Blüthgen, 1941](#)), Ilam ([Giordani Soika, 1970](#); [Ebrahimi & Carpenter, 2008](#)), Kermanshah, Khuzestan ([Giordani Soika, 1970](#)).

General distribution: Eastern (Iran) and Western (Israel, Syria) Palaearctic ([Gusenleitner, 2013](#)).

Remarks: This species has been recorded from Iran as *Hoplomerus leucopus* Blüthgen, 1941 ([Blüthgen, 1941](#)).

Odynerus (Odynerus) melanocephalus (Gmelin, 1790)

Vespa melanocephalus Gmelin, 1790: 2760. Europa (destroyed).

Distribution in Iran: Alborz ([Giordani Soika, 1970](#)), East Azarbaijan, Golestan ([Ebrahimi & Carpenter, 2008](#)), No locality cited ([du Buysson, 1912](#); [Kostylev, 1929](#)).

General distribution: Eastern (Afghanistan, Kazakhstan, Iran, Uzbekistan) and Western (Armenia, Caucasus, Cyprus, Europe, Iraq, Israel, Jordan, Russia, Syria, Turkey) Palaearctic ([Antropov & Fateryga, 2017](#)).

Remarks: This species has been recorded from Iran as *Hoplomerus armeniacus* Morawitz, 1885 ([Kostylev, 1929](#)). The subspecies *Odynerus melanocephalus armeniacus* (Morawitz, 1885) has been recorded from Iran ([Ebrahimi & Carpenter, 2008](#)).

Odynerus (Spinicoxa) albopictus de Saussure, 1856

Odynerus (Epipona div. Hoplopus) albopictus de Saussure, 1856: 312, ♂. – L'ile de Rhodes [Greece].

Distribution in Iran: East Azarbaijan ([Ebrahimi & Carpenter, 2008](#)), Khuzestan ([Blüthgen & Gusenleitner, 1970](#)).

General distribution: Eastern (Iran, Kazakhstan, Turkmenistan, Uzbekistan) and Western (Caucasus, Europe, Iraq, Israel, Jordan, North Africa, Russia, Syria, Turkey) Palaearctic ([Fateryga, 2018](#)).

Remarks: The subspecies *Odynerus albopictus calcaratus* (Morawitz, 1885) has been recorded from Iran ([Blüthgen & Guseinleitner, 1970](#); [Ebrahimi & Carpenter, 2008](#)).

Odynerus (Spinicoxa) nigrospinosis (Morawitz, 1895)

Hoplomerus nigrospinosis Morawitz, 1895: 436, ♀♂. -Transcaspia, Serachs [Turkmenistan].

Distribution in Iran: Khuzestan ([Blüthgen & Guseinleitner, 1970](#)).

General distribution: Eastern (Iran - [Blüthgen & Guseinleitner, 1970](#); Turkmenistan) and Western (Armenia, Egypt, Israel, Turkey) Palaearctic ([Guseinleitner, 2013](#)).

Genus: *Onychopterocheilus* Blüthgen, 1955

Onychopterocheilus Blüthgen, 1955: 407. Type species: *Odynerus daw* Dusmet, 1903, by original designation.

Onychopterocheilus (Acutopterocheilus) spheciformis (Guseinleitner, 1970)

Pterocheilus spheciformis Guseinleitner, 1970: 9, ♀. - Iran.

Distribution in Iran: Khuzestan ([Blüthgen & Guseinleitner, 1970](#)).

General distribution: Eastern Palaearctic (Iran) ([Guseinleitner, 2013](#)).

Onychopterocheilus (Onychopterocheilus) hellenicus (Morawitz, 1885)

Pterochilus hellenicus Morawitz, 1885: 137, ♀. - Greece.

Distribution in Iran: No locality cited ([Guseinleitner, 2013](#)).

General distribution: Eastern (Iran) and Western (Balkans, Lebanon, Italy, Turkey, Syria) Palaearctic ([Guseinleitner, 2013](#)).

Remarks: The subspecies *Onychopterocheilus hellenicus syriacus* (Blüthgen, 1952) has been recorded from Iran ([Guseinleitner, 2013](#)).

Onychopterocheilus mavromoustakisi (Giordani Soika, 1970)

Pterocheilus mavromoustakisi Giordani Soika, 1970: 46, ♀. - Iran.

Distribution in Iran: Alborz ([Giordani Soika, 1970](#)).

General distribution: Eastern (Iran) and Western (Lebanon, Turkey) Palaearctic ([Guseinleitner, 2013](#)).

Remarks: This species has been recorded from Iran as *Pterocheilus mavromoustakisi* Giordani Soika, 1970 ([Giordani Soika, 1970](#)).

Onychopterocheilus stiziformis (Guseinleitner, 1970)

Pterocheilus stiziformis Guseinleitner, 1970: 7, ♀. - Iran.

Distribution in Iran: Sistan-o Baluchestan ([Blüthgen & Guseinleitner, 1970](#)).

General distribution: Afrotropical (UAE), Eastern Palaearctic (Iran) ([Guseinleitner, 2013](#)).

Genus: *Paragymnomerus* Blüthgen, 1938

Paragymnomerus Blüthgen, 1938 (1937): 286. Type species: *Odynerus spiricornis* Spinola, 1808, by original designation.

***Paragymnomerus amitinorum* Blüthgen, 1952**

Paragymnomerus amitinorum Blüthgen, 1952: 4, ♀♂. – Tiberias [Israel].

Distribution in Iran: No locality cited ([Gusenleitner, 2013](#)).

General distribution: Eastern (Iran) and Western (Croatia, Greece, Israel, Turkey) Palaearctic ([Gusenleitner, 2013](#)).

***Paragymnomerus mammillatus* (Morawitz, 1885)**

Hoplomerus mammillatus Morawitz, 1885: 149, ♀. – Iran.

Distribution in Iran: Semnan ([Morawitz, 1885](#)).

General distribution: Eastern Palaearctic (Iran) ([Gusenleitner, 2013](#)).

Remarks: This species has been recorded from Iran as *Hoplomerus mammillatus* Morawitz, 1885 ([Morawitz, 1885](#)).

***Paragymnomerus spiricornis* (Spinola, 1808)**

Odynerus spiricornis Spinola, 1808: 257, ♂. – Genuam [Italy].

Distribution in Iran: No locality cited ([Gusenleitner, 2013](#)).

General distribution: Eastern (Iran - [Gusenleitner, 2013](#); Turkestan - [van der Vecht & Fischer, 1972](#)) and Western (Armenia, Europe, Turkey) Palaearctic ([Gusenleitner, 2013](#)).

Genus: *Paraleptomenes* Giordani Soika, 1970

Paraleptomenes Giordani Soika, 1970: 79. Type species: *Paraleptomenes nurseanus* Giordani Soika, 1970, by original designation.

***Paraleptomenes miniatus* (de Saussure, 1855)**

Odynerus miniatus de Saussure, 1855: 249, ♀. – Les Indes orientales.

Distribution in Iran: Kerman ([Blüthgen & Gusenleitner, 1970](#); [Pashaei Rad et al., 2010](#)).

Distribution: Afrotropical (Mauritius), Eastern Palaearctic (China - [Tan et al., 2018](#); Iran, Pakistan), Oriental (India, Taiwan) ([Rafi et al., 2017](#)).

Remarks: The subspecies *Paraleptomenes miniatus miniatus* (de Saussure, 1855) has been recorded from Iran ([Pashaei Rad et al., 2010](#)).

Genus: *Paravespa* Radoszkowski, 1886

Paravespa Radoszkowski, 1886: 46. Type species: *Hoplomerus komarowi* Radoszkowski, 1886 (= *Hoplomerus quadricolor* Morawitz, 1885), by monotypy.

***Paravespa grandis* (Morawitz, 1885)**

Hoplomerus grandis Morawitz, 1885: 159, ♂. – Persia boreali, Scharud [Iran].

Distribution in Iran: Semnan ([Morawitz, 1885](#)), Sistan-o Baluchestan ([Hamzavi et al., 2019](#)); No locality cited ([Ebrahimi & Carpenter, 2008](#)).

General distribution: Eastern (Afghanistan - [Gusenleitner, 2013](#); Iran, Turkmenistan - [Morawitz, 1885](#)) and Western (Azerbaijan - [Kokujev, 1912](#); Lebanon - [Gusenleitner, 2013](#); Turkey - [Giordani Soika, 1970](#)) Palaearctic.

Remarks: This species has been recorded from Iran as two separate species including *Hoplomerus grandis* Morawitz, 1885 and *Hoplomerus persa* Morawitz, 1885 ([Morawitz, 1885](#)). The subspecies *Paravespa grandis caucasica* (Kokujev, 1912) has been recorded from Iran ([Ebrahimi & Carpenter, 2008](#)).

***Paravespa mimetica* (Von Schulthess, 1924)**

Odynerus mimeticus Von Schulthess, 1924: 284, ♀♂. – Jericho [Israel].

Distribution in Iran: Khuzestan ([Giordani Soika, 1970](#); [Ebrahimi & Carpenter, 2008](#)).

General distribution: Afro-tropical (Saudi Arabia), Eastern (Iran, Turkmenistan) and Western (Armenia - [Bialynicki-Birula, 1926](#); Israel, Turkey) Palaearctic ([Gusenleitner, 2013](#)).

***Paravespa quadricolor* (Morawitz, 1885)**

Hoplomera quadricolor Morawitz, 1885: 146, ♀. – Territorio Transcaspico, Krasnowodsk [Turkmenistan].

Distribution in Iran: Sistan-o Baluchestan ([Bialynicki-Birula, 1926](#); [Ebrahimi & Carpenter, 2008](#)); No locality cited ([Giordani Soika, 1970](#)).

General distribution: Eastern (Afghanistan, Iran, Turkmenistan) and Western (Turkey - [Gusenleitner, 1988b](#)) Palaearctic ([Gusenleitner, 2013](#)).

Remarks: This species has been recorded from Iran as *Hoplomerus quadricolor* Morawitz, 1885 ([Bialynicki-Birula, 1926](#)).

***Paravespa rex* (von Schulthess, 1924)**

Odynerus (Hoplopus) rex von Schulthess, 1924: 284, 285, ♀♂. – Turkestan, Kisilkum.

Distribution in Iran: Alborz ([Giordani Soika, 1970](#)).

General distribution: Eastern (Iran, Kazakhstan, Kyrgyzstan, Uzbekistan) and Western (Russia, Turkey) Palaearctic ([Antropov & Fateryga, 2017](#)).

Genus: *Parodontodynerus* Blüthgen, 1938

Parodontodynerus Blüthgen, 1938 (1937): 280. Type species: *Eumenes ephippium* Klug, 1817, by original designation.

***Parodontodynerus aramaeus* Blüthgen, 1955**

Parodontodynerus aramaeus Blüthgen, 1955: 27, ♀. – Israel.

Distribution in Iran: Fars, Khuzestan ([Giordani Soika, 1961](#)).

General distribution: Eastern (Iran) and Western (Turkey, Syria - [Giordani Soika, 1970](#); Israel) Palaearctic ([Gusenleitner, 2013](#)).

Remarks: This species has originally been recorded from Iran as *Odontodynerus ephippium eremicus* Giordani Soika, 1961 ([Giordani Soika, 1961](#)).

***Parodontodynerus ephippium* (Klug, 1817)**

Eumenes ephippium Klug, 1817: 264. – Dalmatiae insulae [Croatia].

Distribution in Iran: Alborz ([Giordani Soika, 1970](#)), Zanjan ([Pashaei Rad et al., 2006](#); [Abbasi et al., 2008](#)).

General distribution: Eastern (Iran, Kazakhstan, Turkmenistan, Uzbekistan) and Western (Armenia, Azerbaijan, Europe, Lebanon, Russia, Turkey) Palaearctic ([Fateryga et al., 2019](#)).

***Parodontodynerus rufinus* (Kostylev, 1940)**

Odynerus (Parodontodynerus) ephippium rufinus Kostylev, 1940: 30, ♀. – Iran.

Distribution in Iran: Kerman ([Kostylev, 1940](#)).

General distribution: Eastern Palaearctic (Iran) ([Gusenleitner, 2013](#)).

Remarks: This species has been recorded from Iran as *Odynerus (Parodontodynerus) ephippium rufinus* Kostylev, 1940 ([Kostylev, 1940](#)).

Genus: *Pseudepipona* de Saussure, 1856

Pseudepipona de Saussure 1856: 309. Type species: *Odynerus herrichii* Saussure, 1856, by monotypy.

***Pseudepipona (Deuterepipona) tricolor* Gusenleitner, 1976**

Pseudepipona (Pseudepipona) tricolor Gusenleitner, 1976: 115, ♀♂. – Daghestan [Russia].

Distribution in Iran: No locality cited ([Gusenleitner, 2013](#)).

General distribution: Eastern (Iran, Kazakhstan) and Western (Russia) Palaearctic ([Fateryga et al., 2017](#)).

***Pseudepipona (Pseudepipona) herrichii* (de Saussure, 1856)**

Odynerus (*Epipona* div. *Pseudepipona*) *herrichi* de Saussure, 1856: 309, ♀♂. – Regensburg [Germany].

Distribution in Iran: No locality cited ([Gusenleitner, 1998b](#)).

General distribution: Eastern Palaearctic (China; Iran - [Gusenleitner, 1998b](#); Kazakhstan, Kyrgyzstan, Mongolia, Russia, Turkmenistan, Uzbekistan), Nearctic (North America), Western Palaearctic (Armenia, Azerbaijan, Caucasus, Europe, North Africa, Russia, Syria, Turkey) ([Fateryga & Mokrousov, 2019](#)).

***Pseudepipona (Pseudepipona) lativentris* (de Saussure, 1855)**

Pseudepipona (*Odynerus* div. *Epsilon*) *lativentris* de Saussure, 1855: 275, ♀. – Le Midi de la France. Montpellier [France].

Distribution in Iran: Tehran ([Ebrahimi & Carpenter, 2008](#)).

General distribution: Eastern (Central Asia; Iran - [Ebrahimi & Carpenter, 2008](#)) and Western (Armenia, Europe, Lebanon, Israel, North Africa, Turkey) Palaearctic ([Gusenleitner, 2013](#)).

***Pseudepipona (Pseudepipona) peculiaris* (Morawitz, 1895)**

Odynerus (*Lionotus*) *peculiaris* Morawitz, 1895: 469, ♀. – Mons Bogdo [Russia].

Distribution in Iran: Semnan ([Giordani Soika, 1970](#); [Ebrahimi & Carpenter, 2008](#)).

General distribution: Eastern (Iran, Kazakhstan) and Western (Russia) Palaearctic ([Antropov & Fateryga, 2017](#)).

***Pseudepipona (Pseudepipona) priesneri* Gusenleitner, 1970**

Pseudepipona priesneri Gusenleitner, 1970: 10, ♀♂. – Saudi Arabia.

Distribution in Iran: Khuzestan ([Blüthgen & Gusenleitner, 1970](#)).

General distribution: Afrotropical (Saudi Arabia), Eastern Palaearctic (Iran) ([Blüthgen & Gusenleitner, 1970](#)).

***Pseudepipona (Pseudepipona) sellata* (Morawitz, 1885)**

Lionotus sellatus Morawitz, 1885: 172, ♀♂. – Deserto Kirgisorum. Mons Bogdo [Russia].

Distribution in Iran: Southern Iran ([van der Vecht & Fischer, 1972](#)).

General distribution: Eastern (Iran, Kazakhstan, Turkmenistan) and Western (Azerbaijan, Russia, Turkey) Palaearctic ([Fateryga, 2017](#)).

Genus: *Pseudodontodynerus* Blüthgen, 1939

Pseudodontodynerus Blüthgen, 1939: 249. Type species: *Odynerus pretiosus* Dusmet, 1928, by monotypy.

***Pseudodontodynerus peculiariventris* (Giordani Soika, 1952)**

Odontodynerus peculiariventris Giordani Soika, 1952: 50, ♀. – Tiberias [Israel].

Distribution in Iran: Fars ([Giordani Soika, 1970](#)), Sistan-o Baluchestan ([Hamzavi et al., 2019](#)).

General distribution: Eastern (Iran) and Western (Israel) Palaearctic ([Gusenleitner, 2013](#)).

Remarks: This species has been recorded from Iran as *Pseudodontodynerus peculiariventris salsus* (Giordani Soika, 1970) ([Giordani Soika, 1970](#)).

Genus: *Pseudosymmorphus* Blüthgen, 1938

Pseudosymmorphus Blüthgen, 1938 (1937): 293. Type species *Odynerus hindenburgi* Dusmet, 1917, by original designation.

***Pseudosymmorphus angulatus* Gusenleitner & Ockermüller, 2018**

Pseudosymmorphus angulatus Gusenleitner & Ockermüller, 2018: 1092, ♀♂. – Iran.

Distribution in Iran: Lorestan ([Gusenleitner & Ockermüller, 2018](#)).

General distribution: Eastern Palaearctic (Iran) ([Gusenleitner & Ockermüller, 2018](#)).

***Pseudosymmorphus capillus* Gusenleitner & Ockermüller, 2018**

Pseudosymmorphus capillus Gusenleitner & Ockermüller, 2018: 1094, ♀. – Iran.

Distribution in Iran: Fars ([Gusenleitner & Ockermüller, 2018](#)).

General distribution: Eastern Palaearctic (Iran) ([Gusenleitner & Ockermüller, 2018](#)).

Genus: *Psiliglossa* S. S. Saunders, 1872

Psiliglossa S.S Saunders, 1872: 42. New name for *Stenoglossa* de Saussure.

***Psiliglossa algeriensis* E. Saunders, 1905**

Psiloglossa algeriensis E. Saunders, 1905: 400, ♀♂. – Biskar [Algeria].

Distribution in Iran: Hormozgan ([Gusenleitner, 1973a, 2011a](#)); No locality cited ([Gusenleitner, 2013](#)).

General distribution: Afrotropical (UAE - [Gusenleitner, 2010b](#)), Eastern (Iran) and Western (Algeria, Israel; Jordan - [Gusenleitner, 2011a](#); Libya, Tunisia; Turkey - [Gusenleitner, 2011a](#)) Palaearctic ([Castro & Dvořák, 2009](#)).

Remarks: This species has been recorded from Iran as *Psiliglossa anatolica* Giordani Soika, 1979 ([Gusenleitner 2013](#)).

***Psiliglossa pulchra* Morawitz, 1895**

Psiloglossa pulchra Morawitz, 1895: 410, ♀. Transcaspia: Serachs [Turkmenistan].

Distribution in Iran: Kohgiloyeh-va Boyerahmad ([Gusenleitner, 2004](#)); No locality cited ([Ebrahimi & Carpenter, 2008](#)).

General distribution: Eastern (Iran - [Ebrahimi & Carpenter, 2008](#); Kazakhstan, Kyrgyzstan, Turkmenistan) and Western (Armenia) Palaearctic ([Mokrousov & Zryannin, 2015](#)).

Remarks: The subspecies *Psiliglossa pulchra zhelochovtsevi* Panfilov, 1968 has been recorded from Iran ([Gusenleitner, 2004](#)).

Genus: *Pterocheilus* Klug, 1805

Pterocheilus Klug, 1805: 143. Type species: *Vespa phalerata* Panzer, 1797. Designated by Blanchard, 1840.

***Pterochelilus chobauti* Dusmet, 1928**

Pterochelilus chobauti Dusmet, 1928: 109, ♀. – Algeria.

Distribution in Iran: Sistan-o Baluchestan ([Blüthgen & Guseinleitner, 1970](#)).

General distribution: Afro-tropical (Oman - [Guichard, 1986](#); Saudi Arabia), Eastern (Iran) and Western (Algeria, Egypt; Israel - [Giordani Soika, 1970](#); Morocco; Syria - [Giordani Soika, 1970](#); Tunisia) Palaearctic ([Guseinleitner, 2013](#)).

Remarks: The subspecies *Pterochelilus chobauti chlorodynerooides* Guseinleitner, 1970 has been recorded from Iran ([Blüthgen & Guseinleitner, 1970](#)).

***Pterochelilus persicus* Guseinleitner & Fallahzadeh, 2013**

Pterochelilus persicus Guseinleitner & Fallahzadeh, 2013: 109, ♀. – Iran.

Distribution in Iran: Fars ([Guseinleitner et al., 2013](#)).

General distribution: Eastern Palaearctic (Iran) ([Guseinleitner, 2013](#)).

***Pterochelilus tenebricosus* Guseinleitner, 1998**

Pterochelilus tenebricosus Guseinleitner, 1998: 503, ♀. – Iran.

Distribution in Iran: Fars ([Guseinleitner, 1998c](#)).

General distribution: Eastern Palaearctic (Iran) ([Guseinleitner, 2013](#)).

Genus: *Raphiglossa* S. S. Saunders, 1850

Raphiglossa Saunders, 1850: 71. Type species: *Raphiglossa eumenoides* Saunders, 1850. Designated by Ashmead, 1902.

***Raphiglossa eumenoides* S. S. Saunders, 1850**

Raphiglossa eumenoides S. S. Saunders, 1850: 17, ♀♂. – Greece.

Distribution in Iran: Alborz ([Guseinleitner, 2000b](#)).

General distribution: Eastern (Iran) and Western (Armenia, Azerbaijan, Europe, Lebanon, Russia; Turkey - [Giordani Soika, 1970](#)) Palaearctic ([Fateryga et al., 2019](#)).

Remarks: The subspecies *Raphiglossa eumenoides caucasica* Giordani Soika, 1970 has been recorded from Iran ([Guseinleitner, 2000b](#)).

***Raphiglossa formosa* Kostylev, 1940**

Raphiglossa formosa Kostylev, 1940: 139, ♂. – Kabadian [Tajikistan].

Distribution in Iran: Fars ([Giordani Soika, 1970](#)).

General distribution: Eastern (Iran, Tajikistan) and Western (Syria) Palaearctic ([Guseinleitner, 2013](#)).

***Raphiglossa irana* Giordani Soika, 1970**

Raphiglossa irana Giordani Soika, 1970: 28, ♂. – Iran.

Distribution in Iran: Alborz ([Giordani Soika, 1970](#)).

General distribution: Eastern Palaearctic (Iran) ([Guseinleitner, 2013](#)).

Genus: *Rhynchium* Spinola, 1806

Rhynchium Spinola, 1806: 186. Type species: *Rygchium* (!) *europaeum* Spinola, 1806 (= *Vespa oculata* Fabricius, 1781), by monotypy.

***Rhynchium acromum* Giordani Soika, 1952**

Rhynchium acromum Giordani Soika, 1952: 49, ♀. - Karachi [Pakistan].

Distribution in Iran: Kerman ([Pashaei Rad et al., 2010](#)), Sistan-o Baluchestan ([Blüthgen & Gusenleitner, 1970](#)).

General distribution: Eastern Palaearctic (Iran - [Blüthgen & Gusenleitner, 1970](#); Pakistan - [Giordani Soika, 1952](#)).

***Rhynchium cyanopterum* de Saussure, 1852**

Rhynchium cyanopterum de Saussure, 1852: 108, ♀♂. - Egypte.

Distribution in Iran: No locality cited ([du Buysson, 1912](#))

General distribution: Afrotropical (Southern Sahara to West Africa; Saudi Arabia, UAE - [Guichard, 1986](#); Oman - [Gusenleitner, 2005](#); Yemen), Eastern (Iran - [du Buysson, 1912](#)) and Western (Egypt, Jordan, Israel) Palaearctic ([Gusenleitner, 2013](#)).

***Rhynchium oculatum* (Fabricius, 1781)**

Vespa oculata Fabricius, 1781: 463, Italia.

Distribution in Iran: Bushehr ([du Buysson, 1912](#)), Hormozgan ([Blüthgen & Gusenleitner, 1970](#); [Giordani Soika, 1970](#); [Ebrahimi & Carpenter, 2008](#); [Khoobdel et al., 2014](#)), Kerman ([Pashaei Rad et al., 2010](#)).

General distribution: Afrotropical (Saudi Arabia; UAE - [Gusenleitner, 2010b](#); Oman, Yemen - [Giordani Soika, 1957](#)), Eastern (Afghanistan, Iran) and Western (Cyprus, Egypt, Europe, Israel, Jordan, Malta, Turkey) Palaearctic, Oriental (India) ([Gusenleitner, 2013](#)).

Remarks: This species has been recorded from Iran as *Rhynchium distinguendum* du Buysson, 1912 ([du Buysson, 1912](#)), which is considered as a subspecies, *Rhynchium oculatum distinguendum* du Buysson, 1913 ([Blüthgen & Gusenleitner, 1970](#); [Giordani Soika, 1970](#); [Pashaei Rad et al., 2010](#)).

Genus: *Stenancistrocerus* de Saussure, 1863

Stenancistrocerus de Saussure, 1863: 216. Type species: *Odynerus atropos* Lepeletier, 1841. Designated by Bequaert, 1925.

***Stenancistrocerus (Paratropancistrocerus) biblicus* (Giordani Soika, 1952)**

Tachyancistrocerus biblicus Giordani Soika, 1952: 38, ♀♂. - Israel.

Distribution in Iran: Sistan-o Baluchestan ([Hamzavi et al., 2019](#)).

General distribution: Eastern (Iran - [Hamzavi et al., 2019](#)) and Western (Israel - [Gusenleitner, 2013](#)) Palaearctic.

Genus: *Stenodynerus* de Saussure, 1863

Stenodynerus de Saussure, 1863: 228. Type species: *Odynerus chinensis* Saussure, 1863. Designated, by Bohart, 1939.

***Stenodynerus bluethgeni* van der Vecht, 1971**

Stenodynerus bluethgeni van der Vecht, 1971: 131, ♀♂. - France.

Distribution in Iran: No locality cited ([Gusenleitner, 2000a](#)).

General distribution: Eastern (China, Iran, Kazakhstan, Turkmenistan) and Western (Caucasus; Cyprus - [Gusenleitner, 1981](#); Europe, Russia, Turkey) Palaearctic ([Antropov & Fateryga, 2017](#)).

***Stenodynerus chevrieranus* (de Saussure, 1855)**

Odynerus (*Odynerus* div. *Epsilon*) *chevrieranus* de Saussure, 1855: 268, ♀♂. – Les environs de Genève [Switzerland].

Distribution in Iran: East Azarbaijan ([Ebrahimi & Carpenter, 2008](#)), Zanjan ([Abbasi et al., 2008; Pashaei Rad et al., 2010](#)).

General distribution: Eastern (Iran, Kazakhstan) and Western (Azerbaijan, Europe, Russia, Turkey) Palaearctic ([Fateryga et al., 2019](#)).

***Stenodynerus chitgarensis* Giordani Soika, 1970**

Stenodynerus chitgarensis Giordani Soika, 1970: 108, ♀♂. – Iran.

Distribution in Iran: Alborz ([Giordani Soika, 1970](#)), Tehran ([Giordani Soika, 1970; Ebrahimi & Carpenter, 2008](#)), Zanjan ([Abbasi et al., 2008; Pashaei Rad et al., 2010](#)).

General distribution: Eastern (Iran, Kyrgyzstan, Uzbekistan) and Western (Azerbaijan, Turkey) Palaearctic ([Fateryga et al., 2019](#)).

***Stenodynerus cyrus* Gusenleitner, 2012**

Stenodynerus cyrus Gusenleitner, 2012: 1131, ♀. – Iran.

Distribution in Iran: Kerman ([Gusenleitner, 2012a](#)).

General distribution: Eastern Palaearctic (Iran) ([Gusenleitner, 2013](#)).

***Stenodynerus fastidiosissimus* (de Saussure, 1855)**

Odynerus (*Odynerus* div. *Epsilon*) *fastidiosissimus* de Saussure, 1855: 265, ♀♂. – L'Algérie ou l'Europe méridionale.

Distribution in Iran: Kohgiloyeh-va Boyerahmad ([Gusenleitner, 2012a](#)); No locality cited ([Gusenleitner, 2000a](#)).

General distribution: Eastern (Central Asia, Iran) and Western (Iraq, Israel, Libya, North Africa, Mediterranean Islands, Southern Europe, Russia ([Antropov & Fateryga, 2017](#)); Turkey) Palaearctic ([Castro & Dvořák, 2009](#)).

Remarks: The subspecies *Stenodynerus fastidiosissimus difficilis* (Morawitz, 1867) has been recorded from Iran ([Gusenleitner, 2000a](#)).

***Stenodynerus punctifrons* (Thomson, 1874)**

Odynerus (*Leionotus*) *Lionotuspunctifrons* Thomson, 1874: 57, ♀. – Helvetia.

Distribution in Iran: No locality cited ([Kim & Yamane, 2004](#)).

General distribution: Eastern (Kazakhstan, Kyrgyzstan, Mongolia, Russia) and Western (Caucasus, Europe, Russia, Turkey) Palaearctic ([Fateryga & Mokrousov, 2019](#)).

***Stenodynerus sapidus* Giordani Soika, 1970**

Stenodynerus sapidus Giordani Soika, 1970: 110, ♀. – Iran.

Distribution in Iran: Alborz ([Giordani Soika, 1970](#)).

General distribution: Eastern (Iran) and Western (Russia) Palaearctic ([Fateryga & Popov, 2017](#)).

***Stenodynerus trotzinai* (Morawitz, 1895)**

Odynerus trotzinai Morawitz, 1895: 463, ♀. – Transcaspia [Turkmenistan].

Distribution in Iran: Sistan-o Baluchestan ([Hamzavi et al., 2019](#)).

General distribution: Eastern (Iran - [Hamzavi et al., 2019](#); Pakistan - [Gusenleitner, 2006c](#); Turkmenistan - [Morawitz, 1895](#)) and Western (Turkey - [Yildirim & Gusenleitner, 2012](#)) Palaearctic.

***Stenodynerus xanthomelas* (Herrich-Schäffer, 1839)**

Odynerus xanthomelas Herrich-Schäffer, 1839: 13, ♀♂.

Distribution in Iran: No locality cited ([Ebrahimi & Carpenter, 2008](#)).

General distribution: Eastern (China, Iran) and Western (Caucasus, Europe, Russia, Turkey) Palaearctic ([Fateryga, 2018](#)).

Genus: *Symmorphus* Wesmael, 1836

Symmorphus Wesmael, 1836: 45. Type species: *Odynerus elegans* Wesmael, 1833. Designated by Richards, 1935.

***Symmorphus (Symmorphus) crassicornis* (Panzer, 1798)**

Vespa crassicornis Panzer, 1798: 53, ♀. – Austria.

Distribution in Iran: No locality cited ([Ebrahimi & Carpenter, 2008](#)).

General distribution: Oriental (India), Eastern (China, Iran, Kazakhstan; Kyrgyzstan - [Cumming, 1989](#); Mongolia; Tajikistan- [Cumming, 1989](#)) and Western (Caucasus, Europe, Russia, Turkey) Palaearctic ([Fateryga, 2018](#)).

***Symmorphus (Symmorphus) gracilis* (Brullé, 1832)**

Odynerus gracilis Brullé 1832: 362, ♂. – foret de Koubeh [Greece].

Distribution in Iran: No locality cited ([Ebrahimi & Carpenter, 2008](#)).

General distribution: Eastern (Iran, Kazakhstan, Tajikistan, Turkmenistan) and Western (Caucasus, Europe, Israel, Lebanon, Russia, Syria, Turkey) Palaearctic ([Fateryga, 2018](#)), Oriental (India - [Cumming, 1989](#)).

***Symmorphus (Symmorphus) murarius* (Linnaeus, 1758)**

Vespa muraria Linnaeus, 1758: 573, ♀. – Insecten von Teutschland.

Distribution in Iran: No locality cited ([Ebrahimi & Carpenter, 2008](#)).

General distribution: Eastern (Iran, Kazakhstan, Korea, Russia) and Western (Caucasus, Europe, North Africa, Russia, Turkey) Palaearctic ([Fateryga, 2018](#)).

Genus: *Syneuodynerus* Blüthgen, 1951

Syneuodynerus Blüthgen, 1951: 67. Type species: *Odynerus egregius* Herrich-Schäffer, 1839, by original designation.

***Syneuodynerus egregius* (Herrich-Schäffer, 1839)**

Odynerus egregius Herrich-Schäffer, 1839: 10, ♀. – Spain.

Distribution in Iran: Hormozgan ([Gusenleitner, 1997b](#)).

General distribution: Eastern (Iran, Turkmenistan) and Western (Caucasus, Cyprus, Europe, Israel, Jordan, North Africa, Russia, Syria, Turkey) Palaearctic ([Fateryga, 2018](#)).

Remarks: The occurrence of this species in Iran was indicated by the subspecies *Syneuodynerus egregius egredior* Gusenleitner, 1970 ([Gusenleitner, 1997b](#)).

Genus: *Tachyancistrocerus* Giordani Soika, 1952

Tachyancistrocerus Soika, 1952: 37. Type species: *Odynerus rhodensis* Saussure, 1855, by original designation.

***Tachyancistrocerus komarowi* (Morawitz, 1885)**

Ancistrocerus komarowi Morawitz, 1885: 175, ♀. – Territorio transcaspico. Asschabad [Ashkhabad, Turkmenistan].

Distribution in Iran: Bushehr ([du Buysson, 1912](#); [Giordani Soika, 1943](#)), Fars, Khuzestan ([Giordani Soika, 1970](#)), Kerman ([Pashaei Rad et al., 2010](#)).

General distribution: Afrotropical (Oman), Eastern (Afghanistan, Iran, Kazakhstan, Pakistan, Turkmenistan) and Western (Turkey) Palaearctic ([Gusenleitner, 2006c](#)).

Remarks: This species has been recorded from Iran as *Odynerus komarowi* Morawitz, 1885 ([du Buysson, 1912](#)) and as *Pseudonortonia bushirensis* Giordani Soika, 1943 ([Giordani Soika, 1943](#)).

***Tachyancistrocerus pallidenotatus* Giordani Soika, 1970**

Tachyancistrocerus pallidenotatus Giordani Soika, 1970: 90, ♀. – Iran.

Distribution in Iran: Fars ([Giordani Soika, 1970](#)).

General distribution: Eastern Palaearctic (Iran) ([Gusenleitner, 2013](#)).

***Tachyancistrocerus quabosi* Giordani Soika, 1979**

Tachyancistrocerus quabosi Giordani Soika, 1979: 278.

Distribution in Iran: Sistan-o Baluchestan ([Hamzavi et al., 2019](#)).

General distribution: Afrotropical (Oman, UAE - [Gusenleitner, 2013](#)), Eastern Palaearctic (Iran - [Hamzavi et al., 2019](#)).

***Tachyancistrocerus schmidti* (Kokujev, 1913)**

Odynerus (Ancistrocerus) schmidti Kokujev, 1913: 4, ♀. – Caucasus (Geok-tapa) [Azerbaijan].

Distribution in Iran: Alborz, Khorasan-e Razavi, Tehran ([Giordani Soika, 1970](#)).

General distribution: Eastern (Afghanistan, Iran, Kyrgyzstan, Turkmenistan, Uzbekistan) and Western (Armenia, Azerbaijan, Russia, Turkey) Palaearctic ([Fateryga & Mokrousov, 2019](#)).

Subfamily: Masarinae Latreille, 1802

Genus: *Celonites* Latreille, 1802

Celonites Latreille, 1802: 368. Type species: *Masaris apiformis* Fabricius, 1793 (= *Vespa abbreviata* Villers, 1789), by monotypy.

***Celonites clarus* Gusenleitner, 1973**

Celonites clarus Gusenleitner, 1973: 60, 66, ♀. – Iran.

Distribution in Iran: Tehran ([Gusenleitner, 1973b](#)).

General distribution: Eastern Palaearctic (Iran) ([Gusenleitner, 1973b](#); [Carpenter, 2001](#)).

***Celonites crenulatus* Morawitz, 1888**

Celonites crenulatus Morawitz, 1888: 267, ♀. – Territorio transcaspico Kiltitschinar [Iran].

Distribution in Iran: No locality cited ([Morawitz, 1888](#)).

General distribution: Eastern Palaearctic (Iran, Turkmenistan) ([Carpenter, 2001](#)).

Celonites cyprius de Saussure, 1854

Celonites cyprius de Saussure, 1854: 4, ♂. – Cyprus.

Distribution in Iran: Alborz, Fars, Khorasan-e Razavi ([Gusenleitner, 1973b](#)), Tehran ([Ebrahimi & Carpenter, 2008](#)).

General distribution: Eastern (Iran) and Western (Armenia, Cyprus, Greece, Israel, Lebanon, Syria, Turkey) Palaearctic ([Carpenter, 2001](#)).

Remarks: The subspecies *Celonites cyprius smyrnensis* Richards, 1962 has been recorded from Iran ([Gusenleitner, 1973b](#); [Ebrahimi & Carpenter, 2008](#)).

Celonites discretus Gusenleitner, 1973

Celonites discretus Gusenleitner, 1973: 62, ♀. – Iran.

Distribution in Iran: Fars ([Gusenleitner, 1973b](#)).

General distribution: Eastern Palaearctic (Iran) ([Carpenter, 2001](#)).

Celonites iranus Gusenleitner, 2018

Celonites iranus Gusenleitner, 2018: 306, ♀. – Iran.

Distribution in Iran: Kerman ([Gusenleitner, 2018](#)).

General distribution: Eastern Palaearctic (Iran) ([Gusenleitner, 2018](#)).

Celonites jousseaumei du Buysson, 1906

Celonites jousseaumei du Buysson, 1906: 104, ♀♂. – Obock [Djibouti].

Distribution in Iran: Hormozgan ([Ebrahimi & Carpenter, 2008](#)).

General distribution: Afrotropical (Djibouti, Mali, Oman, Qatar, Senegal, Sudan, Saudi Arabia, UAE, Yemen), Eastern (Iran - [Ebrahimi & Carpenter, 2008](#)) and Western (Algeria, Israel) Palaearctic ([Carpenter, 2001](#)).

Celonites longipilis Gusenleitner, 1973

Celonites longipilis Gusenleitner, 1973: 59, ♀. – Iran.

Distribution in Iran: Fars ([Gusenleitner, 1973b](#)).

General distribution: Eastern Palaearctic (Iran) ([Gusenleitner, 1973b](#); [Carpenter, 2001](#)).

Celonites persicus Richards, 1962

Celonites persicus Richards, 1962: 241, ♀. – Iran.

Distribution in Iran: Khuzestan ([Richards, 1962](#)).

General distribution: Eastern Palaearctic (Iran) ([Richards, 1962](#); [Carpenter, 2001](#)).

Celonites pictus Richards, 1962

Celonites pictus Richards, 1962: 226, ♂. – Senegal.

Distribution in Iran: Hormozgan ([Gusenleitner, 1992](#)).

General distribution: Afrotropical (Oman, Saudi Arabia, Senegal), Eastern (Iran) and Western (Morocco) Palaearctic ([Carpenter, 2001](#)).

Remarks: The subspecies *Celonites pictus rufiventris* Gusenleitner, 1992 was described from Iran ([Gusenleitner, 1992](#)).

Celonites rothschildi du Buysson, 1906

Celonites rothschildi du Buysson, 1906: 105, ♂. – Afrique orientale anglaise: Lesammise, Rendile [Kenya].

Distribution in Iran: Isfahan ([Gusenleitner, 1992](#)).

General distribution: Afrotopical (Kenya), Eastern Palaearctic (Iran) ([Carpenter, 2001](#)).

***Celonites semenovi* Kostylev, 1935**

Celonites (Eucelonites) semenovi Kostylev, 1935: 112 - Persia [Iran].

Distribution in Iran: No locality cited ([Kostylev, 1935](#)).

General distribution: Eastern Palaearctic (Iran) ([Kostylev, 1935; Carpenter, 2001](#)).

***Celonites tauricus* Kostylev, 1935**

Celonites (Celonites) abbreviatus tauricus Kostylev, 1935: 108, ♀. - [Crimea].

Distribution in Iran: Alborz ([Mauss et al., 2016](#)).

General distribution: Eastern (Iran) and Western (Armenia, Azerbaijan, Cyprus, Europe, Russia, Syria, Turkey) Palaearctic ([Fateryga, 2018](#)).

Genus: *Ceramius* Latreille, 1810

Ceramius Latreille, 1810: 329,438. Type species: *Ceramius fonscolombei* Latreille, 1810, by original designation.

***Ceramius caucasicus* André, 1884**

Ceramius caucasicus André, 1884: 820, ♂. - Caucase [Armenia].

Distribution in Iran: Alborz ([Gusenleitner, 1973b](#)), East Azarbaijan ([Ebrahimi & Carpenter, 2008](#)).

General distribution: Eastern (Iran - [Gusenleitner, 1973b](#)) and Western (Armenia, Turkey) Palaearctic ([Carpenter, 2001](#)).

Genus: *Jugurtia* de Saussure, 1854

Jugurtia de Saussure, July 1854: 19, 37, 42–44. Type species: *Celonites oraniensis* Lepeletier, 1841, by subsequent designation of Ashmead, 1902.

***Jugurtia escalerae* Meade-Waldo, 1910**

Jugurtia escalerae Meade-Waldo, 1910: 33, ♀. - Iran.

Distribution in Iran: Khuzestan ([Meade-Waldo 1910](#)), Kordestan ([Ebrahimi & Carpenter, 2008](#)), Sistan-o Baluchestan ([Blüthgen & Gusenleitner, 1970](#)).

General distribution: Eastern (Iran) and Western (Turkey) Palaearctic ([Carpenter, 2001](#)).

***Jugurtia eurycara* Kostylev, 1935**

Jugurtia eurycara Kostylev, 1935: 94, ♀♂. - Caucasus.

Distribution in Iran: Alborz ([Gusenleitner, 1973b](#)), Markazi ([Ebrahimi & Carpenter, 2008](#)).

General distribution: Eastern (Iran) and Western (Armenia, Turkey) Palaearctic ([Carpenter, 2001](#)).

***Jugurtia irana* Kostylev, 1935**

Jugurtia irana Kostylev, 1935: 94, ♀♂. - Iran.

Distribution in Iran: Alborz ([Gusenleitner, 1973b](#)), Khorasan, Sistan-o Baluchestan ([Kostylev, 1935](#)).

General distribution: Eastern Palaearctic (Iran) ([Carpenter, 2001](#)).

***Jugurtia zarudnyi* Kostylev, 1935**

Jugurtia zarudnyi Kostylev, 1935: 93, ♀♂. – Iran.

Distribution in Iran: Hormozgan ([Gusenleitner, 1973b, 1992](#)), Kerman, Khorasan ([Kostylev, 1935](#)).

General distribution: Eastern Palaearctic (Iran) ([Carpenter, 2001](#)).

Genus: *Masaris* Fabricius, 1793

Masaris Fabricius, 1793: 6, 283, 284. Type species: *Masaris vespiformis* Fabricius, 1793, by subsequent monotypy.

***Masaris agnolii* Selis, 2019**

Masaris agnolii Selis, 2019: 395, ♂. – Iran.

Distribution in Iran: Charmahal-o Bakhtiari ([Selis, 2019](#)).

General distribution: Eastern Palaearctic (Iran) ([Selis, 2019](#)).

***Masaris carli* von Schulthess, 1922**

Masaris carli von Schulthess, 1922: 404.

Distribution in Iran: Fars ([Gusenleitner, 1992, 2002](#)).

General distribution: Eastern Palaearctic (Iran, Kazakhstan, Uzbekistan) ([Carpenter, 2001](#)).

Genus: *Quartinia* André, 1884

Quartinia André, 1884: 811, 822. Type species: *Quartinia dilecta* André, by monotypy.

***Quartinia araxana* Giordani Soika, 1960**

Quartinia araxana Giordani Soika, 1960: 133, ♀♂. – Caucaso.

Distribution in Iran: Alborz ([Gusenleitner, 1985a](#)).

General distribution: Eastern (Iran) and Western (Armenia, Turkey) Palaearctic ([Carpenter, 2001](#)).

***Quartinia atrata* Gusenleitner, 1998**

Quartinia atrata Gusenleitner, 1998: 503, ♀. – Iran.

Distribution in Iran: Fars ([Gusenleitner, 1998c](#)).

General distribution: Eastern Palaearctic (Iran) ([Carpenter, 2001](#)).

***Quartinia irana* Gusenleitner, 2012**

Quartinia irana, Gusenleitner, 2012: 319, ♀ – Iran.

Distribution in Iran: Markazi ([Gusenleitner, 2012b](#)).

General distribution: Eastern Palaearctic (Iran) ([Gusenleitner, 2012b](#)).

***Quartinia soikai* Richards, 1964 [1962]**

Quartinia soikai Richards, 1964 (1962): 84, ♀. – Turkey.

Distribution in Iran: Alborz ([Gusenleitner, 1973b](#)).

General distribution: Eastern (Iran) and Western (Greece, Turkey) Palaearctic ([Carpenter, 2001](#)).

***Quartinia syriaca* Richards, 1964 (1962)**

Quartinia syriaca Richards, 1964 (1962): ♀♂. – Syria.

Distribution in Iran: Fars ([Gusenleitner, 1973b](#)).

General distribution: Eastern (Iran) and Western (Lebanon, Syria) Palaearctic ([Carpenter, 2001](#)).

Remarks: The subspecies *Quartinia syriaca nitens* Gusenleitner, 1973 was described from Iran ([Gusenleitner, 1973b](#)).

Subfamily Polistinae Latreille, 1802**Genus: *Parapolybia* de Saussure, 1854**

Parapolybia de Saussure, 1854: 207. Type species: *Polybia (Parapolybia) indica* Saussure, 1854, by subsequent designation of Bingham, 1897.

***Parapolybia escalerae* (Meade-Waldo, 1911)**

Polybia escalerae Meade-Waldo, 1911: 109, ♀. – Iran.

Distribution in Iran: Bushehr, Kerman, Lorestan ([Ebrahimi & Carpenter, 2008](#)), Khuzestan ([Meade-Waldo, 1911](#)).

General distribution: Eastern (Iran, Pakistan) and Western Palaearctic (Turkey) ([Yildirim & Kojima, 1999](#)).

***Parapolybia persica* (Meade-Waldo, 1911)**

Polybia persica Meade-Waldo, 1911: 108, ♀. – Iran.

Distribution in Iran: Khuzestan ([Meade-Waldo, 1911](#)).

General distribution: Eastern Palaearctic (Iran) ([Kojima & Carpenter, 1997](#)).

Genus: *Polistes* Latreille, 1802

Polistes Latreille, 1802, 363. Type species: *Vespa gallica* Linnaeus, 1767, by subsequent designation of Latreille, 1810.

***Polistes (Gyrostoma) olivaceus* (De Geer, 1773)**

Vespa olivacea De Geer, 1773: 582.

Distribution in Iran: Hormozgan ([Khoobdel et al., 2014](#)).

General distribution: Afrotropical (Madagascar, Mauritius, Oman, Réunion, Tanzania), Asutralasian (Australia - exotic), Eastern Palaearctic (Afghanistan, Iran, Japan (?), Pakistan), Neotropical (Chile - exotic), Oceanic (Fiji, French Polynesia, Hawaii - exotic, Marquesas, Marianas, New Zealand - exotic, Samoa, Tonga, Tuamotu Archipelago, New Caledonia), Oriental (Cambodia, China, India, Indonesia, Laos, Malaysia, Myanmar, Nepal, Philippines, Singapore, Sri Lanka, Taiwan, Thailand, Vietnam), Western Palaearctic (Egypt) ([Carpenter, 1996](#)).

***Polistes (Gyrostoma) rothneyi* Cameron, 1900**

Polistes rothneyi Cameron, 1900: 410, ♂. – Barrackpore, Bengal (Rothney).

Distribution in Iran: Kerman ([Nematy et al., 2016](#)).

General distribution: Eastern Palaearctic (Iran - [Nematy et al., 2016](#)), Oriental (India, Nepal) ([Carpenter, 1996](#)).

Polistes (Gyrostoma) wattii Cameron, 1900

Polistes wattii Cameron, 1900: 416, ♀. – Bengal.

Distribution in Iran: Bushehr, Fars, Ilam, Kermanshah, ([Ebrahimi & Carpenter, 2008](#)), Hormozgan ([Blüthgen & Gusenleitner, 1970](#); [Guiglia, 1976](#); [Ebrahimi & Carpenter, 2008](#)), Kerman ([Ebrahimi & Carpenter, 2008](#); [Pashaei Rad et al., 2010](#)), Khuzestan, Sistan-o Baluchestan ([Blüthgen & Gusenleitner, 1970](#); [Ebrahimi & Carpenter, 2008](#)).

General distribution: Afrotropical (Mauritius, Oman, Saudi Arabia, UAE), Oriental (India), Eastern (Afghanistan, China, Iran, Pakistan; Tajikistan, Turkmenistan - [Castro & Dvořák, 2010](#)) and Western (Iraq) Palaearctic ([Carpenter, 1996](#)).

Remarks: This species has been recorded from Iran as *Polistes (Megapolistes) wattii* Cameron, 1900 ([Blüthgen & Gusenleitner, 1970](#)).

Polistes (Polistes) associa Kohl, 1898

Polistes associa Kohl, 1889: 89, ♂. – Poros [Greece].

Distribution in Iran: Guilan ([Castro & Dvořák, 2010](#)).

General distribution: Oriental (India ?), Eastern (China (?); Iran, Pakistan, Tajikistan) and Western (Azerbaijan, Europe, Israel, Russia, Turkey) Palaearctic ([Antropov & Fateryga, 2017](#)).

Polistes (Polistes) atrimandibularis Zimmermann, 1930

Polistes atrimandibularis Zimmermann, 1930: 611, ♂. – Toblach, [Italy].

Distribution in Iran: No locality cited ([Carpenter, 1996](#)).

General distribution: Eastern (Iran) and Western (Armenia, Europe, Egypt, Russia ([Antropov & Fateryga, 2017](#)), Turkey) Palaearctic ([Carpenter, 1996](#)).

Polistes (Polistes) biglumis (Linnaeus, 1758)

Vespa biglumis Linnaeus, 1758: 573, ♀. – Europe.

Distribution in Iran: Semnan ([Blüthgen & Gusenleitner, 1970](#)).

General distribution: Oriental (India), Eastern (Iran, Kazakhstan, Kyrgyzstan, Korea, Mongolia, Pakistan, Turkmenistan) and Western (Caucasus, Europe, Iraq, Israel, North Africa, Syria, Russia, Turkey) Palaearctic ([Antropov & Fateryga, 2017](#)).

Polistes (Polistes) bischoffi Weyrauch, 1937

Polistes bischoffi Weyrauch, 1937: 274, ♀♂. – Italy.

Distribution in Iran: No locality cited ([Antropov & Fateryga, 2017](#)).

General distribution: Eastern (Iran, Kazakhstan, Mongolia) and Western (Europe, North Africa, Russia, Turkey) Palaearctic ([Antropov & Fateryga, 2017](#)).

Remarks: According to [Neumeyer et al. \(2015\)](#) this species might be *P. albellus* Giordani Soika, 1976.

Polistes (Polistes) bucharensis Erichson, 1849

Polistes bucharensis Erichson, 1849: 307, ♀. – Uzbekistan.

Distribution in Iran: Alborz, Khorasan-e Razavi ([Guiglia, 1977](#)), Fars ([Guiglia, 1976, 1977](#)), Golestan ([Weyrauch, 1939](#)), Kordestan, Markazi ([Ebrahimi & Carpenter, 2008](#)), South Khorasan ([Blüthgen & Gusenleitner, 1970](#)), Tehran ([Blüthgen & Gusenleitner, 1970; Guiglia, 1977](#)), Zanjan ([Abbasi et al., 2008](#); [Pashaei Rad et al., 2010](#)).

General distribution: Eastern (China, Iran, Mongolia, Uzbekistan) and Western (Azerbaijan, Cyprus, Egypt, Greece, Israel, Turkey) Palaearctic ([Schmid-Egger et al., 2017](#)).

Remarks: This species has been recorded from Iran as *Polistes iranus* Guiglia, 1976 ([Guiglia, 1976, 1977](#); [Abbasi et al., 2008](#); [Pashaei Rad et al., 2010](#)), as *Polistes gallica* var. *ornata* Weyrauch, 1938, as *Polistes (Polistes) dominulus bucharensis* Erichson, 1849 ([Abbasi et al., 2008](#); [Pashaei Rad et al., 2010](#)) and as *Polistes gallicus bucharensis* Erichson, 1849 ([Blüthgen & Gusenleitner, 1970](#)).

Polistes (Polistes) chinensis (Fabricius, 1793)

Vespa chinensis Fabricius 1793: 261.

Distribution in Iran: No locality cited ([Antropov & Fateryga, 2017](#)).

General distribution: Oceania (New Zealand - exotic), Eastern (China, Iran, Japan, Kazakhstan, Korea, Mongolia, Russia) and Western (Russia, Turkey) Palaearctic ([Antropov & Fateryga, 2017](#)).

Polistes (Polistes) dominula (Christ, 1791)

Vespa dominula Christ, 1791: 229, ♀♂. - Germany.

Distribution in Iran: East Azarbaijan, Kordestan, Markazi ([Ebrahimi & Carpenter, 2008](#)).

General distribution: Afrotropical (Benin - exotic, Ethiopia, South Africa - exotic), Australasian (Australia - exotic), Nearctic (Canada - exotic, U.S.A - exotic), Neotropical (Chile - exotic), Oriental (India), Eastern (Afghanistan, Caucasus, central Asia, Iran, Mongolia, Pakistan, Russia) and Western (Europe, North Africa, Russia, Syria, Turkey) Palaearctic ([Carpenter, 1996](#)).

Polistes (Polistes) foederata Kohl, 1898

Polistes foederata Kohl, 1898: 90, ♂. - Helenendorf [Azerbaijan].

Distribution in Iran: Mazandaran ([Guiglia, 1976](#)).

General distribution: Eastern (Iran - [Guiglia, 1976](#)) and Western (Caucasus ([Guiglia, 1976](#)), Azerbaijan, Europe, Turkey) Palaearctic ([Schmid-Egger et al., 2017](#)).

Remarks: This species has been recorded from Iran as *Polistes (Leptopolistes) foederatus* Kohl ([Guiglia, 1976](#)).

Polistes (Polistes) gallicus (Linnaeus, 1767)

Vespa gallica Linnaeus, 1767: 949, Holotype ♂. - Europa australi [S Europe].

Distribution in Iran: Ardabil, East Azarbaijan, Fars, Golestan, Guilan, Hamadan, Hormozgan, Ilam, Isfahan, Kerman, Khuzestan, Kohgiloyeh-va Boyerahmad, Lorestan, Markazi, North Khorasan, Qom, ([Ebrahimi & Carpenter, 2008](#)), Kermanshah, Kordestan ([du Buysson, 1912](#); [Ebrahimi & Carpenter, 2008](#)), Khorasan-e Razavi ([Blüthgen & Gusenleitner, 1970](#); [Ebrahimi & Carpenter, 2008](#)), Mazandaran ([Guiglia, 1976, 1977](#); [Ebrahimi & Carpenter, 2008](#)), Sistan-o Baluchestan ([Blüthgen & Gusenleitner, 1970](#); [Ebrahimi & Carpenter, 2008](#)), South Khorasan ([Blüthgen & Gusenleitner, 1970](#)), Tehran ([Ebrahimi & Carpenter, 2008](#)), Zanjan ([Abbasi et al., 2008](#)), Various regions ([Esmaili & Rastegar, 1974](#); [Pashaei Rad et al., 2010](#)); No locality cited ([Morice, 1921](#)).

General distribution: Afrotropical (Ethiopia), Eastern (Afghanistan, China, Iran, Mongolia) and Western (Armenia, Azerbaijan, Europe, Israel, North Africa, Russia, Turkey, Turkmenistan, Ukraine) Palaearctic ([Carpenter, 1996](#)).

Remarks: According to [Schmid-Egger et al. \(2017\)](#) this species might be *P. mongolicus* du Buysson, 1911 *in part*. This species was recorded from Iran as *Polistes (Leptopolistes) omissus* Weyr, 1939 ([Blüthgen & Gusenleitner, 1970](#); [Guiglia, 1976](#)) and as *Polistes gallica* (Linnaeus, 1767) ([Esmaili & Rastegar, 1974](#)) from Iran.

Polistes (Polistes) indicus Stolfa, 1934

Polistes indicus Stolfa, 1934: 47, ♀♂. – Pakistan.

Distribution in Iran: Sistan-o Baluchestan ([Blüthgen & Gusenleitner, 1970](#)).

General distribution: Afrotropical (Oman, UAE), Eastern (Afghanistan, Iran, Pakistan), and Western (Iraq) Palaearctic ([Siddiqui et al., 2015](#)).

Remarks: This species has been recorded from Iran as *Polistes (Leptopolistes) indicus* Stolfa, 1934 ([Blüthgen & Gusenleitner, 1970](#)).

Polistes (Polistes) nimpha (Christ, 1791)

Vespa nimpha Christ, 1791: 232, ♀♂. – Kronberg, Taunus [Germany].

Distribution in Iran: Guilan, Tehran ([Blüthgen & Gusenleitner, 1970](#)), Mazandaran ([Ebrahimi & Carpenter, 2008](#)), Zanjan ([Abbasi et al., 2008](#); [Pashaei Rad et al., 2010](#)).

General distribution: Oriental (India), Eastern (central Asia, China, Iran, Mongolia, Pakistan) and Western (Europe, Caucasus, Iraq, Israel, Madeiras, North Africa, Turkey) Palaearctic ([Carpenter, 1996](#)).

Remarks: The occurrence of this species in Iran was indicated by the subspecies *Polistes nimpha irakensis* Gusenleitner, 1976 ([Abbasi et al., 2008](#); [Pashaei Rad et al., 2010](#)).

Polistes (Polistes) semenowi Morawitz, 1889

Polistes semenowi Morawitz, 1889: 552, syntypes ♀. – Turkmenistan.

Distribution in Iran: Tehran ([Ebrahimi & Carpenter, 2008](#)), Mazandaran ([Guiglia, 1976](#); [Ebrahimi & Carpenter, 2008](#)).

General distribution: Eastern (Iran, Turkmenistan) and Western (Caucasus, Europe; Israel - Castro & Dvořák, 2009; North Africa, Russia, Turkey) Palaearctic ([Carpenter, 1996](#)).

Remarks: This species has been recorded from Iran as *Sulcopolistes semenowi* (Morawitz, 1889) ([Guiglia, 1976](#)) and as *Polistes sulcifer* Zimmermann, 1930 ([Ebrahimi & Carpenter, 2008](#)).

Polistes sp.

Distribution in Iran: No locality cited ([Esmaili & Rastegar, 1974](#)).

Subfamily: Vespinae Latreille, 1802

Genus: *Dolichovespula* Rohwer, 1916

Dolichovespula Rohwer, 1916: 642. Type species: *Vespa maculata* Linnaeus, 1763, by original designation.

Dolichovespula omissa (Bischoff, 1931)

Vespa omissa Bischoff, 1931: 6, ♀. – Germany.

Distribution in Iran: Tehran ([Dvořák et al., 2012](#)).

General distribution: Eastern (Iran) and Western (Europe, Russia, Turkey) Palaearctic ([Carpenter & Kojima, 1997](#)).

***Dolichovespula sylvestris* (Scopoli, 1763)**

Vespa sylvestris Scopoli, 1763: 309 - Slovenia.

Distribution in Iran: East Azerbaijan ([Ebrahimi & Carpenter, 2008](#)), Golestan ([Dvořák et al., 2012](#)), Guilan, Khuzestan ([Archer, 1981](#)), Kohgiloyeh-va Boyerahmad ([Castro & Dvořák, 2009](#)), Mazandaran ([Ebrahimi & Carpenter, 2008; Dvořák et al., 2012](#)), Tehran ([Archer, 1981; Ebrahimi & Carpenter, 2008; Dvořák et al., 2012](#)), West Azerbaijan ([Blüthgen & Gusenleitner, 1970; Archer, 1981](#)); No locality cited ([Birula, 1930](#)).

General distribution: Oriental (India), Eastern (Afghanistan, central Asia, China, Iran, Mongolia, Pakistan) and Western (Armenia, Europe, Russia, Turkey) Palaearctic ([Carpenter & Kojima, 1997](#)).

Remarks: This species has been recorded from Iran as *Dolichovespula sylvestris sumptuosa* (du Buysson, 1905) ([Birula 1930; Blüthgen & Gusenleitner, 1970; Archer, 1981](#)).

Genus: *Vespa* Linnaeus, 1758

Vespa Linnaeus, 1758: 343, 572. Type species: "Vespa crabro, Fab." [= *Vespa crabro* Linnaeus, 1758], by subsequent designation of Latreille, 1810.

***Vespa crabro* Linnaeus, 1758**

Vespa crabro Linnaeus, 1758: 572. "in Europae" (holotype ♀. - Linnaean Society, London).

Distribution in Iran: Golestan ([Ebrahimi & Carpenter, 2008, 2012; Dvořák et al., 2012](#)), Guilan ([Perez, 1910; Morice, 1921; Ebrahimi & Carpenter, 2008](#)), Ilam ([Dvořák et al., 2012](#)), Mazandaran ([Ebrahimi & Carpenter, 2008, 2012; Dvořák et al., 2012](#)), North of Iran ([Esmaili & Rastegar, 1974](#)).

General distribution: Eastern Palaearctic (China, Iran, Japan, Kazakhstan, Korea, Mongolia), Nearctic (North America - exotic), Neotropical (Guatemala - exotic), Oriental (Taiwan), Western Palaearctic (Caucasus, Europe, Russia, Turkey) ([Carpenter & Kojima, 1997](#)).

Remarks: This species has been recorded from Iran as *Vespa crabro caspica* Perez, 1910 ([Perez, 1910; Esmaili & Rastegar, 1974](#)).

***Vespa orientalis* Linnaeus, 1771**

Vespa orientalis Linnaeus, 1771: 540, "Oriente" (holotype ♀. - Linnaean Society, London).

Distribution in Iran: Alborz ([Guiglia, 1976, 1977](#)), Ardabil ([Ebrahimi & Carpenter, 2008, 2012; Dvořák et al., 2012](#)), Bushehr ([Ebrahimi & Carpenter, 2008, 2012; Dvořák et al., 2012](#)), East Azerbaijan ([Ebrahimi & Carpenter, 2008, 2012; Dvořák et al., 2012](#)), Fars ([Guiglia, 1976, 1977; Ebrahimi & Carpenter, 2008, 2012; Dvořák et al., 2012](#)), Golestan ([Ebrahimi & Carpenter, 2008, 2012; Dvořák et al., 2012](#)), Guilan ([Dvořák et al., 2012](#)), Hamadan ([Ebrahimi & Carpenter, 2008, 2012; Dvořák et al., 2012](#)), Hormozgan ([Ebrahimi & Carpenter, 2008, 2012; Dvořák et al., 2012; Khoobdel et al., 2014](#)), Ilam ([Dvořák et al., 2012](#)), Isfahan ([Blüthgen & Gusenleitner, 1970; Dvořák et al., 2012](#)), Kerman ([Blüthgen & Gusenleitner, 1970; Ebrahimi & Carpenter, 2008, 2012; Dvořák et al., 2012](#)), Kermanshah ([Ebrahimi & Carpenter, 2008; Dvořák et al., 2012](#)), Khorasan-e Razavi ([Guiglia, 1976, 1977; Ebrahimi & Carpenter, 2008, 2012; Dvořák et al., 2012](#)), Khuzestan ([du Buysson, 1912; Ebrahimi & Carpenter, 2008, 2012; Dvořák et al., 2012](#)), Kohgiloyeh-va Boyerahmad ([Ebrahimi & Carpenter, 2008; Dvořák et al., 2012](#)), Kordestan ([Ebrahimi & Carpenter, 2008; Dvořák et al., 2012](#)), Lorestan ([du Buysson, 1912; Ebrahimi & Carpenter, 2008, 2012; Dvořák et al., 2012](#)), Markazi ([Ebrahimi & Carpenter, 2008, 2012; Dvořák et al., 2012](#)), Mazandaran ([Dvořák et al., 2012](#)),

North Khorasan ([Blüthgen & Gusenleitner, 1970](#); [Ebrahimi & Carpenter, 2008](#)), Qazvin ([Morice, 1921](#); [Ebrahimi & Carpenter, 2008, 2012](#)), Qom ([Ebrahimi & Carpenter, 2008](#)), Semnan ([Dvořák et al., 2012](#)), Sistan-o Baluchestan ([Blüthgen & Gusenleitner, 1970](#); [Ebrahimi & Carpenter, 2008, 2012](#); [Dvořák et al., 2012](#)), South Khorasan ([Dvořák et al., 2012](#)), Tehran ([du Buysson, 1912](#); [Blüthgen & Gusenleitner, 1970](#); [Ebrahimi & Carpenter, 2008, 2012](#); [Dvořák et al., 2012](#)), West Azerbaijan ([Dvořák et al., 2012](#)), Yazd ([Dvořák et al., 2012](#)), Zanjan ([Abbasi et al., 2008](#); [Ebrahimi & Carpenter, 2012](#); [Dvořák et al., 2012](#)), Different regions of Iran ([Esmaili & Rastegar, 1974](#)).

General distribution: Afrotropical (Bahrein, Ethiopia, Oman, Saudi Arabia, Somalia, UAE, Yemen; Madagascar – exotic), Oriental (India, Nepal), Eastern (Afghanistan, China, Iran, Pakistan, Tajikistan, Turkmenistan, Uzbekistan) and Western (Algeria, Cyprus, Egypt, Europe, Iraq, Israel, Jordan, Lebanon, Libya, Russia, Syria, Turkey) Palaearctic ([Carpenter & Kojima, 1997](#)).

Remarks: This species has been recorded from Iran as *Vespa orientalis aegyptiaca* André, 1884 ([Morice, 1921](#)).

Genus: *Vespula* Thomson, 1869

Vespula Thomson, 1869: 79. Type species: *Vespa austriaca* Panzer, 1799, by subsequent designation of Ashmead, 1902.

Vespula germanica (Fabricius, 1793)

Vespa germanica Fabricius 1793: 256, Type unknown; “Kiliae” (Germany).

Distribution in Iran: East Azerbaijan ([Ebrahimi & Carpenter, 2008](#); [Dvořák et al., 2012](#)), Fars ([Ebrahimi & Carpenter, 2008](#); [Dvořák et al., 2012](#)), Golestan ([Dvořák et al., 2012](#)), Guilan ([Morice, 1921](#); [Blüthgen & Gusenleitner, 1970](#); [Dvořák et al., 2012](#)), Hamadan ([Dvořák et al., 2012](#)), Hormozgan ([Khoobdel et al., 2014](#)), Isfahan ([Ebrahimi & Carpenter, 2008](#); [Dvořák et al., 2012](#)), Kerman ([Ebrahimi & Carpenter, 2008](#); [Dvořák et al., 2012](#)), Kermanshah ([du Buysson, 1912](#); [Morice, 1921](#); [Dvořák et al., 2012](#)), Khorasan-e Razavi ([Guiglia, 1976, 1977](#); [Ebrahimi & Carpenter, 2008](#); [Dvořák et al., 2012](#)), Khuzestan ([Dvořák et al., 2012](#)), Kohgiloyeh-va Boyerahmad ([Ebrahimi & Carpenter, 2008](#); [Dvořák et al., 2012](#)), Kordestan ([du Buysson, 1912](#); [Ebrahimi & Carpenter, 2008](#); [Dvořák et al., 2012](#)), Lorestan ([Ebrahimi & Carpenter, 2008](#); [Dvořák et al., 2012](#)), Markazi ([Ebrahimi & Carpenter, 2008](#); [Dvořák et al., 2012](#)), Mazandaran ([Ebrahimi & Carpenter, 2008](#); [Dvořák et al., 2012](#)), North Khorasan ([Blüthgen & Gusenleitner, 1970](#); [Ebrahimi & Carpenter, 2008](#)), Qazvin ([Morice, 1921](#); [Dvořák et al., 2012](#)), Qom ([Ebrahimi & Carpenter, 2008](#)), Semnan ([Dvořák et al., 2012](#)), Sistan-o Baluchestan ([Ebrahimi & Carpenter, 2008](#)), South Khorasan ([Morice, 1921](#); [Blüthgen & Gusenleitner, 1970](#); [Dvořák et al., 2012](#)), Tehran ([Ebrahimi & Carpenter, 2008](#); [Dvořák et al., 2012](#)), West Azerbaijan ([Ebrahimi & Carpenter, 2008](#); [Dvořák et al., 2012](#)), Yazd ([Dvořák et al., 2012](#)), Zanjan ([Abbasi et al., 2008](#); [Dvořák et al., 2012](#)), Different regions of Iran ([Esmaili & Rastegar, 1974](#)).

General distribution: Afrotropical (Ascension Island, South Africa – exotic), Asutralasian (Australia – exotic), Eastern Palaearctic (Afghanistan, China, Iran, Kazakhstan, Korea, Mongolia, Pakistan, Tajikistan, Turkmenistan, Uzbekistan), Nearctic (Canada and U.S.A. – exotic), Neotropical (Argentina and Chile – exotic), Oceanic (New Zealand – exotic), Oriental (India, Taiwan), Western Palaearctic (Algeria, Caucasus, Europe, Iceland – exotic, Israel, Morocco, Russia, Syria, Tunisia, Turkey) ([Carpenter & Kojima, 1997](#)).

Remarks: This species has been recorded from Iran as *Paravespula germanica* (Fabricius, 1793) ([Blüthgen & Guseinleitner, 1970](#)).

Vespula vulgaris (Linnaeus, 1758)

Vespa vulgaris Linnaeus, 1758: 572, Europa (lectotype ♀. – Linnaean Society, London).

Distribution in Iran: Khorasan-e Razavi ([Dvořák et al., 2012](#)), North Khorasan ([Archer, 1989](#)).

General distribution: Australasian (Australia – exotic), Neotropical (Argentina and Chile - exotic), Oceanic (French Polynesia and New Zealand - exotic), Oriental (India), Eastern (China, Iran, Japan, Kazakhstan, Kyrgyzstan, Korea, Mongolia) and Western (Europe, Georgia, Iceland - exotic, Russia, Turkey) Palaearctic ([Carpenter & Kojima, 1997](#)).

Irrelevant records

A few vespid species are recorded from Iran on the basis of specimens which have never gone through the relevant identification by the experts mentioned as co-authors. The masarine species listed by [Abd-Rabou et al. \(2005\)](#) are obviously elements of the Afrotropical region ([Gess & Gess, 2010](#)) or have restricted distribution areas in southwestern Europe ([Mauss & Castro, 2000](#), Mauss in litt. 2019), so that their occurrence in Iran is particularly unlikely. Considering the unreliable data that appeared in those publications, all the species records from these surveys are excluded from the fauna of Iran, until they may be found and confirmed in future reputable studies ([Table 1](#)).

Discussion

A total of 231 vespid species (235 species including unidentified taxa) belonging to 55 genera and 4 subfamilies are recorded from various parts of Iran. The subfamilies Eumeninae, Masarinae, Polistinae and Vespinae are represented by 184 species in 45 genera, 24 species in 5 genera, 17 species in 2 genera, and 6 species in 3 genera, respectively ([Fig. 1](#)). Two species, *Stenancistrocerus transcaspicus* (Kostylev, 1935) and *Pseudepipona tricolor* Guseinleitner, 1976 were originally described from the Transcaspien region (Farab) and from the Caucasus region (Russia: Dagestan), respectively. These species are listed in [Guseinleitner \(2013\)](#) as elements of the Iranian fauna. Since those areas are now separated from Iran the above mentioned species were also excluded from this checklist.

There are 48 species belonging to the following genera that are exclusively recorded from Iran. *Alastor* Lepeletier (4 species), *Ancistrocerus* Wesmael (1), *Antepipona* de Saussure (2), *Chlorodynerus* Blüthgen (3), *Eumenes* Latreille (1), *Euodynerus* Dalla Torre (5), *Eustenancistrocerus* Blüthgen (4), *Extraepipona* Guseinleitner (1), *Hemipterochilus* Ferton (1), *Labochilus* Blüthgen (1), *Leptochilus* de Saussure (3), *Onychopterocheilus* Blüthgen (1), *Paragymnomerus* Blüthgen (1), *Parodontodynerus* Blüthgen (1), *Pterocheilus* Klug (2), *Raphiglossa* S. S. Saunders (1), *Stenodynerus* de Saussure (1), *Tachyancistrocerus* Giordani Soika (1), *Pseudosymmorphus* Blüthgen (2), *Celonites* Latreille (6), *Jugurtia* de Saussure (2), *Masaris* Fabricius (1), *Quartinia* André (2) and *Parapolybia* de Saussure (1). Fifteen new species have been described during the time period 2000–2019 on the basis of the material collected from Iran, including *Eustenancistrocerus spinosissimus* Guseinleitner, 2000; *Labochilus orientalis* Guseinleitner, 2001; *Eumenes modestus* Guseinleitner, 2006; *Ancistrocerus tenebrosus* Guseinleitner, 2010; *Euodynerus amplus* Guseinleitner, 2011; *Stenodynerus cyrus* Guseinleitner, 2012; *Quartinia irana* Guseinleitner, 2012; *Pterocheilus persicus* Guseinleitner & Fallahzadeh, 2013; *Eustenancistrocerus iranicus* Guseinleitner, 2013; *Extraepipona occulta* Guseinleitner, 2014;


Leptochilus simurgh Gусенлеитнер, 2016; *Celonites iranus* Gусенлеитнер, 2018; *Pseudosymmorphus angulatus* Gусенлеитнер & Оккермюller, 2018; *Pseudosymmorphus capillus* Gусенлеитнер & Оккермюller, 2018 and *Masaris agnolii* Selis, 2019. None of these species are recorded from other countries yet.

The genera *Euodynerus*, *Polistes* and *Eumenes* are the most species-rich groups in the vespid fauna of Iran with 18, 15 and 12 species respectively (Fig. 2). On the other hand, some genera including *Acanthodynerus* Gусенлеитнер, *Antodynerus* de Saussure, *Brachydynerus* Blüthgen, *Brachypipona* Gусенлеитнер, *Cephalochilus* Blüthgen, *Cyphodynerus* van der Vecht, *Extraepipona* Gусенлеитнер, *Gymnomerus* Blüthgen, *Knemodynerus* Blüthgen, *Labochilus* Blüthgen, *Paraleptomenes* Giordani Soika, *Pseudodontodynerus* Blüthgen, *Stenancistrocerus* de Saussure, *Syneodynerus* Blüthgen and *Ceramius* Latreille are each represented by a single species in Iran. The monotypic genus, *Extraepipona* represented by *Extraepipona occulta* Gусенлеитнер is from Fars province in the South-central Iran. The main studies have been focused on the areas of South-east, North-central, South and North-central of the country (Blüthgen & Gусенлеитнер, 1970; Giordani Soika, 1970; Gусенлеитнер, 1973b; Ebrahimi & Carpenter, 2008), represented by Sistan-o Baluchestan (26 genera, 42 species), Alborz (23 genera, 42 species), Fars (22, 39) and Tehran (16, 38) provinces, respectively. Some other provinces in the south-west (Khuzestan: 21, 34) and south-east (Kerman: 21, 31) were also explored considerably (Table 2). In his long series of taxonomic papers, Gусенлеитнер (1973b, 1985b, 1986b, 1995a, 1998c, 2000c, 2001, 2006a, 2010a, 2011a, 2012a, 2012b, 2014, 2016, 2018) described 20 new species from 10 provinces, mainly from south and south-eastern areas in Fars, Hormozgan and Kerman provinces, with 6, 4 and 3 new species, respectively. The provincial distribution for many species is unclear, since the locality data are not recorded properly, or local names were used. Some of these species were recovered by the subsequent studies which provide the locality data. Furthermore, there were not sufficient data to assign the (five) species (Kohl & Handlirsch, 1889; Kostylev, 1935; Gусенлеитнер, 1972) recorded from the former Khorasan province, until it was subdivided into three separate provinces in 2004: Khorasan-e Razavi, North Khorasan and South Khorasan.

More than 47 subspecies belonging mainly to the subfamily Eumeninae (43) (Fig. 1) are recorded from various regions of Iran (Blüthgen & Gусенлеитнер, 1970; Giordani Soika, 1970; Gусенлеитнер, 1972; Kurzenko, 1977; Gусенлеитнер, 2000b; 2004; Abbasi et al., 2008; Ebrahimi & Carpenter, 2008; Gусенлеитнер, 2010a; Pashaei Rad et al., 2010; Gусенлеитнер, 2013). Some of these subspecies were later raised to separate species or synonymized (Kostylev, 1937, 1940; Giordani Soika, 1961; Blüthgen & Gусенлеитнер, 1970; Gусенлеитнер, 1972; Ebrahimi & Carpenter, 2008). Further discussion about validity of these subspecies attributed to the knowledge about evolutionary patterns of the vespid lineages (Cumming, 1989; Carpenter, 2003). Additional investigations on both zoogeography and habitat preference, supplemented with molecular data are needed to define the taxonomic situation of these subspecies. A similar situation was also suggested for the sphecid fauna of Iran with many subspecies (Jahantigh et al., 2017). An overall analysis of the vespid fauna of Iran (Table 3) indicates the majority of species (77.1%) are distributed only in the (Eastern and Western) Palaearctic regions. The rest include the species that have a broader area of distribution, expanded into the Afrotropical Region, Oriental Region or both. A higher percentage of the species (11.3%) are found in Afrotropics compared to the Oriental Region (4.8%). There are few Palaearctic species that are found also in the Nearctic (1 species - *Pseudepipona herrichii*) or Neotropical region (1 species - *Eumenes dubius*).

Table 1. Irrelevant records of Vespidae species that should be excluded from the fauna of Iran.

Excluded species	References	General distribution*
<i>Ceramius bicolor</i> (Thunberg, 1815)	Abd-Rabou et al., 2005	Afrotropical
<i>Ceramius capicola</i> Brauns, 1902	Abd-Rabou et al., 2005	Afrotropical
<i>Ceramius fonscolombei</i> Latreille, 1810	Abd-Rabou et al., 2005	Western Palaearctic (Atlanto-Mediterranean)
<i>Ceramius hispanicus</i> Dusmet, 1909	Abd-Rabou et al., 2005	Western Palaearctic (Iberian)
<i>Ceramius tuberculifer</i> de Saussure, 1853	Abd-Rabou et al., 2005	Western Palaearctic: (Atlanto-Mediterranean)
<i>Jugurtia brauni</i> (Schulthess, 1922)	Abd-Rabou et al., 2005	Afrotropical
<i>Masarina familiaris</i> Richards, 1962	Abd-Rabou et al., 2005	Afrotropical
<i>Dolichovespula media</i> (Retzius, 1783)	Abd-Rabou et al., 2005; Sakenin et al., 2010b	Eastern and Western Palaearctic
<i>Dolichovespula saxonica</i> (Fabricius, 1793)	Abd-Rabou et al., 2005; Sakenin et al., 2010b	Eastern and Western Palaearctic
<i>Vespa rufa</i> (Linnaeus, 1758)	Abd-Rabou et al., 2005; Sakenin et al., 2010a, 2010b	Eastern and Western Palaearctic, Oriental

**Figure 1.** A comparision of the number of genera and species of four subfamilies of Vespidae in Iran.




Figure 2. Number of recorded species for the Vespidae genera in Iran. Genera with fewer than two species are excluded.

On the other hand, there are some species (*Polistes olivaceus*, *Polistes dominula*, *Vespa germanica*), that are widely distributed in various biogeographical regions both in the northern and southern Hemisphere. These species expanded their distribution through the association with human global transportation (Liebert et al., 2006; Masciocchi & Corley, 2013; Hammer & Jensen, 2019).

Table 2. Number of recorded genera and species of Vespidae from Iranian provinces.

Locality (Provinces)	Area (km ²)	Eumeninae		Masarinae		Polistinae		Vespinae		Total	
		Gen.*	sp.**	Gen.	sp.	Gen.	sp.	Gen.	sp.	Gen.	sp.
Alborz	5.128	17	33	4	7	1	1	1	1	23	42
Ardabil	179.02	2	2	0	0	1	1	1	1	4	5
Bushehr	22.820	4	4	0	0	2	2	1	1	7	7
Charmahal-o Bakhtiari	16.305	0	0	1	1	0	0	0	0	1	1
East Azerbaijan	45.757	10	17	1	1	1	2	3	3	15	24
Fars	122.779	16	27	3	7	1	3	2	2	22	39
Golestan	20.206	3	5	0	0	1	2	3	4	7	11
Guilan	14.086	4	4	0	0	1	3	3	4	8	11
Hamadan	19.329	3	3	0	0	1	1	3	2	7	6
Hormozgan	70.743	9	13	2	3	1	3	2	2	14	21
Ilam	20.023	1	1	0	0	1	2	1	2	3	5
Isfahan	106.754	2	2	1	1	1	1	2	2	6	6
Kerman	179.469	15	23	2	2	2	4	2	2	21	31
Kermanshah	24.867	2	2	0	0	1	2	2	2	5	6
Khorasan-e Razavi	127.223	6	10	1	1	1	2	2	3	10	16
Khuzestan	63.928	14	25	2	2	2	4	3	3	21	34
Kohgiloyeh-va Boyerahmad	15.498	3	3	0	0	1	1	3	3	7	7
Kordestan	29.048	3	6	1	1	1	3	2	2	7	12
Lorestan	28.199	5	6	0	0	2	2	2	2	9	10
Markazi	29.088	9	15	2	2	1	3	2	2	14	22
Mazandaran	23.725	6	9	0	0	1	4	3	4	10	17
North Khorasan	28.341	2	2	0	0	1	1	2	3	5	6
Qazvin	15.559	4	4	0	0	0	0	2	2	6	6
Qom	11.514	1	1	0	0	1	1	2	2	4	4
Semnan	97.193	5	5	0	0	1	1	2	2	8	8
Sistan-o Baluchestan	178.850	22	35	1	2	1	3	2	2	26	42
South Khorasan	138.845	4	7	0	0	1	2	2	2	7	11
Tehran	13.688	11	28	1	2	1	4	3	4	16	38
West Azarbaijan	37.393	0	0	0	0	0	0	3	3	3	4
Yazd	73.551	1	1	0	0	0	0	2	2	3	3
Zanjan	21.783	9	16	0	0	1	3	2	2	12	21

*Gen. = genus; **Sp. = species

Table 3. Biogeographical distribution of the vespid species in Iran.

Area of distribution	No of species	Frequency (%)
Palaearctic	178	77.1
Palaearctic and Afrotropical	26	11.3
Palaearctic and Oriental	11	4.8
Palaearctic, Afrotropical and Oriental	7	3
Holarctic (Palaearctic and Nearctic)	1	0.4
Palaearctic and Neotropical	1	0.4

While many areas remained unexplored, the known Vespidae of Iran indicate a rich fauna that needs further studies in different habitats. No comparison was done to determine the faunal similarity with the neighbouring countries, but the number of species (Turkey: 298 (species and subspecies) (Yildirim & Guseinleitner, 2012), Russia: 197 (Antropov & Fateryga, 2017), Pakistan: 79 (Rafi et al., 2017)) indicates a moderately well developed knowledge about the vespid fauna of Iran.

Acknowledgements

This work was supported by the Grant No. UOZ-GR-9618-6, University of Zabol to E. Rakhshani. James M. Carpenter was supported by his institute. Our sincere thanks to Volker Mauss (State Museum of Natural History Stuttgart) for editing both early and final version of the manuscript.

Conflict of Interests

The authors declare that there is no conflict of interest regarding the publication of this paper.

References

- Abbasi, R., Rad, S.P., Ebrahimi, E. & Sheidaei, M. (2008) Faunistic study of vespid wasps in Zanjan Province (Northwest of Iran) with some ecological measures. *Environmental Sciences*, 6 (1), 65–74.
- Abd-Rabou, S., Ghahari, H., Mauss, V. & Plant, J. (2005) New records of Apidae, Andrenidae, Sphecidae and Vespidae (Hymenoptera) from Iran. *Egyptian Journal of Agricultural Research*, 83 (4), 1613–1618¹.
- Antropov, A.V. & Fateryga, A.V. (2017) Family Vespidae. In: Lelej, A.S., Proshchalykin, M.Yu. & Loktionov, V.M. (Eds). Annotated Catalogue of the Hymenoptera of Russia. Volume I. Symphyta and Apocrita: Aculeata. *Proceedings of the Zoological Institute RAS*, Suppl. 6, 175–196.
- Archer, M. (1981) Taxonomy of the sylvestris group (Hymenoptera: Vespidae, *Dolichovespula*) with the introduction of a new name and notes on distribution. *Insect Systematics & Evolution*, 12 (2), 187–193.
<https://doi.org/10.1163/187631281794709944>
- Archer, M.E. (1989) *A Key to the World Species of the Vespinae (Hymenoptera)*. Part I. Keys, Checklist and Distribution. The University College of Ripon and York St John, York, 41 pp.
- Białyński-Birula, A. (1926) Ein Beitrag zur Kenntnis der paläarktischen *Hoplomerus*-Arten (Hymenoptera - Diptera). *Bulletin de l'Académie des Sciences de l'URSS*, 20 (10–11), 885–906.
- Birula, A. (1930) Über die russischen Wespen und ihre geographische Verbreitung (Dritter Beitrag). *Annuaire du Musée Zoologique de l'Académie des Sciences de l'URSS*, 31, 291–339.
- Blüthgen, P. (1941) Die Untergattungen *Hoplomerus* und *Monoploemerus* der Gattung *Hoplomerus* (Hym. Vespidae Eumenidinae). *Archiv für Naturgeschichte* (N.F.), 10 (3), 305–344.
- Blüthgen, P. (1956) Untersuchungen über palaarktische Eumenidae: III. Zur Kenntnis der paläarktischen *Pterocheilus*-Arten, IV. Die Untergattungen *Alastor* Lep. und *Parastalor* Blüthg. der Gattung *Alastor* Lep. *Mitteilungen aus dem Zoologischen Museum in Berlin*, 32, 117–149.
- Blüthgen, P. (1961) Ergebnisse der Deutschen Afghanistan-Expedition 1956 der Landessammlungen für Naturkunde Karlsruhe. *Beiträge zur Naturkundlichen Forschung in Südwestdeutschland*, 19, 277–287.

¹ V. Mauss and J. Plant did not publish this article and repudiate to be coauthors of this publication (Mauss in litt. 2019).

- Blüthgen, P. & Guseinleitner, J. (1970) Über Faltenwespen aus dem Iran (Hym., Diptoptera), *Stuttgarter Beiträge Naturkunde*, 223, 1–13.
- Carpenter, J.M. (1996) Distributional checklist of species of the genus *Polistes* (Hymenoptera, Vespidae, Polistinae, Polistini). *American Museum Novitates*, 3188, 1–39.
- Carpenter, J.M. (2001) Checklist of species of the subfamily Masarinae (Hymenoptera: Vespidae). *American Museum Novitates*, 3325, 1–39.
[https://doi.org/10.1206/0003-0082\(2001\)325%3C0001:COSOTS%3E2.0.CO;2](https://doi.org/10.1206/0003-0082(2001)325%3C0001:COSOTS%3E2.0.CO;2)
- Carpenter, J.M. (2003) Return to the subspecies concept in the genus *Zeta* (Hymenoptera: Vespidae; Eumeninae). *Boletín del Museo Nacional de Historia Natural del Paraguay*, 14 (1–2), 19–24.
- Carpenter, J.M. & Kojima, J. (1997) Checklist of the species in the subfamily Vespinae (Insecta: Hymenoptera: Vespidae). *Natural History Bulletin of Ibaraki University*, 1, 51–92.
- Carpenter, J.M., Guseinleitner, J. & Madl, M. (2010) A catalogue of the Eumeninae (Hymenoptera: Vespidae) of the Ethiopian region excluding Malagasy subregion. Part II: genera *Delta* de Saussure 1885 to *Zethus* Fabricius 1804 and species incertae sedis. *Linzer Biologische Beiträge*, 42 (1), 251–253.
- Castro, L. & Dvořák, L. (2009) New and noteworthy records of vespid wasps (Hymenoptera: Vespidae) from the Palaearctic region (II). *Boletín de la Sociedad Entomológica Aragonesa*, 44, 295–304.
- Castro, L. & Dvořák, L. (2010) New and noteworthy records of vespid wasps (Hymenoptera: Vespidae) from the Palaearctic region (III). *Acta Musei Moraviae, Scientiae Biologicae* (Brno), 95 (2), 37–53.
- Cumming, J.M. (1989) Classification and evolution of the eumenine wasp genus *Symmorphus* Wesmael (Hymenoptera: Vespidae). *The Memoirs of the Entomological Society of Canada*, 121, 5–168.
<https://doi.org/10.4039/entm121148fv>
- de Saussure, H. (1854–1856) *Études sur la famille des Vespides III: La monographie des Masariens et un supplément à la monographie des Euméniens*. Genève (Kessmann & CHerbuliez), 352 pp.
- Dover, C. (1925) [1924] Further notes on the Indian Diplopterous wasps. *Journal of the Asiatic Society of Bengal* (new series), 20, 289–305.
- Dover, C. (1929) Wasps and bees in the Raffles Museum, Singapore. *Bullettin of Raffles Museum*, 2, 43–70.
- Dover, C. & Rao, H.S. (1922) A note on the Diplopterous Wasps in the Collection of the Indian Museum. *Journal of Asiatic Society of Bengal*, [N.S.], 18, 235–249.
- du Buysson, R. (1904) [1905] Monographie des Guêpes ou Vespa. *Annales de la Société Entomologique de France*, 73 (3), 485–634.
- du Buysson, R. (1912) Les Vespides, Euménides and Scoliides. *Annales d'Histoire Naturelle Délégation en Perse II. Entomologie*, 1, 77–83.
- Dvořák, L., Ghahari, H., Carpenter, J. & Abbasi, R. (2012) On the distribution and taxonomy of vespine wasps of Iran (Hymenoptera: Vespidae: Vespinae). *Acta Musei Moraviae, Scientiae Biologicae* (Brno), 97 (2), 69–86.
- Ebrahimi, E. & Carpenter, J.M. (2008) Catalog of the vespid wasps of Iran (Hymenoptera, Vespidae). *Zootaxa*, 1785, 1–42.
<https://doi.org/10.11646/zootaxa.1785.1.1>
- Ebrahimi, E. & Carpenter, J.M. (2012) Distribution pattern of the hornets *Vespa orientalis* and *V. crabro* in Iran: (Hymenoptera: Vespidae). *Zoology in the Middle East*, 56 (1), 63–66.
<https://doi.org/10.1080/09397140.2012.10648942>
- Esmaili, M. & Rastegar, R. (1974) Identified species of aculeate Hymenoptera of Iran. *Journal of the Entomological Society of Iran*, 2 (2), 41–52.

- Farahbakhsh, Gh. (1961) *A Checklist of Economically Important Insects and Other Enemies of Plants and Agricultural Products in Iran*. Iranian Ministry of Agriculture, Department of Plant Protection, Tehran, 153 pp.
- Fateryga, A.V. & Mokrousov, M.V. (2019) New records of eumenine wasps (Hymenoptera: Vespidae: Eumeninae) from Russia with description of a new species of *Leptochilus* de Saussure, 1853. *Zootaxa*, 4612 (3), 412–422.
<https://doi.org/10.11646/zootaxa.4612.3.7>
- Fateryga, A.V. & Popov, I.B. (2017) New records of Vespidae and Megachilidae (Hymenoptera) in Russia. *Ekosistemy*, 9, 86–89.
- Fateryga, A.V. (2010) Trophic relations between vespid wasps (Hymenoptera, Vespidae) and flowering plants in the Crimea. *Entomological Review*, 90 (6), 698–705.
<https://doi.org/10.1134/S0013873810060047>
- Fateryga, A.V. (2017) New records of solitary vespid wasps (Hymenoptera: Vespidae: Eumeninae, Masarinae) from Russia and adjacent countries. *Far Eastern Entomologist*, 334, 1–16.
- Fateryga, A.V. (2018) Wasps of the family Vespidae (Hymenoptera) of the Crimean Peninsula. *Entomofauna*, 39 (1), 193–233.
- Fateryga, A.V., Ivanov, S.P. & Budashkin, Y.I. (2017) First data on the bionomics of *Brachyodynerus magnificus magnificus* (Morawitz, 1867) (Hymenoptera: Vespidae: Eumeninae). *Zootaxa*, 4312 (2), 343–354.
<https://doi.org/10.11646/zootaxa.4312.2.9>
- Fateryga, A.V., Proshchalykin, M.Y., Aliyev, K.A. & Maharramov, M.M. (2019) To the knowledge of eumenine wasps (Hymenoptera: Vespidae: Eumeninae) of Nakhchivan Autonomous Republic of Azerbaijan. *Far Eastern Entomologist*, 379, 25–32.
<https://doi.org/10.25221/fee.379.2>
- Gess, S.K. & Gess, F.W. (2010) Pollen wasps and flowers in southern Africa. *SANBI Biodiversity Series*, 18, 1–147.
- Giordani Soika, A. (1943) Eumenini paleartici nuovi o poco noti (Hym.). *Bollettino del Museo Civico di Storia Naturale di Venezia*, 3 (1), 1–14.
- Giordani Soika, A. (1949) Revisione degli *Eumenes* appartenenti al sottogenere *Katamenes* (Hymenoptera: Vespidae). *Bollettino della Società Entomologica Italiana*, 79 (3–4), 40–47.
- Giordani Soika, A. (1952) Sulle caratteristiche biogeografiche della Palestina, Arabia ed Egitto, con un contributo alla conoscenza degli Zethini ed Eumenini della Palestina. *Bollettino del Museo Civico di Storia Naturale di Venezia*, 6 (1), 5–62.
- Giordani Soika, A. (1957) Expedition to South-West Arabia 1937–8. 31- Hymenoptera: Vespidae. *British Museum (Natural History) Expedition to South-West Arabia*, 1 (31), 471–484.
- Giordani Soika, A. (1958) [1957] Biogeografia, Evoluzione e Sistematica dei Vespidi Solitari della Polinesia Meridionale. *Bollettino del Museo Civico di Storia Naturale di Venezia*, 10, 183–221.
- Giordani Soika, A. (1961) Notulae Vespidiologicae XVII–XVIII. *Atti della Società Italiana di Scienze Naturali e del Museo Civico di Storia Naturale in Milano*, 100, 373–388.
- Giordani Soika, A. (1966) Notulae vespidiologicae XXVI. Eumenidi nuovi o poco noti della penisola iberica. *Bollettino del Museo Civico di Storia Naturale di Venezia*, 17 (1964), 93–95.
- Giordani Soika, A. (1968) Le Specie Paleartiche del Genere *Cyrtolabus* V.d.Vecht (Hym. Eumenidae). *Bollettino del Museo Civico di Storia Naturale di Venezia*, 18, 109–124.
- Giordani Soika, A. (1970) Contributo alla conoscenza degli Eumenidi del Medio Oriente. *Bollettino del Museo Civico di Storia Naturale di Venezia*, 41, 27–183.
- Giordani Soika, A. (1973) Descrizione di nuovi Eumenidi. *Bollettino del Museo Civico di Storia Naturale di Venezia*, 24, 97–131.
- Giordani Soika, A. (1979) Notulae vespidiologicae, XL: Eumenidi paleartici nuovi o poco noti. *Bollettino del Museo Civico di Storia Naturale di Venezia*, 30, 247–263.

- Goulet, H. & Huber, J.T. (1993) *Hymenoptera of the World: An Identification Guide to Families of Hymenoptera*. Canada Communication Group – Publishing, Ottawa, 667 pp.
- Guichard, K.M. (1986) [1985] Wasps of the Family Eumenidae (Hymenoptera: Vespoidea) of the Arabian Peninsula. *Fauna of Saudi Arabia*, 7, 202–229.
- Guiglia, D. (1976) [1977] Le vespe sociali (Missione A. Giordani Soika in Iran, 7). *Bollettino del Museo Civico di Storia Naturale di Venezia*, 28, 99–104.
- Guiglia, D. (1976) Contributo alla conoscenza dei Polistini dell'Iran (Hymenoptera, Vespidae). *Bollettino del Museo Civico di Storia Naturale di Venezia*, 81, 195–201.
- Gusenleitner, J. (1972) Übersicht über die derzeit bekannten westpaläarktischen Arten der Gattung *Eumenes* Latreille 1802 (Hym. Vespoidea). *Bulletino del Museo Civico di Storia Naturale di Venezia*, 22 (23), 67–117.
- Gusenleitner, J. (1973a) Bemerkenswertes über Faltenwespen. V. (Diptera, Hymenoptera). *Nachrichtenblatt der Bayerischen Entomologen*, 22, 118–120.
- Gusenleitner, J. (1973b) Über Masaridae aus dem nahen Osten (Vespoidea, Hymenoptera). *Bulletino della Museo Civico di Storia Naturale di Venezia*, 24, 55–69.
- Gusenleitner, J. (1979) Die *Microdynerus*- und *Pseudomicrodynerus*-Arten des Balkans und der Türkei. *Linzer Biologische Beiträge*, 11 (1), 75–94.
- Gusenleitner, J. (1981) Revision der paläarktischen *Stenodynerus*-Arten (Hymenoptera, Eumenidae). *Polskie Pismo Entomologiczne*, 51, 209–305.
- Gusenleitner, J. (1985a) Bemerkenswertes über Faltenwespen VIII (Hymenoptera, Vespoidea). *Nachrichtenblatt der Bayerischen Entomologen*, 34, 105–110.
- Gusenleitner, J. (1985b) Neue *Leptochilus*-Arten aus dem Mittelmeergebiet und dem Iran (Hymenoptera, Eumenidae). *Entomofauna*, 6 (9), 81–101.
- Gusenleitner, J. (1986a) Beitrag zur Kenntnis der paläarktischen *Antepipona*-Arten (Hym., Eumenidae). *Annalen des Naturhistorischen Museums in Wien*, Serie B, 88/89, 565–577.
- Gusenleitner, J. (1986b) Drei neue *Alastor*-Arten aus dem Iran (Hymenoptera, Eumenidae). *Entomofauna*, 7 (2), 29–39.
- Gusenleitner, J. (1987) Über Eumenidae aus Nepal (Hymenoptera Vespoidea). *Linzer Biologische Beiträge*, 19 (1), 255–270.
- Gusenleitner, J. (1988a) Neue und bemerkenswerte Faltenwespen aus der Türkei (Hymenoptera, Vespoidea). *Linzer Biologische Beiträge*, 20 (2), 713–737.
- Gusenleitner, J. (1988b) Über Eumenidae aus Thailand, mit einer Bestimmungstabelle für orientalische Labus-Arten (Hymenoptera, Vespoidea). *Linzer Biologische Beiträge*, 20 (1), 173–198.
- Gusenleitner, J. (1994) Über Eumeniden aus dem Nahen Osten und Arabien sowie Beschreibung einer neuen Subspecies aus Zanzibar. *Linzer Biologische Beiträge*, 26 (1), 307–324.
- Gusenleitner, J. (1992) Bemerkenswerte Funde von Masaridae in Griechenland und im Nahen Osten (Vespoidea, Hymenoptera). *Linzer Biologische Beiträge*, 24 (1), 59–63.
- Gusenleitner, J. (1995a) Bemerkungen über die Gattung *Leptochilus* Saussure 1852 und Beschreibung neuer Arten dieser Gattung (Hymenoptera, Vespoidea, Eumenidae). *Linzer Biologische Beiträge*, 27, 169–181.
- Gusenleitner, J. (1995b) Bestimmungstabellen mittel- und südeuropäischer Eumeniden (Vespoidea, Hymenoptera). Teil 4: Die Gattung *Ancistrocerus* Wesmael 1836 mit einem Nachtrag zum Teil 1: Die Gattung *Leptochilus* Saussure. *Linzer Biologische Beiträge*, 27 (2), 753–775.
- Gusenleitner, J. (1997a) Bestimmungstabellen mittel- und südeuropäischer Eumeniden (Vespoidea, Hymenoptera) Teil 7. Die Gattungen *Microdynerus* Thomson 1874 und *Eumicrodynerus* Gusenleitner, 1972. *Linzer Biologische Beiträge*, 29 (2), 779–797.
- Gusenleitner, J. (1997b) Die Gattungen *Syneuodynerus* Blüthgen 1951 und *Intereuodynerus* gen. nov. im Nahen Osten (Hymenoptera, Eumenidae). *Linzer Biologische Beiträge*, 29 (2), 763–769.

- Gusenleitner, J. (1998a) Bestimmungstabellen mittel- und südeuropäischer Eumeniden (Vespoidea, Hymenoptera), Teil 8. Die Gattungen *Odynerus* Latreille 1802, *Gymnomerus* Blüthgen 1938, *Paragymnomerus* Blüthgen 1938 und *Tropidodynerus* Blüthgen 1939.). *Linzer Biologische Beiträge*, 30 (1), 163–181.
- Gusenleitner, J. (1998b) Bestimmungstabellen mittel- und südeuropäischer Eumeniden (Vespoidea, Hymenoptera), Teil 9. Die Gattung *Pseudepipona* Saussure. *Linzer Biologische Beiträge*, 30 (2), 487–495.
- Gusenleitner, J. (1998c) Über Faltenwespen aus dem asiatischen Raum (Hymenoptera, Eumenidae, Masaridae). *Linzer Biologische Beiträge*, 30, 503–513.
- Gusenleitner, J. (2000a) Bestimmungstabellen mittel- und südeuropäischer Eumeniden (Vespoidea, Hymenoptera), Teil 13. Die Gattung *Stenodynerus* Saussure 1863. *Linzer Biologische Beiträge*, 32 (1), 29–41.
- Gusenleitner, J. (2000b) Bestimmungstabellen mittel- und südeuropäischer Eumeniden (Vespoidea, Hymenoptera) Teil 14. Der Gattungsschlüssel und die bisher in dieser Reihe nicht behandelten Gattungen und Arten. *Linzer Biologische Beiträge*, 32 (1), 43–65.
- Gusenleitner, J. (2000c) Neue Eumenidae aus Syrien, der Türkei und dem Iran (Hymenoptera, Vespoidea). *Linzer Biologische Beiträge*, 32 (2), 927–938.
- Gusenleitner, J. (2001) Über die Gattung *Labochilus* Blüthgen 1939 (Hymenoptera, Vespoidea, Eumenidae). *Linzer Biologische Beiträge*, 33 (2), 677–687.
- Gusenleitner, J. (2002) Über die Arten der Gattung *Masaris* Fabricius 1793 aus Zentralasien (Hymenoptera, Vespoidea, Masaridae). *Linzer Biologische Beiträge*, 34 (1), 321–333.
- Gusenleitner, J. (2004) Ein Beitrag zur Kenntnis und Verbreitung palaearktischer Eumenidae (Hymenoptera: Vespoidea). *Linzer Biologische Beiträge*, 36 (2), 1093–1105.
- Gusenleitner, J. (2005) Über Faltenwespen aus dem Oman Teil 2 (Hymenoptera, Vespidae, Eumenidae, Masaridae). *Linzer Biologische Beiträge*, 37 (2), 1199–1201.
- Gusenleitner, J. (2006a) Eine neue Eumenes-Art aus dem Iran (Hymenoptera: Vespidae, Eumeninae). *Linzer Biologische Beiträge*, 38 (2), 1361–1363.
- Gusenleitner, J. (2006b) Über Aufsammlungen von Faltenwespen in Indien (Hymenoptera: Vespidae). *Linzer Biologische Beiträge*, 38 (1), 677–695.
- Gusenleitner, J. (2006c) Über Eumeninae, aufgesammelt in Pakistan (Hymenoptera: Vespidae). *Linzer Biologische Beiträge*, 38 (2), 1295–1305.
- Gusenleitner, J. (2008) Bestimmungstabellen mittel- und südeuropäischer Eumeniden (Vespoidea, Hymenoptera) 4. Nachtrag. *Linzer Biologische Beiträge*, 40 (2), 1505–1507.
- Gusenleitner, J. (2010a) Bemerkenswertes über Faltenwespen X (Hymenoptera: Vespidae: Eumeninae, Masarinae). *Linzer Biologische Beiträge*, 42 (2), 1347–1352.
- Gusenleitner, J. (2010b) Order Hymenoptera, family Vespidae. *Arthropod fauna of the UAE*, 3, 422–467.
- Gusenleitner, J. (2011a) Bemerkenswertes über Faltenwespen XI (Hymenoptera: Vespidae: Eumeninae: Masarinae). *Linzer Biologische Beiträge*, 43 (1), 323–329.
- Gusenleitner, J. (2011b) Bemerkenswertes über Faltenwespen XII (Hymenoptera: Vespidae: Eumeninae). *Linzer Biologische Beiträge*, 43 (2), 1127–1134.
- Gusenleitner, J. (2011c) Eine Aufsammlung von Faltenwespen aus Laos im Biologiezentrum Linz (Hymenoptera: Vespidae: Vespinae, Stenogastrinae, Polistinae, Eumeninae). *Linzer Biologische Beiträge*, 43 (2), 1351–1368.
- Gusenleitner, J. (2012a) Bemerkenswertes über Faltenwespen XII (Hymenoptera: Vespidae: Eumeninae). *Linzer Biologische Beiträge*, 44 (2), 1129–1136.
- Gusenleitner, J. (2012b) Neue Masarinae aus der paläarktischen Region (Hymenoptera: Vespidae: Masarinae). *Linzer Biologische Beiträge*, 44, 319–326.
- Gusenleitner, J. (2013) Die Gattungen der Eumeninae im Nahen Osten, in Nordafrika und in Arabien (Hymenoptera: Vespidae: Eumeninae). *Linzer Biologische Beiträge*, 45 (1), 5–107.

- Gusenleitner, J. (2014) Eine neue Gattung und Art der Eumeninae aus dem Iran (Hymenoptera: Vespidae, Eumeninae). *Linzer Biologische Beiträge*, 46 (1), 537–540.
- Gusenleitner, J. (2016) Über Eumeninae aus dem Nahen Osten (Hymenoptera: Vespidae: Eumeninae). *Linzer Biologische Beiträge*, 48 (1), 89–97.
- Gusenleitner, J. (2018) Neue asiatische Faltenwespen (Hymenoptera, Vespidae: Eumeninae, Masarinae). *Linzer Biologische Beiträge*, 50 (1), 303–308.
- Gusenleitner, J. (2019) ZOBODAT: Zoological-Botanical Database (Vespoidea) (version 4.0, Oct 2011). In: Roskov Y., Ower G., Orrell T., Nicolson D., Bailly N., Kirk P.M., Bourgoin T., DeWalt R.E., Decock W., Nieukerken E. van, Zarucchi J. & Penev L. (Eds.), *Species 2000 & ITIS Catalogue of Life, 2019 Annual Checklist*. Available from: <https://www.catalogueoflife.org/annual-checklist/2019>. Naturalis, Leiden, the Netherlands.
- Gusenleitner, J. & Madl, M. (2009) Notes on Vespidae (Hymenoptera) of Mauritius. *Entomofauna*, 30 (27), 465–471.
- Gusenleitner, J., Fallahzadeh, M., Haghghi, A. & Dousti, A. (2013) Two new species of Eumeninae from Iran (Hymenoptera: Vespidae). *Linzer Biologische Beiträge*, 45 (1), 109–116.
- Gusenleitner, J. & Ockermüller, E. (2018) Neue Eumeninae aus der Paläarktischen Region (Hymenoptera: Vespidae: Eumeninae). *Linzer Biologische Beiträge*, 50 (2), 1091–1098.
- Hammer, S. & Jensen, J.K. (2019) The invasion of two species of social wasps (Hymenoptera, Vespidae) to the Faroe Islands. *BioInvasions Records*, 8 (3), 558–567.
- Hamzavi, F., Carpenter, J. M. & Ting-Jing, L. (2019) A contribution to the study of Eumeninae (Hymenoptera, Vespidae) in South-Eastern Iran. *Journal of Insect Biodiversity and Systematics*, 5 (4), 393–398.
- Jahantigh, F., Rakhshani, E. Mokhtari, A. & Ramroodi, S. (2017) Catalogue of Ampulicidae, Crabronidae and Sphecidae of Iran (Hymenoptera, Apoidea). *Zootaxa*, 4307 (1), 001–096. <https://doi.org/10.11646/zootaxa.4307.1.1>
- Khoobdel, M., Tavassoli, M., Salari, M. & Firozi, F. (2014) The stinging Apidae and Vespidae (Hymenoptera: Apocrita) in Iranian islands, Qeshm, Abu-Musa, Great Tunb and Lesser Tunb on the Persian Gulf. *Asian Pacific Journal of Tropical Biomedicine*, 4, 258–262. <https://doi.org/10.12980/APJT.B.4.2014C1153>
- Kim, J.K., & Yamane, S. (2004) Revision of the Genus *Stenodynerus* Saussure (Hymenoptera, Vespidae, Eumeninae) in the Far East, with Descriptions of a New Species and a New Subspecies from Taiwan. *Japanese Journal of Systematic Entomology*, 10 (2), 235–264.
- Kohl, F. & Handlirsch, A. (1889) Transcaspische Hymenopteren. *Verhandlungen der kaiserlich-königlichen Zoologisch-Botanischen Gesellschaft in Wien*, 39, 267–286.
- Kojima, J.I. & Carpenter, J.M. (1997) Catalog of species in the polistine tribe Ropalidiini (Hymenoptera, Vespidae). American Museum novitates; no. 3199. *American Museum Novitates*, 3199, 1–96.
- Kokujev, N.R. (1912) Hymenoptera nova e Caucaso, collecta a dom. A.B. Schelkonikov. Mitteilungen des Kaukasischen Museums, 7 (1), 1–6.
- Kostylev, G. (1929) Materialien zur Kenntnis der Vespiden-Fauna Armeniens. *Revue Russe d'Entomologie*, 23 (1–2), 76–82.
- Kostylev, G. (1935) Materialien zur Kenntnis der Masariden-Fauna der Paläarktis. *Archives du Musée Zoologique de l'Université de Moscow*, 2, 85–116.
- Kostylev, G. (1937) [1936] Quelques espèces nouvelles ou peu connues de la famille des Vespidés. *Archives du Musée Zoologique de l'Université de Moscow*, 3, 221–226.
- Kostylev, G. (1939) Nouvelles espèces des genres *Eumenes* Latr. et *Labus* Sauss. du sudouest de l'Asie. *Archives du Musée Zoologique de l'Université de Moscow*, 5, 161–168.

- Kostylev, G. (1940) Espèces nouvelles et peu connues de Vespides, d'Euménides et de Masarides paléarctiques (Hymenoptera). II. *Bulletin de la Société des Naturalistes de Moscou, Section Biologique (NS)*, 49, 137–154.
- Kumar, G.P., Carpenter, J.M., Castro, L. & Sureshan, P.M. (2017) A taxonomic review of the Indian species of the genus *Eumenes* Latreille (Hymenoptera: Vespidae: Eumeninae). *Zootaxa*, 4317 (3), 469–498.
<https://doi.org/10.11646/zootaxa.4317.3.3>
- Kumar, P.G., Castro, L. & Sheikh, A.H. (2019) Additions to the knowledge of the genus *Katamenes* Meade-Waldo, 1910 (Hymenoptera, Vespidae, Eumeninae) in India with a key and a checklist to the region's species. *Boletín de la Asociación Española de Entomología*, 43 (1–2), 15–26.
- Kurzenko, N.V. (1977) Eumenid wasps (Hymenoptera, Eumenidae) of the Mongolian People's Republic and adjacent regions of China and Southern Siberia. *Insects of Mongolia*, 5, 537–582.
- Kurzenko, N.V. (1978) *Solitary Vespid Wasps of the Family Eumenidae (Hymenoptera) of the fauna of the USSR*. Thesis manuscript to acquire a scientific degree of Candidate of Biological Science. Institute of Biology and Soil Science of the Far East Center of the Academy of Sciences of USSR, Vladivostok, 326 pp.
- Liebert, A.E., Gamboa, G.J., Stamp, N.E., Curtis, T.R., Monnet, K.M., Turillazzi, S. & Starks, P.T. (2006) Genetics, behavior and ecology of a paper wasp invasion: *Polistes dominulus* in North America. *Annales Zoologici Fennici* 43, 595–624.
- Mahmood, K., Ullah, M., Aziz, A., Hasan, S.A. & Inayatullah, M. (2012) To the knowledge of Vespidae (Hymenoptera) of Pakistan. *Zootaxa*, 3318, 26–50.
<https://doi.org/10.11646/zootaxa.3318.1.2>
- Masciocchi, M. & Corley, J. (2013) Distribution, dispersal and spread of the invasive social wasp (*Vespa germanica*) in Argentina. *Austral Ecology*, 38 (2), 162–168.
<https://doi.org/10.1111/j.1442-9993.2012.02388.x>
- Mauss, V. & Castro, L. (2000) The geographical and altitudinal distribution of the pollen wasp *Ceramius hispanicus* (Hymenoptera, Vespidae), endemic on the Iberian Peninsula. *Graellsia*, 56, 87–95.
- Mauss, V., Fateryga, A. & Prosi, R. (2016) Taxonomy, distribution and bionomics of *Celonites tauricus* Kostylev, 1935, stat. n. (Hymenoptera, Vespidae, Masarinae). *Journal of Hymenoptera Research*, 48, 33–66.
<https://doi.org/10.3897/JHR.48.6884>
- Meade-Waldo, G. (1910) V. New species of Diptoptera in the collection of the British Museum. *Journal of Natural History*, 5 (25), 30–51.
<https://doi.org/10.1080/00222931008692723>
- Meade-Waldo, G. (1911) X. New species of Diptoptera in the collection of the British Museum. *Journal of Natural History*, 7 (37), 98–113.
<https://doi.org/10.1080/00222931108692910>
- Mokrousov, M.B. & Zryanin, V.A. (2015) Materials on the early spring wasps and ants fauna of Uzbekistan (Hymenoptera: Vespoidea: Chrysididae, Scoliidae, Pompiloidea, Vespoidea, Apoidea [Spheciformes], Formicoidea). *Eversmannia*, 5, 36–48.
- Morawitz, F. (1885) Eumenidarum species novae. *Horae Societatis Entomologicae Rossicae*, 19, 135–181.
- Morawitz, F. (1888) Hymenoptera aculeata nova. *Horae Societatis Entomologicae Rossicae*, 22, 224–302.
- Morawitz, F. (1895) Materialien zu einer Vespidenfauna des Russischen Reiches. *Horae Societatis Entomologicae Rossicae*, 29, 407–493.
- Morice, F. (1921) Annotated lists of aculeate Hymenoptera (except Heterogyna) and chrysids recently collected in Mesopotamia and north-west Persia. *The Journal of the Bombay Natural History Society*, 28, 192–203.

- Nematy, A., Khayrandish, M., Ebrahimi, E. & Carpenter, J.M. (2016) First report of the paper wasp *Polistes rothneyi* Cameron (Hym.: Vespidae: Polistinae) from Iran. *Journal of Entomological Society of Iran*, 36 (3), 237–239. [In Persian with English summary].
- Neumeyer, R. & Castro, L. (2015) New data on the distribution of *Polistes bischoffi* Weyrauch, 1937 and *Polistes helveticus* Neumeyer, 2014, a synonym of *Polistes albellus* Giordani Soika, 1976 n. stat. (Hymenoptera: Vespidae). *Boletín de la Sociedad Entomológica Aragonesa*, 57, 205–2016.
- Nguyen, L.T., Dang, H.T., Kojima, J.I. & Carpenter, J.M. (2014) An annotated distributional checklist of solitary wasps of the subfamily Eumeninae (Hymenoptera: Vespidae) of Vietnam. *Entomologica Americana*, 120 (1), 7–18.
<https://doi.org/10.1664/13-RA-010.1>
- Pashaei Rad, Sh., Abbasi, R. & Ebrahimi, E. (2006) Two new records for the fauna of solitary wasps of the subfamily Eumeninae (Hymenoptera: Vespidae) in Iran. *Zoology in the Middle East*, 39, 117–119.
<https://doi.org/10.1080/09397140.2006.10638195>
- Pashaei Rad, Sh., Abbasi, R., Soleimani, G. & Dvořák, L. (2010) New and supplementary information on the vespid fauna of Iran (Hymenoptera: Vespidae). *Zoology in the Middle East*, 50 (1), 95–100.
<https://doi.org/10.1080/09397140.2010.10638417>
- Perez, J. (1910) Notes sur les Vespides. *Actes de la Société Linnéenne de Bordeaux*, 64, 1–20.
- Pickett, K.M. & Carpenter, J.M. (2010) Simultaneous analysis and the origin of eusociality in the Vespidae (Insecta: Hymenoptera). *Arthropod Systematics & Phylogeny*, 68 (1), 3–33.
- Rafi, M.A., Carpenter, J. M., Qasim, M., Shehzad, A., Zia, A., Khan, M.R., Mastoi, M.I., Naz, F., Ilyas, M. & Shah, M. (2017) The vespid fauna of Pakistan. *Zootaxa*, 4362 (1), 1–28.
<https://doi.org/10.11646/zootaxa.4362.1.1>
- Richards, O.W. (1962) *A Revisional Study of the Masarid Wasps (Hymenoptera, Vespoidea)*. British Museum (Natural History), 294 pp.
- Ross, K.G. & Matthews, R.W. (1991) *The Social Biology of Wasps*. Cornell University Press, Ithaca, NY.
- Sakenin, H., Samin, N. & Bagriacik, N. (2010a) A contribution to the aculeate Hymenoptera (Insecta) from Iran. *Efflattounia*, 10, 15–20.
- Sakenin, H., Havaskary, M., Samin, N., Shirdel, F. & Imani, S. (2010b) Odonata and Vespidae (Hymenoptera) from Arasbaran and Vicinity, Northwestern Iran. *ISSAAS 2009 International congress, Agriculture for Better Living and Global Economy. Nong Nooch Tropical Botanical Garden and Resort, Pattaya, Thailand*, 11–15 January, pp. 184.
- Salehi Fard, M., Modarres Awal, M. & Guseleinleitner, J. (2013) First Record of *Allodynerus dignotus* Morawitz, 1895 (Hym: Vespidae: Eumeninae) from Iran. *Journal of Plant Protection*, 26 (4), 486–488. [In Persian with English summary]
- Saunders, E. (1905) Hymenoptera Aculeata collected in Algeria by the Rev. Alfred Edwin Eaton, M.A., F.E.S. and the Rev. Francis David Morice, M.A., F.E.S. Part II. Diptoptera. *Transactions of the Entomological Society of London*, 399–417.
- Schmid-Egger, C., Achterberg, K.V., Neumeyer, R., Morinière, J. & Schmidt, S (2017) Revision of the West Palaearctic *Polistes* Latreille, with the descriptions of two species—an integrative approach using morphology and DNA barcodes (Hymenoptera, Vespidae). *ZooKeys*, 713, 53–112.
<https://doi.org/10.3897/zookeys.713.11335>
- Selis, M. (2019) A new species of *Masaris* Fabricius from Iran (Hymenoptera: Vespidae: Masarinae). *Zootaxa*, 4567 (2), 395–400.
<https://doi.org/10.11646/zootaxa.4567.2.12>
- Siddiqui, J.A., Bodlah, I., Carpenter, J.M., Naeem, M., Ahmad, M. & Bodlah, M.A. (2015) Vespidae (Hymenoptera) of the Pothwar region of Punjab, Pakistan. *Zootaxa*, 3914 (5), 501–524.
<https://doi.org/10.11646/zootaxa.3914.5.1>
- Tan, J.L., Carpenter, J.M. & van Achterberg, C. (2018) An illustrated key to the genera of Eumeninae from China, with a checklist of species (Hymenoptera, Vespidae). *ZooKeys*, 740, 109–149.
<https://doi.org/10.3897/zookeys.740.22654>

- van der Vecht, J. & Fischer, F.C.J. (1972) Palaearctic Eumenidae. *Hymenopterum Catalogus (Novo Editio)*, 8, 1-197.
- Weyrauch, W. (1939) Zur Systematik der palaearktischen Polistinen auf biologischer Grundlage. *Archiv für Naturgeschichte*, 8 (2), 145-197.
- Wickwar, O.S. (1908) Hymenoptera new to Ceylon, with descriptions of new species. *Spolia Zeylanica*, 5, 115-122.
- Yildirim, E. & Kojima, J. (1999) Distributional checklist of the species of the family Vespidae (Insecta: Hymenoptera; Aculeata) of Turkey. *Natural History Bulletin of Ibaraki University*, 3, 19-50.
- Yildirim, E. & Gusenleitner, J. (2009) The present situation of the Vespidae-fauna (Hymenoptera, Aculeata) of Turkey. *Linzer Biologische Beiträge*, 41 (1), 937-941.
- Yildirim, E. & Gusenleitner, J. (2012) Contribution to the knowledge of the Vespidae (Hymenoptera, Aculeata) of Turkey, with a checklist of the Turkish species. *Turkish Journal of Zoology*, 36 (3), 361-374.
- <https://doi.org/10.3906/zoo-1012-62>

فهرست به روز زنبورهای خانواده Vespidae (Hymenoptera) در ایران

زهرا رحمانی^۱، احسان رخشانی^{۲*} و جیمز مایکل کارپنتر^۳

۱ گروه گیاه‌پژوهشی، دانشکده کشاورزی، دانشگاه زابل، زابل، ایران.

۲ موزه تاریخ طبیعی، بخش جانورشناسی بی‌مهرگان، نیویورک، آمریکا.

* پست الکترونیکی نویسنده مسئول مکاتبه: rakhshani@uoz.ac.ir

تاریخ دریافت: ۱۱ دی ۱۳۹۸ ، تاریخ پذیرش: ۲۷ دی ۱۳۹۸ ، تاریخ انتشار: ۱۱ بهمن ۱۳۹۸

چکیده: ۲۳۱ گونه از بال‌غشاییان خانواده Vespoidea (Hymenoptera, Vespidae) ایران متعلق به ۵۵ جنس و چهار زیرخانواده شامل Eumeninae (۴۵ جنس، ۱۸۴ گونه)، Masarinae (۵ جنس، ۲۴ گونه)، Polistinae (۲ جنس، ۱۷ گونه) و Vespinae (۳ جنس، ۶ گونه) فهرست شدند. ارزیابی کلی از الگوی پراکنش گونه‌های این خانواده در ایران، نشان دهنده یک فون مرکب متأثر از مناطق بیogeografیایی مختلف است. ۱۱۱ گونه در هر دو منطقه‌ی شرق و غرب پالئارکتیک یافت شده‌اند، در حالی که ۶۷ گونه تنها در منطقه‌ی شرق پالئارکتیک انتشار دارند. تعداد اندکی از گونه‌ها (۱۴ گونه، ۶/۱٪) متعلق به جنس‌های مختلف، در نواحی مرکزی و غربی آسیا پراکنش داشته و گزارشی از حضور آنها در اروپا (غرب پالئارکتیک) و خاور دور وجود ندارد. گونه‌هایی که در هر دو منطقه اورینتال و آفریقاییکال یافت شده‌اند، به ترتیب ۱۱/۷ و ۱۵/۶ درصد از فون زنبورهای وسپیده ایران را شامل می‌شوند. گونه‌های متعددی (۴۸٪، ۲۰/۸٪) منحصرًا از ایران گزارش شده و هنوز هیچ گزارشی از پراکنش این گونه‌ها در سایر کشورها ثبت نشده است. بیشترین تعداد گونه‌ها به ترتیب از استان‌های سیستان و بلوچستان (۴۲ گونه، ۱۸/۲٪)، البرز (۴۲ گونه، ۱۸/۲٪)، فارس (۳۹ گونه، ۱۶/۹٪) و تهران (۳۸ گونه، ۱۶/۵٪) به عنوان مناطق شاخص از جنوب شرق و بخش‌های مرکزی شمال و جنوب کشور گزارش شده‌اند.

واژگان کلیدی: کاتالوگ، پراکنش، زنبورهای سرخ، زنبورهای کوزه‌گر، زنبورهای کاغذساز، زنبورهای گردخوار، زنبورهای زرد

Authors' Brief Biographical Notes



Zahra Rahmani (Ph.D. Candidate)

Department of Plant Protection, College of Agriculture, University of Zabol, P.O. Box 98615-538, I.R. Iran;

E-mail: zahrarahmani@uoz.ac.ir

Senior author is a Ph.D. candidate who has started her project on Taxonomy of Vespidae, then focused on a revisional works on parasitic hymenoptera of the family Pteromalidae (Chalcidoidea). She published many papers from her Ph.D. thesis, to be continued until finalization of the project.

https://www.researchgate.net/profile/Zahra_Rahmani16



Ehsan Rakhshani (Ph.D.)

Department of Plant Protection, College of Agriculture, University of Zabol, P.O. Box 98615-538, I.R. Iran;

E-mail: rakhshani@uoz.ac.ir

Iranian entomologist, who works on Taxonomy of Hymenoptera, especially on parasitic wasps of the family Braconidae and recently on Aculeata in Eastern parts of Iran. He is the author and co-author of more than 200 journal papers, jointly published with researchers from about 30 countries.

https://www.researchgate.net/profile/Ehsan_Rakhshani



James Michael Carpenter (Ph.D.)

Division of Invertebrate Zoology, American Museum of Natural History, Central Park West at 79th Street, New York, NY 10024, USA;

E-mail: carpente@amnh.org

He is the Peter J. Solomon Family Curator, Chair of the Division of Invertebrate Zoology and Professor in the Richard Gilder Graduate School of the American Museum of Natural History. He is the author of more than 250 scientific publications, and has been publishing on Vespidae for four decades including many joint papers with Iranian researchers.

https://www.researchgate.net/profile/James_Carpenter3