



New data on the digger wasps fauna of families Pemphredonidae, Psenidae and Philanthidae (Hymenoptera: Apoidea) in Iran

Shohreh Rezaei¹, Majid Fallahzadeh^{1*}, Abu Fazel Dousti¹ & Nazila Saghaei²

¹ Department of Entomology, Jahrom Branch, Islamic Azad University, Jahrom, Iran.

² Department of Entomology, Marvdasht Branch, Islamic Azad University, Marvdasht, Iran.

ABSTRACT. The digger wasps fauna (Hymenoptera: Apoidea) of Fars province in southern Iran was partly investigated. Members of three families: Pemphredonidae (5 species of 3 genera), Psenidae (3 species of 2 genera) and Philanthidae (2 species of 2 genera) were collected and documented. The following eight species: *Diodontus brevilabris* de Beaumont, 1967, *D. crassicornis* Gribodo, 1894, *D. insidiosus* Spooner, 1938, *Passaloecus corniger* Shuckard, 1837, *Pemphredon inornata* Say, 1824, *Mimesa punctipleuris* (Gussakovskij, 1937), *Psenulus laevigatus* (Schenck, 1857) and *Cerceris spinifera haladai* K. Schmidt, 2000, are new to the Iranian wasp fauna. In addition, two species: *Psenulus schencki* (Tournier, 1889) and *Philanthus variegatus* Spinola, 1839, are new records for Fars province. For each species, we include available distributional data as well as comments on its taxonomy and geographical distribution.

Received:

02 April, 2020

Accepted:

23 April, 2020

Published:

12 May, 2020

Subject Editor:

Ali Asghar Talebi

Key words: Hymenoptera, fauna, new records, Iran

Citation: Rezaei, Sh., Fallahzadeh, M., Dousti, A.F. & Saghaei, N. (2020) New data on the digger wasps fauna of families Pemphredonidae, Psenidae and Philanthidae (Hymenoptera: Apoidea) in Iran. *Journal of Insect Biodiversity and Systematics*, 6 (2), 195–203.

Introduction

Apid wasps (Apoidea excluding Apidae) are the largest group of the aculeate wasps (Hymenoptera: Aculeata), comprising over 10,000 described species worldwide (Aguiar et al., 2013; Pulawski, 2020), which were usually assigned to four families Ampulicidae, Crabronidae, Heterogynaidae and Sphecidae *sensu stricto* or placed all in one family Sphecidae *sensu lato* (for history of classification and details see Pulawski, 2020). Brothers (1999) and Melo (1999) re-classified Apoidea and found that Sphecidae *sensu lato* are paraphyletic with relation to apoid bees. Therefore, Apoidea were splitted into five monophyletic families: Apidae, Ampulicidae, Crabronidae, Heterogynaidae and Sphecidae. Based on recent molecular and phylogenetic studies, Debevec et al. (2012) demonstrated that two families Ampulicidae and Heterogynaidae are sequential sister taxa to Sphecidae and Crabronidae. Later, Branstetter et al. (2017) suggested that Heterogynaidae and paraphyletic Sphecidae *sensu stricto* placed in a polyphyletic Crabronidae. Most recently,

Corresponding author: Majid Fallahzadeh, E-mail: mfallahm@yahoo.com

Copyright © 2020, Rezaei et al. This is an open access article distributed under the terms of the Creative Commons Attribution License (CC BY NC 4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

[Sann et al. \(2018\)](#) established the polyphyletic nature of the digger wasp family "Crabonidae", and they suggested splitting it into eight families: Ammoplanidae, Astatidae, Bembecidae, Crabronidae, Mellinidae, Pemphredonidae, Philanthidae, and Psenidae.

During the last two decades, numerous papers have been published on the fauna of Iranian digger wasps, the representatives are: [Ebrahimi, 2000a, 2000b, 2005, 2008, 2014](#); [Dollfuss, 2006, 2008](#); [Schmid-Egger, 2004, 2019](#); [Fallahzadeh et al., 2006, 2009, 2018](#); [Pulawski, 2007](#); [Ghazi-Soltani et al., 2009, 2010a, 2010b, 2010c](#); [Atbaei et al., 2015](#); [Rezaei & Fallahzadeh, 2015](#); [Schmid-Egger et al., 2016](#); [Sadeghi et al., 2016, 2018a, 2018b, 2019](#); [Jahantigh et al., 2017](#); [Khosroabadi et al., 2019](#)). The main purpose of the present study was to increase our knowledge about the fauna of Iranian Pemphredonidae, Psenidae and Philanthidae in the Fars province.

Material and methods

Specimens for the current study were collected using Malaise traps at seven locations with four elevation levels in the Fars province, Iran: about 2000 (Dasht-e Arzhan and Dalin), about 1500 (Shiraz, Kavar and Zafar Abad), about 700 (Larestan: Karmostaj), and about 500 (Larestan: Hormood Abbasi) meters above the sea level, respectively. Sampling localities are briefly described in Table 1. Voucher specimens are deposited at the Insect Collection of Jahrom Branch, Islamic Azad University, Iran (JIAU) and the Institute of Biodiversity and Ecosystem Research, Sofia, Bulgaria (IBER). Family-level classification is based on the most recent classification of Apoidea suggested by [Sann et al. \(2018\)](#). Nomenclature and general distributional for each species were adapted from [Antropov et al. \(2017\)](#) and [Pulawski \(2020\)](#). Data under Material examined for the recorded species are presented in the following order: number of specimens examined, locality, and date of collection. The GPS coordinates, altitudes of sampling localities, and collector names are given in [Table 1](#). The genera and species are listed alphabetically under each family.

Table 1. List of the sampling localities in Fars province, Iran.

No.	Locality	Coordinate	Altitude in m a.s.l.	Collector
01	Dasht-e Arzhan	29°39'39.0" N 51°59'03.4" E	2152	Sh. Rezaei
02	Dalin	30°02'15.0" N 52°07'54.7" E	2019	Sh. Rezaei
03	Shiraz (Bustan-e Jannat)	29°36'52.0" N 52°28'09.0" E	1584	Sh. Rezaei
04	Kavar	29°11'52.0" N 52°42'06.0" E	1547	Sh. Rezaei
05	Zafar Abad	29°24'06.0" N 52°35'01.0" E	1514	Sh. Rezaei
06	Larestan (Karmostaj)	27°31'55.0" N 54°26'01.0" E	781	Sh. Rezaei
07	Larestan (Hormood Abbasi)	27°32'06.0" N 54°59'02.0" E	506	Sh. Rezaei

Results

Digger wasps collected in the current study represent three families, 7 genera and 10 species as follows:

Family: Pemphredonidae

Genus: *Diodontus* Curtis, 1834

***Diodontus brevilabris* de Beaumont, 1967**

Material examined: Iran, Fars province: Dalin (loc. 02, [Table 1](#)), 19-25.vi.2019, 1♀.

Distribution: Caucasus, Europe, Iran (new record), Kyrgyzstan, Russia, Turkey, Turkmenistan.

Remarks: Because of morphological similarity (see [Budrys, 1996](#)) and the lack of regional keys, this group is considered to be difficult for identification. Recently, [Olszewski et al. \(2016\)](#) keyed Central and Eastern European species while [Budrys et al. \(2019\)](#) keyed the species of *D. minutus* (Fabricius, 1793) species-group of the Mediterranean region. The later key can be used partially to identify fauna from southern Iran. *Diodontus brevilabris* was originally described from Turkey ([de Beaumont, 1967](#)).

***Diodontus crassicornis* Gribodo, 1894**

Material examined: Iran, Fars province: Dasht-e Arzhan (loc. 01, [Table 1](#)), 7-20.vi.2019, 1♂; Kavar (loc. 04, [Table 1](#)), 16-26.vii.2019, 1♂.

Distribution: Algeria, Egypt, Iran (new record), Spain, Tunisia.

Remarks: Iran represents the northeastern limit of this species distribution.

***Diodontus insidiosus* Spooner, 1938**

Material examined: Iran, Fars province: Larestan (loc. 06, [Table 1](#)), 21-30.iv.2018, 1♂.

Distribution: Europe, Iran (new record), Kazakhstan, Russia.

Genus: *Passaloecus* Shuckard, 1837

***Passaloecus corniger* Shuckard, 1837**

Material examined: Iran, Fars province: Shiraz (loc. 03, [Table 1](#)), 24-30.iv.2018, 1♀; 1-9.vi.2019, 1♀; 1-10.ix.2019, 1♀; 22-30.ix.2019, 1♀.

Distribution: Europe, Iran (new record), Japan, Kazakhstan, North Africa, Russia, Turkey.

Genus: *Pemphredon* Latreille, 1796

***Pemphredon inornata* Say, 1824**

Material examined: Iran, Fars province: Shiraz (loc. 03, [Table 1](#)), 1-7.v.2018, 1♀; 17-23.vii.2018, 3♀♀; 21-27.viii.2018, 1♀; Dasht-e Arzhan (loc. 01, [Table 1](#)), 22-31.v.2019, 3♀♀.

Distribution: Central Asia, Europe, China, Iran (new record), Japan, Korean Peninsula, Mongolia, North Africa, North America, Russia, Turkey.

Family: Psenidae

Genus *Mimesa* Shuckard, 1837

***Mimesa punctipleuris* (Gussakovskij, 1937)**

Material examined: Iran, Fars province: Larestan (loc. 07, [Table 1](#)), 29.v.2015-29.6.2015, 1♂.

Distribution: China, Iran (new record), Kazakhstan, Kyrgyzstan, Mongolia, Russia.

Remarks: No review study on Iranian *Mimesa* species is available, therefore the study of this fauna is very difficult. Identification is based on [de Beaumont \(1937\)](#) and [Budrys \(1985\)](#).

Genus: *Psenulus* Kohl, 1896

Psenulus laevigatus (Schenck, 1857)

Material examined: Iran, Fars province: Shiraz (loc. 03, [Table 1](#)), 10-20.iv.2018, 1♀; 1-9.vi.2019, 1♀; Zafar Abad (loc. 05, [Table 1](#)), 18-25.vi.2019, 2♀♀.

Distribution: Europe, Iran (new record), Japan, Korean Peninsula, Russia, Turkey.

Psenulus schencki (Tournier, 1889)

Material examined: Iran, Fars province: Shiraz (loc. 03, [Table 1](#)), 24-30.iv.2018, 1♂; 17-23.vii.2018, 1♂; 21-27.viii.2018, 1♂.

Distribution: Iran: Khorasan-e Razavi ([Samin et al., 2018](#)), Fars (new record), Europe, Georgia, North Africa, Turkey.

Family: Philanthidae

Genus: *Cerceris* Latreille, 1802

Cerceris spinifera haladai K. Schmidt, 2000

Material examined: Iran, Fars province: Zafar Abad (loc. 05, [Table 1](#)), 11-21.v.2019, 1♀; 12-20.vii.2019, 1♀.

Distribution: Iran (new record), Turkey.

Genus: *Philanthus* Fabricius, 1790

Philanthus variegatus Spinola, 1839

Material examined: Iran, Fars province: Larestan (loc. 06, [Table 1](#)), 21-30.iv.2019, 1♂.

Distribution: Iran: Qazvin ([de Beaumont, 1957](#)), Ardabil ([de Beaumont, 1957; Ebrahimi, 2014](#)), East Azerbaijan ([Sakenin et al., 2011](#)), Kerman, Mazandaran ([Ebrahimi, 2014](#)), Fars (new record), Afghanistan, Armenia, Azerbaijan, Central Asia, China, Iraq, Israel, Kazakhstan, Kuwait, North Africa, Oman, Pakistan, Russia, Saudi Arabia, Syria, Turkey, United Arab Emirates.

Discussion

The digger wasps (former family Crabronidae) in Fars and Hormozgan provinces in southern Iran have received increasing attention in recent years ([Fallahzadeh et al., 2006, 2009, 2018; Atbaei et al., 2015; Rezaei & Fallahzadeh, 2015; Schmid-Egger et al., 2016; Sadeghi et al., 2016, 2018a, 2018b, 2019; Khosroabadi et al., 2019](#)). Before 2015, the number of known species from the Fars province was 28 (see [Atbaei et al., 2015](#)). Since then, the number of known species of these wasps from the Fars province increased significantly to 158 species. In the current study, eight species: *Diodontus brevilabris*, *D. crassicornis*, *D. insidiosus*, *Passaloecus corniger*, *Pemphredon inornata*, *Mimesa punctipleuris*, *Psenulus laevigatus* and *Cerceris spinifera haladai*, were added to the Iranian wasp fauna. In addition, two species: *Psenulus schencki* and *Philanthus variegatus*, were new records for the Fars province.

The genus *Diodontus* includes 79 species on all major continents while the Palaearctic fauna has more than half of the known species ([Pulawski, 2020](#)). Only five species of the genus *Diodontus* had been recorded from Iran prior to the present study: *D. hyalipennis*

Kohl, 1892, *D. minutus* (Fabricius, 1793), *D. spinicollis* Gussakovskij, 1933, *D. temporalis* Kohl, 1901 and *D. tristis* (Vander Linden, 1829) ([Gussakovskij, 1933](#); [de Beaumont, 1957](#); [Ebrahimi, 2005, 2014](#); [Atbaei et al., 2015](#); [Rezaei & Fallahzadeh, 2015](#)). With addition of the three species reported in the present study, the number of *Diodontus* known from Iran increases to eight.

Passaloecus, a small genus distributed worldwide, contains about 40 described species. The genus appears to be especially well represented in the Palaearctic region from where more than half of the known species have been recorded ([Pulawski, 2020](#)). So far, the *Passaloecus* species recorded from Iran are *Passaloecus gracilis* (Curtis, 1834), *P. pictus* Ribaut, 1952 and *P. turionum* Dahlbom, 1844 ([de Beaumont, 1957](#); [Ebrahimi, 2005, 2014](#); [Atbaei et al., 2015](#); [Rezaei & Fallahzadeh, 2015](#)), and now we added *P. corniger* to the fauna of Iran.

[Dollfuss \(1995\)](#) revised and keyed out the world species of *Pemphredon*. It is a Holarctic genus with more than 45 described species of which about 28 occur in the Palaearctic region ([Pulawski, 2020](#)). Up to now, only *P. austriaca* Kohl, 1888, *P. lethifer* (Shuckard, 1837), *P. rugifer* (Dahlbom, 1844) and *P. morio* Vander Linden, 1829 have been reported from Iran ([Gussakovskij, 1933](#); [Dollfuss, 1995](#); [Ebrahimi, 2005, 2014](#); [Atbaei et al., 2015](#); [Rezaei & Fallahzadeh 2015](#); [Samin et al., 2018](#)), and now we added *P. inornata* to the fauna of Iran.

Mimesa is an Old World genus with 76 described species, of which more than 40 are found in the Palaearctic region ([Pulawski, 2020](#)). *Mimesa punctipleuris* was originally described from Mongolia ([Gussakovskij, 1937](#)). [Ma et al. \(2008\)](#) reviewed the Chinese species, while [Schmid-Egger \(2014\)](#) revised and keyed out the species from North Africa and the Arabian Peninsula. Currently, six species: *M. caucasica* Maidl, 1914, *M. crassipes* A. Costa, 1871, *M. grandii* Maidl, 1933, *M. jacobsoni* ([Gussakovskij, 1937](#)), *M. punctipleuris* and *M. scheuchli*, are known from Iran (for previous records see [Jahantigh et al., 2017](#)).

Psenulus, a cosmopolitan genus, currently consists of 161 species, most of which are found in the Oriental and Palaearctic regions ([Pulawski, 2020](#)). *Psenulus schencki* was hitherto recorded from northeastern Iran. Therefore, its presence in southern Iran extends the known distribution of this species in the Iranian plateau.

Cerceris is a worldwide, large genus containing about 867 species, of which about 270 have been reported from the Palaearctic region ([Pulawski, 2020](#)). Recently, [Sadeghi et al. \(2019\)](#) reviewed all of *Cerceris* of Iran and listed 53 species and subspecies. *Cerceris spinifera haladai* was originally described from Turkey and has not been recorded thereafter.

Philanthus is a cosmopolitan genus that currently includes 134 nominal species ([Pulawski, 2020](#)). The present record of *Philanthus variegatus* from southern Iran reveals the wide distribution of this species on the Iranian plateau.

At present, considering the results of this work, the number of known species from the Fars province is about 168. Indeed, the 168 included species are only a fraction of what must occur in the Fars province. On the other hand, comprehensive research has not been conducted elsewhere in the country and many provinces are poorly known and not well investigated. The species reported from Fars are several times as much as from any other part of Iran, and a simple comparison shows that other areas need further investigation (see [Jahantigh et al., 2017](#)). Therefore, further researches on the diversity, biology and behavior of digger wasps in other areas are strongly recommended.

Acknowledgments

The research was supported by Islamic Azad University, Jahrom Branch, Jahrom, Iran. We are indebted to Dr. Toshko Ljubomirov (Institute of Biodiversity and Ecosystem Research,

Sofia, Bulgaria) for help in identification of the specimens. We are most grateful to Dr. Toshko Ljubomirov, Dr. Wojciech J. Pulawski (California Academy of Science, USA), Dr. Christian Schmid-Egger (Germany) and two anonymous reviewers for their careful reading and useful correction of the initial manuscript that improved it considerably.

Conflict of Interests

The authors declare that there is no conflict of interest regarding the publication of this paper.

References

- Antropov, A.V., Astafurova, Yu.V., Belokobylskij, S.A., Byvaltsev, A.M., Danilov, Yu.N., Dubovikoffm D.A., Fadeev K. I., Fateryga, A.V., Kurzenko, N.V., Lelej, A.S., Levchenko, T.V., Loktionov, V.M., Mokrousov, M.V., Nemkov, P.G., Proshchalykin, M.Yu., Rosa, P., Sidorov, D.A., Sundukov, Yu.N., Yusupov, Z.M. & Zaytseva, L.A. (2017) *Annotated Catalogue of the Hymenoptera of Russia. Volume I. Symphyta and Apocrita: Aculeata*. Zoological Institute RAS, Saint Petersburg. Proceedings of the Zoological Institute RAS, Supplement 6. 475 pp.
- Atbaei, M., Fallahzadeh, M. & Ljubomirov, T. (2015) A contribution to the fauna of Crabronidae (Hymenoptera, Apoidea) in South-Western Iran. *Journal of Insect Biodiversity*, 3 (11), 1–30.
<https://doi.org/10.12976/jib/2015.3.11>
- Aguiar, A.P., Deans, A.R., Engel, M.S., Forshage, M., Huber, J.T., Jennings, J.T., Johnson, N.F., Lelej, A.S., Longino, J.F., Lohrmann, V., Miko, I., Ohl, M., Rasmussen, C., Taeger, A. & Yu, D.S.K. (2013) Order Hymenoptera Linnaeus, 1758. In: Zhang, Z.Q. (ed.) *Animal Biodiversity: An Outline of Higher-Level Classification and Survey of Taxonomic Richness*. Zootaxa, 3703 (1), 51–62.
<http://dx.doi.org/10.11646/zootaxa.3703.1.12>
- Branstetter, M.G., Danforth, B.N., Pitts, J.P., Faircloth, B.C., Ward, P.S., Buffington, M.L., Gates, M.W., Kula, R.R. & Brady, S.G. (2017) Phylogenomic insights into the evolution of stinging wasps and the origins of ants and bees. *Current Biology*, 27, 1019–1025.
<https://doi.org/10.1016/j.cub.2017.03.027>
- Brothers, D.J. (1999) Phylogeny and evolution of wasps, ants and bees (Hymenoptera, Chrysidoidea, Vespoidea and Apoidea). *Zoologica Scripta*, 28, 233–249.
<https://doi.org/10.1046/j.1463-6409.1999.00003.x>
- Budrys, E. (1985) New and little known species of the genus *Mimesa* Shuck. (Hymenoptera, Sphecidae) of Middle Asia and Kazakhstan and a key to species of the USSR fauna]. *Trudy Zoologicheskogo Instituta Akademii Nauk SSSR* 132:58–76.
- Budrys, E. (1996) Morphometric similarity and summary of measurements of Palearctic species of the genus *Diodontus* Curtis (Hymenoptera, Sphecidae). In: Skirkevičius, A. (ed.), *Lietuvos Entomologų Darbai*. Lietuvos Entomologų Draugija, Ekologijos Institutas, Vilnius, pp. 35–47.
- Budrys, E., Budrienė, A., Orlovskytė, S. & Soon, V. (2019) Two new species of *Diodontus* (Hymenoptera: Pemphredonidae) from the western Mediterranean and their phylogenetic relationships. *The Canadian Entomologist*, 151 (5), 558–583. <https://doi.org/10.4039/tce.2019.46>
- de Beaumont, J. (1937) Les Psenini (Hym. Sphecid.) de la région paléarctique. *Mitteilungen der Schweizerischen Entomologischen Gesellschaft*, 17, 33–93.
- de Beaumont, J. (1957) Sphecidae du nord de l'Iran (Hym.). *Mitteilungen der Schweizerischen Entomologischen Gesellschaft*, 30, 127–139.
- de Beaumont, J. (1967) Hymenoptera from Turkey. Sphecidae, I. With Appendix. *Sphex* Linné, Subgenus *Palmodes* Kohl par P. Roth. *Bulletin of the British Museum (Natural History). Entomology*, 19, 253–382.
- Debevec, A.H., Cardinal, S. & Danforth, B.N. (2012) Identifying the sister group to the bees: a molecular phylogeny of Aculeata with an emphasis on the superfamily Apoidea. *Zoologica Scripta*, 41 (5), 527–535. <https://doi.org/10.1111/j.1463-6409.2012.00549.x>

- Dollfuss, H. (1995) A worldwide revision of *Pemphredon* Latreille 1796 (Hymenoptera, Sphecidae). *Linzer Biologische Beiträge*, 27, 905–1019.
- Dollfuss, H. (2006) The Crabroninae Wasps of "Biologiezentrum Linz" Collection in Linz, Austria (Hymenoptera, Apoidea, Crabronidae), Part 2. *Linzer Biologische Beiträge*, 38 (1), 505–532.
- Dollfuss, H. (2008) The Crabronid wasps of the genera *Oxybelus* Latreille 1796 and *Brimocelus* Arnold 1927 of "Biologiezentrum Linz" Collection in Linz, Austria (Hymenoptera, Apoidea, Crabronidae). *Linzer Biologische Beiträge*, 40, 463–505.
- Ebrahimi, E. (2000a) The first report on three orthopteran predator wasps in Iran. *Proceedings of the 14th Iranian Plant Protection Congress, Isfahan, 5-8 September, Isfahan*, Isfahan University of Technology, Iran, Volume I: Pests, p. 361.
- Ebrahimi, E. (2000b) The first record of three predator wasps in Iran. *Proceedings of the 14th Iranian Plant Protection Congress, Isfahan, 5-8 September, Isfahan*, Isfahan University of Technology, Iran, Volume I: Pests, p. 362.
- Ebrahimi, E. (2005) An identification guide to the Sphecidae of Iran (Insecta, Hymenoptera). *Journal of Entomological Society of Iran*, 24 (2), 109–135.
- Ebrahimi, E. (2008) A contribution to the sphecid wasps of Iran (Hymenoptera: Sphecidae), including first record of six species. *Journal of Entomological Society of Iran*, 28 (1), 93–97.
- Ebrahimi, E. (2014) The list of Hymenoptera in the Hayk Mirzayans insect museum, suborder Apocrita. superfamily Apoidea (Spheciformis series). Families Ampulicidae, Sphecidae, Crabronidae. In: Askari, H., Farazmand, H., Hosseini Nezhad, A., Manzari, S., Mirabolfathi, M., Mofidi-Neyestanak, M., Morovati, M., Zand, E. & Zare, R. (eds.), *Insects of Iran*. Publication No. 20. Iranian Research Institute of Plant Protection, Tehran, pp. 1–62.
- Fallahzadeh, M., Ostovan, H. & Saghaei, N. (2009) A contribution to the fauna of Sphecidae and Crabronidae (Hymenoptera) in Fars province, Iran. *Plant Protection Journal*, 1, 234–248.
- Fallahzadeh, M., Ostovan, H. & Shojaei, M. (2006) First record of four sphecid wasps from Iran. *Applied Entomology and Phytopathology*, 73, 43–44.
- Fallahzadeh, M., Ljubomirov, T., Tavakoli Roodi, T. & Saghaei, N. (2018) A further contribution to the Crabronidae fauna (Hymenoptera, Apoidea, Spheciformes) of southern Iran. *Transactions of the American Entomological Society*, 144 (3), 144, 625–636. <https://doi.org/10.3157/061.144.0312>
- Ghazi-Soltani, G., Ebrahimi, E. & Iranipur, S. (2009) Sphecidae (Sphecinae and Crabroninae) wasp fauna of East Azarbaijan province, Iran. *Journal of Insect Science*, 19, 271–282.
- Ghazi-Soltani, G., Iranipur, S. & Ebrahimi, E. (2010a) The wasps of family Sphecidae (Hymenoptera), subfamilies Larrinae, Astatinae, Pemphredoninae, Nyssoninae and Philanthinae in East Azarbaijan province. *Iranian Journal of Plant Protection Science*, 41, 179–186.
- Ghazi-Soltani, G., Khaghaninia, S. & Shahim, K. (2010b) An introduction to sphecid wasps of Horand forest-Iran. *Munis Entomology and Zoology*, 5, 636–641.
- Ghazi-Soltani, G., Ebrahimi, E., Iranipur, S. & Pour Abad, R. (2010c) Sphecid wasps from East Azerbaijan province, Iran (Hymenoptera: Sphecidae). *Munis Entomology and Zoology*, 5, 796–803.
- Gussakovskij, V.V. (1933) Sphecidae et Psammocharidae (Hymenoptera), a cl. N. Zarudnyi in Persia orientali collectae. *Trudy Zoologicheskogo Instituta Akademii Nauk SSSR*, 1, 269–307.
- Gussakovskij, V.V. (1937) *Didineis* Wesm., *Pison* Latr. и *Psen* Latr. (Hymenoptera Sphecoidea) [= Obzor palearkticheskikh vidov rodov *Didineis* Wesm., *Pison* Latr. и *Psen* Latr. (Hymenoptera Sphecoidea)] – Espèces paléarctiques des genres *Didineis* Wesm., *Pison* Latr. et *Psen* Latr. (Hymenoptera Sphecoidea). *Trudy Zoologicheskogo Instituta Akademii Nauk SSSR* (= *Travaux de l'Institut Zoologique de l'Académie des Sciences de l'URSS*) 4, 599–698.
- Jahantigh, F., Rakshani, E., Mokhtari, A. & Ramroodi S. (2017) Catalogue of Ampulicidae, Crabronidae and Sphecidae of Iran (Hymenoptera, Apoidea). *Zootaxa*, 4307 (1), 1–96. <https://doi.org/10.11646/zootaxa.4307.1.1>
- Khosroabadi, M., Fallahzadeh, M., Schmid-Egger, C. & Dousti, A.F. (2019) Two new records of Crabronidae (Hymenoptera: Apoidea) from Iran. *Journal of Entomological Society of Iran*, 38(4), 439–441. <https://doi.org/10.22117/jesi.2018.121805.1222>

- Ma, L., Li, Q. & Chen, X.X. (2008). The genus *Mimesa* in China with descriptions of two new species (Hymenoptera: Apoidea: Crabronidae). *Zootaxa*, 1745 (1), 19–29.
- Melo, G.A.R. (1999) Phylogenetic relationships and classification of the major lineages of Apoidea (Hymenoptera), with emphasis on crabronid wasps. *Scientific Papers. Natural History Museum, The University of Kansas*, 14, 1–55. <https://doi.org/10.5962/bhl.title.4053>
- Olszewski, P., Ljubomirov, T., Wiśniowski, B., Kowalczyk, J.K. & Krzyżyński, M. (2016) New records of the genus *Diodontus* Curtis, 1834 (Hymenoptera: Crabronidae) from Bulgaria, Montenegro and Poland, with a key to Central and Eastern European species. *Zootaxa*, 4061 (2), 164–172. <http://dx.doi.org/10.11646/zootaxa.4061.2.6>
- Pulawski, W.J. (2007) The wasp genus *Tachysphex* Kohl, 1883, of Sahara, Sub-Saharan Africa, the Arabian Peninsula, and Madagascar (Hymenoptera: Apoidea: Crabronidae). *Proceedings of the California Academy of Sciences*, Series 4, 58 (Supplement 1), 1–698.
- Pulawski, W.J. (2020) Catalog of Sphecidae. San Francisco, CA: California Academy of Sciences. Available from: <https://www.calacademy.org/scientists/projects/catalog-of-sphecidae/> [Accessed 20th March 2020].
- Rezaei, S. & Fallahzadeh, M. (2015) New data on the digger wasps (Hymenoptera, Apoidea: Crabronidae) in Southern Iran. *Far Eastern Entomologist*, 303, 1–18.
- Sadeghi, M., Fallahzadeh, M., Ostovan, H., Ljubomirov, T. & Hesami S. (2016) New records of nine crabronid wasps (Hymenoptera: Crabronidae) from Iran. In: Talaei-Hassanlou, R. (ed.) *Proceedings of 22nd Iranian Plan Protection Congress*, 27–30 August, Karaj, College of Agriculture and Natural Resources, University of Tehran, Karaj, Iran, p. 431.
- Sadeghi, M., Fallahzadeh, M., Ostovan, H., Ljubomirov, T. & Hesami S. (2018a) New records of the digger wasps (Hymenoptera: Crabronidae) from Iran. *Far Eastern Entomologist*, 363, 21–24. <https://doi.org/10.25221/fee.363.4>
- Sadeghi, M., Fallahzadeh, M., Ostovan, H., Ljubomirov, T. & Hesami S. (2018b) Additions to the knowledge of the digger wasp (Hymenoptera: Spheciformes: Crabronidae) fauna in the Fars Province of Iran. *Journal of Insect Biodiversity and Systematics* 4, 261–279.
- Sadeghi, M., Fallahzadeh, M., Ostovan, H., Ljubomirov, T. & Hesami S. (2019) Revised checklist of the genus *Cerceris* (Hymenoptera: Crabronidae) of Iran. *Far Eastern Entomologist* 395, 14–22. <https://doi.org/10.25221/fee.395.3>
- Sakenin, H., Samin, N., Bagriacik, N. & Rastegar, J. (2011) A study of the Sphecidae (Hymenoptera) from various regions of Iran. *Calodema*, 161, 1–4.
- Samin, N., Nemkov, P. & Bagriacik, N. (2018) New records of Crabronidae and Sphecidae (Hymenoptera: Apoidea) from Iran. *Calodema*, 651, 1–4.
- Sann, M., Niehuis, O., Peters, R.S., Mayer, C., Kozlov, A., Podsiadlowski, L., Bank, S., Meusemann, K., Misof, B., Bleidorn, C. & Ohl, M. (2018) Phylogenomic analysis of Apoidea sheds new light on the sister group of bees. *BMC Evolutionary Biology*, 18, 1–15. <https://doi.org/10.1186/s12862-018-1155-8>
- Schmid-Egger, C. (2004) Revision of *Bembecinus* (Hymenoptera, Crabronidae) of the Palaearctic region. *Notes fauniques de Gembloux*, 54, 3–69.
- Schmid-Egger, C. (2014) Order Hymenoptera, families Crabronidae and Sphecidae. Further records and the description of new species, In: van Harten, A. (ed.) *Arthropod Fauna of UAE*. Vol. 5. Al Amal Printing Press. United Arab Emirates, Abu Dhabi, 744 pp.
- Schmid-Egger, C. (2019) Review of the genus *Ammatomus* A. Costa, 1859 (Hymenoptera, Crabronidae) from the Palearctic region with description of four new species. *Linzer Biologische Beiträge* 51, 437–457.
- Schmid-Egger, C., Fallahzadeh, M. & Khosroabadi, M. (2016) A new species of *Holotachysphex* de Beaumont, 1940 (Hymenoptera: Apoidea, Crabronidae) from Iran with identification key to species. *Zootaxa*, 4169 (1), 187–193. <https://doi.org/10.11646/zootaxa.4169.1.11>

یافته‌های جدید از زنبورهای حفار خانواده‌های Philanthidae و Psenidae و Pemphredonidae از ایران (Hymenoptera: Apoidea)

شهره رضایی^۱، مجید فلاح زاده^{۲*}، ابوفضل دوستی^۱ و نازیلا سقایی^۲

۱ گروه حشره‌شناسی، واحد جهرم، دانشگاه آزاد اسلامی، جهرم، ایران.

۲ گروه حشره‌شناسی، واحد مرودشت، دانشگاه آزاد اسلامی، مرودشت، ایران.

* پست الکترونیکی نویسنده مسئول مکاتبه: mfalahm@yahoo.com

| تاریخ دریافت: ۱۴ فروردین ۱۳۹۹ | تاریخ پذیرش: ۴ اردیبهشت ۱۳۹۹ | تاریخ انتشار: ۲۳ اردیبهشت ۱۳۹۹ |

چکیده: فون زنبورهای حفار (Hymenoptera: Apoidea) در استان فارس مطالعه شد و نمونه‌هایی از سه خانواده Pemphredonidae (پنج گونه از سه جنس)، Psenidae (سه گونه از دو جنس) و Philanthidae (دو گونه از دو جنس) جمع‌آوری شد که در اینجا ارایه می‌شود. هشت گونه D. *Diodontus brevilabris* de Beaumont, 1967 اینجا ارایه می‌شود. برای فون زنبورهای ایران جدید هستند. علاوه بر این دو گونه Cerceris *spinifera haladai* K. Schmidt *variegatus* (Tournier, 1889) و *Psenulus schencki* (Schenck, 1857) *laevigatus* (Schenck, 1857) *punctipleuris* (Gussakovskij, 1937) برای اولین بار از استان فارس گزارش می‌شوند. اطلاعات در دسترس دار ارتباط با تاکسونومی و پراکنش گونه‌های جمع‌آوری شده ارایه شد.

واژگان کلیدی: بال غشاییان، فون، گزارش‌های جدید، ایران